Published in final edited form as:

Doc Ophthalmol. 2010 August ; 121(1): 29–35. doi:10.1007/s10633-010-9228-z.

Deactivation of the Rod Response in Retinopathy of Prematurity (ROP)

Ronald M. Hansen, Maureen E. Harris, Anne Moskowitz, and Anne B. Fulton

Department of Ophthalmology, Children's Hospital and Harvard Medical School, 300 Longwood Avenue, Boston, MA 02115

Abstract

It is known that retinopathy of prematurity (ROP) alters the activation of rod photoreceptors, but the effect of ROP on deactivation has not been investigated. We studied deactivation using a paired flash procedure in 22 subjects (12 infants and 10 older subjects) with a history of preterm birth and ROP. The amplitude of the rod isolated a-wave response to a flash presented 2 to 120 seconds after a test flash was measured, and the time at which it reached 50% of the single flash amplitude (t_{50}) was determined by linear interpolation. Deactivation results were compared to those in former pre-terms who never had ROP ($n=6$) and term born controls. In infants, t_{50} values of ROP subjects did not differ from those in subjects who never had ROP or term born controls. Among mature ROP subjects, eight of 12 had t_{50} values longer than any control subject. Prolonged deactivation in these mature ROP subjects may indicate lack of maturation of the deactivation process (t_{50}) or progressive compromise of retinal function with increasing age.

Keywords

Electroretinogram; Retinopathy of prematurity; Rod photoreceptor; Deactivation

Introduction

Abnormal activation of rod phototransduction has been reported in subjects with a history of retinopathy of prematurity (ROP) [1]. The more severe the ROP, the lower rod photoreceptor sensitivity, S_{ROD} [1]. S_{ROD} depends on the time constants of the biochemical steps from photon capture by rhodopsin to closure of the channels in the rod outer segment. Low S_{ROD} indicates slow kinetics of the molecular processes in the activation of phototransduction, possibly due to subtle alteration in the rod outer segment [2].

Following activation, the rod must deactivate in a timely manner to prepare for response to the next stimulus. The rod recovers by stepwise deactivation of rhodopsin, transducin, and phosphodiesterase in a series of biochemically complex processes [3–7]. In term born infants and infant rats, the kinetics of both activation and deactivation of the photoresponse are slower than in adults [5,8–14]. If activation in ROP is slower than in age-matched term born control subjects, is deactivation also abnormally slow? We used a paired flash technique to study deactivation in ROP subjects in infancy and at older ages.

Methods

Subjects

Twenty-eight subjects with a history of preterm birth were studied (Table 1). All had been monitored in the newborn intensive care nursery following schedules for examination that were modeled on those used in the multi-center ROP treatment trials [15–17]. Based on a review of the results of these examinations, the subjects were categorized by ROP history as either mild ROP or no ROP. Those with mild, untreated ROP (n=22) had Stage 2 or 3 in Zone II or III that resolved spontaneously [18]. The others (n=6) had serial examinations but never had ROP. Gestational age at birth ranged from 23 to 32 (median 27) weeks and birth weight from 550 to 1590 (median 910) grams. None had active ROP at the time of the ERG test. Thirteen subjects were tested as infants at median age 10 (range 7 to 18) weeks post-term. Fifteen older subjects were tested at median age 13 (range 11 to 21) years. The parameters of rod activation in these preterm subjects have been reported [1]. Results from term born 10 week old infants (n=15) and mature subjects (n=10) provide control data for comparison. Data from the term born infants and most of the mature control subjects (8 of 10) were reported previously [11].

This study conformed to the tenets of the Declaration of Helsinki and was approved by the Children's Hospital Committee on Clinical Investigation. Informed consent was obtained from the parents of the infants and children, assent from the older children, and consent from those 18 years and older.

ERG procedure

Parents stayed with infants and children throughout the procedure. One pupil was dilated with cyclopentolate 1% and the subject dark-adapted for 30 minutes. Then, under dim red light, proparacaine 0.5% was instilled and a bipolar Burian-Allen electrode was placed on the cornea. A ground electrode was placed on the skin over the ipsilateral mastoid.

Twenty four subjects were tested using a Compact 4 system (Nicolet, Madison, WI) and four subjects were tested using an Espion system (Diagnosys, Lowell, MA). Rod responses were recorded using a Wratten 47B filter ($\lambda < 510$ nm) in the Nicolet system and a 470 nm LED (half bandwidth 30 nm) in the Espion system. The gain was 1000 and bandpass in both systems was 0.5 to 1000 Hz. The digitization rate was 2,564 Hz for the Nicolet and 2,000 Hz for the Espion. In adult control subjects, rod activation and deactivation parameters do not differ between the Espion and Nicolet systems [13,19]. Therefore, data obtained using the two systems have been combined.

Deactivation of phototransduction in rods

The recovery of the rod's response to light was evaluated using a paired flash paradigm [5,9, 20]. After a +3.3 log scotopic troland second (scot td s) test flash, an equal intensity probe flash was presented at seven selected inter-stimulus intervals (ISI, 2 to 120 s). For each test-probe pair, the amplitude (R) of the rod-isolated a-wave response to the probe was measured at 8 ms. The response to the probe flash provides a measure of the circulating current in the rods [9, 10,20]. For each ISI, R was expressed as proportion of R_{MAX} , the amplitude of the a-wave to the probe alone. Linear interpolation was used to determine the time, t_{50} , at which a-wave amplitude was half R_{MAX} . Between each test-probe pair, 2 minutes in the dark was allowed. Control experiments in term born subjects showed that 2 minutes were sufficient for full recovery to the amplitude of the dark-adapted response [11]. To obtain the rod isolated response, the amplitude of the a-wave to a photopically matched red flash was subtracted from the response to the probe flash.

Rod ERG

Activation was also studied in all subjects. Responses to full-field, brief (<3 ms), blue stimuli ranging from +2.1 to +3.6 log scot td s were recorded. The rod photoresponse parameters (S_{ROD} and R_{ROD}) were calculated by fit of the Hood and Birch [21] formulation of the Lamb & Pugh model [22,23] to the a-waves. The equation is

$$R(i, t) = [1 - \exp\{-0.5 I S_{\text{ROD}}(t - t_d)^2\}] R_{\text{ROD}}$$

where I is the flash (scot td s), S_{ROD} a sensitivity parameter [$(\text{scot td})^{-1}\text{s}^{-3}$] that depends on the time constants of the steps in activation of phototransduction [23], R_{ROD} the saturated response amplitude (μV), and t_d a brief delay (ms). All three parameters, (S_{ROD} , R_{ROD} , and t_d) were free to vary [24].

Calibrations

Stimuli were measured with a detector and appropriate photopic or scotopic filter (IL 1700, International Light, Newburyport, MA) placed at the position of the subject's cornea. Retinal illuminance varies directly with area of the pupil and the transmissivity of the ocular media and inversely with the square of the posterior nodal distance [25]. We used the area of each infant's dilated pupil, published estimates of the ocular media density, and measures of the axial length of the eye. In summary, equal intensity stimuli produced approximately equal retinal illuminance in both infants and control subject [1,11,26–30]. The maximum intensity stimulus produced a retinal illuminance of approximately +3.3 log scot td s in both infants and mature subjects.

Statistical analyses

For each age group, the Mann Whitney test was used to compare the deactivation parameter (t_{50}) of the ROP and no ROP subjects. Spearman rank order correlation was used to evaluate the relationship between t_{50} and S_{ROD} . The level of significance was $p < 0.01$.

Results

On records such as those in Fig. 1 (left panel), a-wave amplitude was measured 8 ms after the stimulus. For each test-probe flash pair, R/R_{MAX} was plotted as a function of inter-stimulus interval (right panel).

The t_{50} values in infants and mature subjects are plotted in Fig. 2. Table 2 summarizes the median deactivation (t_{50}) and activation (S_{ROD}) parameters of the former preterms and control subjects. The t_{50} values in mature ROP subjects were significantly longer (Mann Whitney $U = 0$, $p < 0.01$) than in those who never had ROP. In eight of the 12 mature ROP subjects, t_{50} was longer than in any of the mature controls. The t_{50} values in ROP infants did not differ (Mann Whitney $U = 15$, ns) from those in prematurely born infants who never had ROP and were similar to those previously reported for term born infants. In both age groups, the t_{50} values of the former preterm subjects who never had ROP were similar to those of the control subjects. The t_{50} values in the mature controls are similar to those in the two subjects tested by Friedburg *et al.* [10] who also used equal intensity test and probe flashes ($\sim +3$ log td s).

In term born subjects, higher values of S_{ROD} are associated with shorter t_{50} values [11]. However, this is not the case in the ROP subjects ($p = -0.262$; ns). In Fig. 3, deactivation (t_{50}) and activation (S_{ROD}) values for each mature ROP subject are shown. In the majority (5 of 7) with normal S_{ROD} ($> 8.2 \text{ isoms}^{-1}\text{s}^{-2}$), deactivation was slower than normal ($t_{50} > 6.2 \text{ s}$);

t_{50} was also abnormal in three of the five mature ROP subjects who had low S_{ROD} . Two had normal t_{50} and low S_{ROD} . Both parameters were normal in only two. Thus, a mismatch occurred in seven of the 12 mature ROP subjects. In contrast, all ten infant ROP subjects had normal t_{50} for age and S_{ROD} was normal in six. For preterm subjects, neither t_{50} nor S_{ROD} varied significantly with either gestational age at birth or birth weight.

Discussion

The results are evidence that the kinetics of deactivation (t_{50}) in the majority of the mature ROP subjects were abnormally slow, even in cases with normal kinetics of activation (S_{ROD}). Furthermore, two with normal t_{50} had low S_{ROD} . In normal development, t_{50} and S_{ROD} are correlated; higher S_{ROD} is associated with more rapid recovery, that is, shorter t_{50} . This is not necessarily the case in disease. For example, re-analysis of data in patients with Smith-Lemli-Opitz Syndrome and mitochondrial disease [31,32] indicated no relationship between t_{50} and S_{ROD} . In the present sample of mature ROP subjects, we found no significant correlation. These data raise the possibility that the ROP rod's capacity to recover from light stimulation did not mature in eight of the subjects with mild ROP, even among those in whom S_{ROD} was normal (Figure 3). Alternatively, the deactivation of the rods in some older ROP subjects may have suffered progressive compromise; long t_{50} is associated with photoreceptor degeneration [9]. S_{ROD} was abnormally low in five subjects due to either lack of maturation or progressive loss. The data do not distinguish between these two explanations. There is evidence of progressive compromise of rod driven post-receptor retinal function in subjects with a history of severe ROP [1,33]. All ROP infants had deactivation kinetics that were normal for age; their t_{50} values were within the range of age matched controls.

In healthy control subjects, activation and deactivation are coupled; higher S_{ROD} values are associated with lower t_{50} values [11,34]. Studies of deactivation in healthy infants [11,34] and adults [5,9,11,34,35] have shown that the brighter the stimuli, the slower the recovery. In healthy infant and adult rat retina, t_{50} varies with the proportion of rhodopsin isomerized [8]. A similar relationship appears to pertain to deactivation in human rods. We have measured [36] the length of peripheral human rod outer segments (ROS) shown in Hendrickson [37]. The peripheral ROS length in the 5 day old infant was 43% of that in the adult. Assuming that there are 7×10^7 molecules of rhodopsin in the adult ROS [23], there are $\sim 3 \times 10^7$ molecules of rhodopsin in the 5 day old ROS. If the axial density of rhodopsin in the ROS is proportional to the rhodopsin content of the retina, the ratio of infant to adult isomerizations is proportional to $(1-10^{-D_{\text{infant}}})/(1-10^{-D_{\text{adult}}})$. The rhodopsin content of the retina in a 10 week old infant is $\sim 68\%$ of that in the adult retina [38]. Therefore, D_{infant} is $\sim 0.68 \times D_{\text{adult}}$ and 1 scot td would isomerize 6.6 molecules of rhodopsin in the infant retina [11]. The $+3.3$ log scot td s flash would isomerize $\sim 13,200$ molecules of rhodopsin (0.044%) in the infant retina compared to $\sim 17,000$ (0.024%) in the adult. Thus, isomerization of a higher proportion of rhodopsin in the infants is consistent with the longer t_{50} values found in infants. This explanation may pertain to the ROP infants. The amount of rhodopsin extractable from the whole retina did not differ between ROP and control rats [2].

The stimuli used in the present study bleach only a small fraction ($<1\%$) of the rhodopsin in the ROS. The double flash method we have used [10] was designed to evaluate the time course of stepwise deactivation of rhodopsin, transducin, and phosphodiesterase. The rate limiting step in recovery of the photoresponse (deactivation) is hydrolysis of guanosine triphosphate (GTP) which is catalyzed by a membrane-associated multi-molecular complex [7]. Slow diffusion of the activation and deactivation proteins in ROS could account for long t_{50} and low S_{ROD} values found in three of the mature ROP subjects. However, a mismatch (normal activation and abnormal deactivation, or abnormal activation and normal deactivation) was found in seven of the mature ROP subjects. Normal kinetics of activation (normal S_{ROD}) is

evidence of normal molecular movements in the disc membrane and in the cytosol leading to closure of the channels in the outer segment membrane [39]. As for the kinetics of deactivation, possibly the re-supply of retinoid is sluggish. Alternatively, those molecules dedicated to recovery may be selectively affected in some ROP subjects. These molecular issues will be more readily approached in rats with oxygen-induced retinopathy [40–44], which is known to model key features of ROP [45]. To our knowledge, deactivation of phototransduction has yet to be studied in rat models of ROP.

Further study of the kinetics of deactivation in mature ROP subjects using a range of stimulus levels, along with measurement of rhodopsin density and kinetics of regeneration [46,47] and dark adaptation, might help explain the abnormal results. The final dark adapted thresholds of some ROP subjects are elevated [48]; the kinetics of dark adaptation and rhodopsin density and regeneration have not been studied.

Acknowledgments

This work was supported by National Eye Institute Grant R01 EY010597.

References

1. Fulton AB, Hansen RM, Moskowitz A, Akula JD. The neurovascular retina in retinopathy of prematurity. *Prog Retin Eye Res* 2009;28(6):452–82. corrigendum, 2010;29:94. [PubMed: 19563909]
2. Fulton AB, Reynaud X, Hansen RM, Lemere CA, Parker C, Williams TP. Rod photoreceptors in infant rats with a history of oxygen exposure. *Invest Ophthalmol Vis Sci* 1999;40(1):168–74. [PubMed: 9888440]
3. Arshavsky VY. Rhodopsin phosphorylation: from terminating single photon responses to photoreceptor dark adaptation. *Trends Neurosci* 2002;25(3):124–6. [PubMed: 11852136]
4. Burns ME, Baylor DA. Activation, deactivation, and adaptation in vertebrate photoreceptor cells. *Annu Rev Neurosci* 2001;24:779–805. [PubMed: 11520918]
5. Pepperberg DR, Birch DG, Hofmann KP, Hood DC. Recovery kinetics of human rod phototransduction inferred from the two-branched a-wave saturation function. *J Opt Soc Am A* 1996;13:586–600.
6. Pugh, EN., Jr; Lamb, TD. Phototransduction in vertebrate rods and cones: Molecular mechanisms of amplification, recovery and light adaptation. In: Stavenga, DG.; de Grip, WJ.; Pugh, EN., Jr, editors. *Handbook of biological physics. Volume 3. Molecular mechanisms of visual transduction*. Elsevier Science; 2000. p. 183–255.
7. Burns ME, Pugh EN Jr. RGS9 concentration matters in rod phototransduction. *Biophys J* 2009;97(6):1538–47. [PubMed: 19751658]
8. Fulton AB, Hansen RM. Recovery of the rod photoresponse in infant rats. *Vision Res* 2003;43(28):3081–5. [PubMed: 14611945]
9. Birch DG, Hood DC, Nusinowitz S, Pepperberg DR. Abnormal activation and inactivation mechanisms of rod transduction in patients with autosomal dominant retinitis pigmentosa and the pro-23-his mutation. *Invest Ophthalmol Vis Sci* 1995;36:1603–1614. [PubMed: 7601641]
10. Friedburg C, Thomas MM, Lamb TD. Time course of the flash response of dark- and light-adapted human rod photoreceptors derived from the electroretinogram. *J Physiol* 2001;534(Pt 1):217–42. [PubMed: 11433004]
11. Hansen RM, Fulton AB. Recovery of the rod photoresponse in infants. *Invest Ophthalmol Vis Sci* 2005;46(2):764–8. [PubMed: 15671311]
12. Pepperberg DR, Cornwall MC, Kahlert M, et al. Light-dependent delay in the falling phase of the retinal rod photoresponse. *Vis Neurosci* 1992;8:9–18. [PubMed: 1739680]
13. Fulton AB, Hansen RM. The development of scotopic sensitivity. *Invest Ophthalmol Vis Sci* 2000;41(6):1588–96. [PubMed: 10798680]
14. Fulton AB, Hansen RM, Findl O. The development of the rod photoresponse from dark-adapted rats. *Invest Ophthalmol Vis Sci* 1995;36(6):1038–45. [PubMed: 7730013]

15. Early Treatment For Retinopathy Of Prematurity Cooperative Group. Revised indications for the treatment of retinopathy of prematurity: results of the early treatment for retinopathy of prematurity randomized trial. *Arch Ophthalmol* 2003;121(12):1684–1694. [PubMed: 14662586]
16. Committee for the Classification of Retinopathy of Prematurity. An international classification of retinopathy of prematurity. *Arch Ophthalmol* 1984;102:1130–1134. [PubMed: 6547831]
17. Cryotherapy for Retinopathy of Prematurity Cooperative Group. Multicenter trial of cryotherapy for retinopathy of prematurity: Three-month outcome. *Arch Ophthalmol* 1988;106:471–479. [PubMed: 2895630]
18. The international classification of Retinopathy of Prematurity revisited. *Arch Ophthalmol* 2005;123:991–999. [PubMed: 16009843]
19. Hansen RM, Fulton AB. Development of the cone ERG in infants. *Invest Ophthalmol Vis Sci* 2005;46(9):3458–62. PMC1467576. [PubMed: 16123452]
20. Lyubarsky AL, Pugh EN Jr. Recovery phase of the murine rod photoresponse reconstructed from electroretinographic recordings. *J Neurosci* 1996;16:563–571. [PubMed: 8551340]
21. Hood DC, Birch DG. Rod phototransduction in retinitis pigmentosa: estimation and interpretation of parameters derived from the rod a-wave. *Invest Ophthalmol Vis Sci* 1994;35(7):2948–61. [PubMed: 8206712]
22. Lamb TD, Pugh EN Jr. A quantitative account of the activation steps involved in phototransduction in amphibian photoreceptors. *J Physiol* 1992;449:719–58. [PubMed: 1326052]
23. Pugh EN Jr, Lamb TD. Amplification and kinetics of the activation steps in phototransduction. *Biochim Biophys Acta* 1993;1141(2–3):111–49. [PubMed: 8382952]
24. Fulton AB, Hansen RM, Petersen RA, Vanderveen DK. The rod photoreceptors in retinopathy of prematurity: an electroretinographic study. *Arch Ophthalmol* 2001;119(4):499–505. [PubMed: 11296015]
25. Pugh, EN, Jr. *Stevens Handbook of Experimental Psychology*. Wiley; New Work: 1988. Vision: Physical and retinal physiology.
26. Brown AM, Dobson V, Maier J. Visual acuity of human infants at scotopic, mesopic and photopic luminances. *Vision Res* 1987;27:1845–1858. [PubMed: 3445474]
27. Hansen, RM.; Fulton, AB. Development of scotopic retinal sensitivity. In: Simons, K., editor. *Early Visual Development, Normal and Abnormal*. Oxford University Press; New York: 1993. p. 130–142.
28. Hansen RM, Fulton AB, Harris SJ. Background adaptation in human infants. *Vision Res* 1986;26(5):771–9. [PubMed: 3750858]
29. Mactier H, Maroo S, Bradnam M, Hamilton R. Ocular biometry in preterm infants: implications for estimation of retinal illuminance. *Invest Ophthalmol Vis Sci* 2008;49(1):453–7. [PubMed: 18172125]
30. Malcolm CA, Hamilton R, McCulloch DL, Montgomery C, Weaver LT. Scotopic electroretinogram in term infants born of mothers supplemented with docosahexaenoic acid during pregnancy. *Invest Ophthalmol Vis Sci* 2003;44(8):3685–91. [PubMed: 12882824]
31. Cooper LL, Hansen RM, Darras BT, et al. Rod photoreceptor function in children with mitochondrial disorders. *Arch Ophthalmol* 2002;120(8):1055–62. [PubMed: 12149059]
32. Elias ER, Hansen RM, Irons M, Quinn NB, Fulton AB. Rod photoreceptor responses in children with Smith-Lemli-Opitz syndrome. *Arch Ophthalmol* 2003;121(12):1738–43. [PubMed: 14662594]
33. Harris ME, Hansen RM, Moskowitz A, Fulton AB. Long term effects of retinopathy of prematurity (ROP) on rod and rod-driven function. *Investigative Ophthalmology and Visual Science*. 2009 ARVO E-Abstract 5310.
34. Pepperberg DP, Birch DG, Hood DC. Electroretinographic determination of human rod flash response *in vivo*. *Methods Enzymol* 2000;316:202–223. [PubMed: 10800677]
35. Pepperberg DR, Birch DG, Hood DC. Photoresponse of human rods *in vivo* derived from paired flash electroretinograms. *Vis Neurosci* 1997;14:73–82. [PubMed: 9057270]
36. Hansen RM, Fulton AB. The course of maturation of rod-mediated visual thresholds in infants. *Invest Ophthalmol Vis Sci* 1999;40(8):1883–6. [PubMed: 10393066]

37. Hendrickson, AE. The morphologic development of human and monkey retina. In: Albert, DM.; Jakobiec, FA., editors. Principles and Practice of Ophthalmology: Basic Sciences. WB Saunders Co; Philadelphia: 1994. p. 561-577.
38. Fulton AB, Dodge J, Hansen RM, Williams TP. The rhodopsin content of human eyes. Invest Ophthalmol Vis Sci 1999;40(8):1878–83. [PubMed: 10393065]
39. Lamb TD, Pugh EN Jr. Phototransduction, dark adaptation, and rhodopsin regeneration the proctor lecture. Invest Ophthalmol Vis Sci 2006;47(12):5137–52. [PubMed: 17122096]
40. Akula JD, Favazza TL, Mocko JA, et al. The anatomy of the rat eye with oxygen-induced retinopathy. Doc Ophthalmol 2010;120(1):41–50. [PubMed: 19820974]
41. Barnett JM, Yanni SE, Penn JS. The development of the rat model of retinopathy of prematurity. Doc Ophthalmol 2010;120(1):3–12. [PubMed: 19639356]
42. Berkowitz BA, Roberts R. Evidence for a critical role of panretinal pathophysiology in experimental ROP. Doc Ophthalmol 2010;120(1):13–24. [PubMed: 19506927]
43. Fletcher EL, Downie LE, Hatzopoulos K, et al. The significance of neuronal and glial cell changes in the rat retina during oxygen-induced retinopathy. Doc Ophthalmol 2010;120(1):67–86. [PubMed: 19763649]
44. Hartnett ME. The effects of oxygen stresses on the development of features of severe retinopathy of prematurity: knowledge from the 50/10 OIR model. Doc Ophthalmol 2010;120(1):25–39. [PubMed: 19639355]
45. Fulton AB, Akula JD, Mocko JA, et al. Retinal degenerative and hypoxic ischemic disease. Doc Ophthalmol 2009;118(1):55–61. [PubMed: 18483822]
46. Alpern M, Pugh EJ. The density and photosensitivity of human rhodopsin in the living retina. Journal of Physiology 1974;237:341–70. [PubMed: 4825455]
47. Fulton A, Hansen R. The relation of rhodopsin and scotopic retinal sensitivity in sector retinitis pigmentosa. Am J Ophthalmol 1988;105:132–140. [PubMed: 3341429]
48. Hansen RM, Fulton AB. Background adaptation in children with a history of mild retinopathy of prematurity. Invest Ophthalmol Vis Sci 2000;41(1):320–4. [PubMed: 10634637]

Fig. 1.

Sample results from a 14 year old subject with a history of mild ROP and t_{50} value (10 s) close to the ROP median. Rod isolated ERG a-wave responses are shown in the left panel; the responses to the single flash and to the probe flash at the indicated inter-stimulus intervals (ISI) are shown. In the right panel, a-wave amplitude (R) expressed as a proportion of the response to the single flash (R_{MAX}) is plotted as a function of ISI. The ISI (arrow) that would produce a half maximum response was determined by linear interpolation.

Fig. 2.

Values of t_{50} and S_{ROD} . Upper panel: Values of t_{50} are plotted for the mild ROP subjects (filled circles), the no ROP subjects (filled triangles), and term born control subjects (open circles). The horizontal bars indicate the median for each group. Lower panel: Values of S_{ROD} are plotted for the same subjects; all features are as described for the upper panel. S_{ROD} values for these subjects were reported previously [1].

Fig. 3.

Deactivation parameter (t_{50}) versus activation parameter (S_{ROD}). In both panels, the shaded bands indicate the range of t_{50} and S_{ROD} values in control subjects; the cross hatched area represents the locus of control data [11]. Data from infant ROP subjects ($n=10$) are plotted in the upper panel and from mature ROP subjects ($n=12$) in the lower panel. The same scales are used in both panels to facilitate comparison across age groups.

Table 1

Subjects with a history of preterm birth [median (range)]

	N	Test age	Gestational age at birth (weeks)	Birth weight (grams)
Infants				
ROP	10	73 (68–81) days	28 (26–31)	970 (626–1170)
No ROP	3	73 (52–131)	29 (25–30)	1230 (740–1360)
Mature				
ROP	12	13 (11–21) years	26 (23–28)	815 (550–1077)
No ROP	3	13 (10–18)	31 (26–32)	1510 (1430–1590)

Table 2Deactivation (t_{50}) and activation (S_{ROD}) parameters [median (range)]

	t_{50} (s)		
	ROP	No ROP	Control
Infant	11.9 (6.9 – 14.7)	10.4 (10.0 – 12.7)	9.3 (7.0 – 15.0)
Mature	7.5 (5.0 – 15.0)	2.9 (2.8 – 4.2)	4.2 (2.6 – 6.2)
	S_{ROD} [(scot td) $^{-1}$ s $^{-3}$]		
	ROP	No ROP	Control
Infant	27.6 (12.4 – 52.5)	39.0 (33.2 – 51.8)	39.9 (22.1 – 72.8)
Mature	77.9 (40.7 – 93.8)	102.7 (98.6 – 106.1)	86.1 (69.7 – 97.8)