Published in final edited form as:

J Steroid Biochem Mol Biol. 2010 July ; 121(1-2): 338–342. doi:10.1016/j.jsbmb.2010.03.046.

Epigenetic silencing of CYP24 in the tumor microenvironment

Candace S. Johnson^{a,*}, Ivy Chung^{b,**}, and Donald L. Trump^b

^a Department of Pharmacology & Therapeutics, Roswell Park Cancer Institute, Buffalo, NY, USA

^b Department of Medicine, Roswell Park Cancer Institute, Elm and Carlton Streets, Buffalo, NY, 14263, USA

Abstract

Calcitriol (1,25 dihydroxycholecalciferol) has significant antitumor activity *in vitro* and *in vivo* in a number of tumor model systems. We developed a system for isolation of fresh endothelial cells from tumors and Matrigel environments which demonstrate that CYP24, the catabolic enzyme involved in vitamin D signaling, is epigenetically silenced selectively in tumor-derived endothelial cells (TDEC). TDEC maintain phenotypic characteristics which are distinct from endothelial cells isolated from normal tissues and from Matrigel plugs (MDEC). In TDEC, calcitriol induces G₀/G₁ arrest, modulates p27 and p21, and induces apoptotic cell death and decreases P-Erk and P-Akt. In contrast, endothelial cells isolated from normal tissues and MDEC are unresponsive to calcitriol-mediated anti-proliferative effects despite intact signaling through the vitamin D receptor (VDR). In TDEC, which is sensitive to calcitriol, the CYP24 promoter is hypermethylated in two CpG island regions located at the 5' end; this hypermethylation may contribute to gene silencing of CYP24. The extent of methylation in these two regions is significantly less in MDEC. Lastly, treatment of TDEC with a DNA methyltransferase inhibitor restores calcitriol-mediated induction of CYP24 and resistance to calcitriol. These data suggest that epigenetic silencing of CYP24 modulates cellular responses to calcitriol.

Keywords

calcitriol; vitamin D; tumor derived endothelial cells; matrigel-derived endothelial cells; CYP 24

1. Introduction

Studies in our laboratory demonstrate that calcitriol (vitamin D or 1,25 dihydroxycholecalciferol), a central factor in bone and mineral metabolism [1,2], has significant antitumor activity *in vitro* and *in vivo* in a number of murine syngeneic and human xenograft tumor model systems [3–6]. Calcitriol induces cell cycle arrest and related kinases, decreases survival signaling molecules and induces modulators of apoptosis in various tumor types [4–10]. Clinically in phase I studies, we have determined the MTD of calcitriol alone and in combination with a number of cytotoxic agents and/or glucocorticoids, where there is synergy [11–20] and we have completed a phase II trial on androgen-independent prostate cancer where the combination of dexamethasone and

*Corresponding author: Tel.: +1 716 845 8300; Fax: +1 716 845 1258. Candace.Johnson@RoswellPark.org.

**Current address: Vascular Biology Program Children's Hospital, Harvard Medical School, Boston, MA 02115

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

calcitriol resulted in an enhanced anti-tumor effect. Induction of CYP24, the enzyme primarily responsible for calcitriol catabolism [21], may be a factor in bioavailability, calcitriol exposure and the anti-proliferative activity pre-clinically and clinically. In addition, the endothelial cells in tumors are sensitive to calcitriol and uniquely modulate CYP24 expression through epigenetic changes [22].

Epigenetic events affect gene expression without alteration in DNA gene sequence and lead to transcriptional gene silencing and inactivation of tumor suppressor genes in human cancer [23]. While many studies document epigenetic changes in tumor cells, only limited data support a role for epigenetic changes in the “normal” cells found in the tumor microenvironment. Epigenetic changes are found in the stromal fibroblasts from normal human breast tissue and breast carcinoma [24,25] and in the tumor stroma and endothelium of localized human prostate cancer [26,27]. Calcitriol also inhibits proliferation of endothelial cells and can inhibit angiogenesis in a number of tumor model systems [28–30]. We have established a method for the isolation of fresh, tumor-derived endothelial cells (TDEC) that maintain phenotypic characteristics which are distinct from endothelial cells isolated from normal tissues [31] and from Matrigel plugs (MDEC) [32]. These endothelial cell populations differ in their response to calcitriol and in their expression of CYP24. We discuss here the epigenetic silencing of CYP24 in these cells and the potential role on the tumor microenvironment in these activities. These unique differences in the epigenetic silencing of calcitriol-induced CYP24 gene expression in endothelial cells from the tumor microenvironment directly impact on calcitriol-mediated signaling pathways and ultimately on therapeutic application.

2. Results / Discussion

2.1. Isolation of TDEC and MDEC

Using an antibody to PECAM or platelet cell adhesion molecule, a positive marker for endothelial cells [31,32], tumor cells are harvested, dissociated, labeled and sorted for PECAM-positive cells. In a dissociated SCC or PC-3 tumor, the sorted population is >95% positive for PECAM and yields $1\text{--}1.5 \times 10^6$ cells from $2\text{--}4 \times 10^7$ dissociated tumor cells. Cells highly fluorescent for PECAM (positive sort) represent 4–7% of the total population of cells from the tumor and have a doubling time of 50 hr as compared to 12–20 hr for most tumor cell lines. These PECAM-positive cells are also positive for the uptake of acetylated low-density lipoprotein (Dil-ac-LDL), factor VIII and demonstrate endothelial-like characteristics by electron microscopy (presence of Weibel-Palade bodies). In addition, cell survival throughout this entire procedure is >90% and PECAM-positive endothelial cells are available for analysis within 6–8 hr of tumor harvest. Cells are utilized fresh or after only 1–2 passages. Cultures maintain a number of phenotype markers through 10 passages or 5–6 weeks in culture. For the appropriate “normal” endothelial cell control, microvascular endothelial cells were isolated from Matrigel plugs that were implanted subcutaneously into mice [32]. In selecting a “normal” EC, it was important to create a situation similar to a tumor without the tumor present to isolate neovascular endothelial cells that were recruited toward an antigenic stimulus. We developed a model to create a “normal” microenvironment where endothelial cells would be recruited, but where the tumor cells would not be present [22,31–34]. Endothelial cells were cultured and propagated in cell culture dishes coated with 1% gelatin. When compared to TDEC, MDEC showed distinct morphologic differences of long, branching and cytoplasmic projections. The doubling times of TDEC and MDEC were 24–28 hours and 32–36 hours, respectively.

2.2 Comparison of VDR between TDEC and MDEC

Using Western blot analysis, both TDEC and MDEC expressed similar basal levels of VDR protein and responded to 10 nM calcitriol by increasing VDR protein expression in a time dependent manner [32]. Although MDEC had higher basal VDR expression compared to TDEC, treatment with calcitriol resulted in increase of VDR protein expression in both cell types across time. An additional middle band was observed in both cell types following calcitriol treatment. These doublet or triplet bands may represent different phosphorylated forms of VDR as indicated in a dephosphorylation assay using λ phosphatase. The more slowly migrating forms of VDR were either reduced or converted to a faster migrating form upon treatment with λ phosphatase, indicating that phosphorylation of VDR is a post-translational process.

To determine whether there is any difference in the VDR mRNA level between the two cells, we amplified the whole coding region of VDR sequence using semi-quantitative RT-PCR and subjected the PCR product for sequencing. The PCR products from both TDEC and MDEC appeared similar in size and quantity. There were no mutations found in either sequence when compared to the VDR from the mouse database. However, dose-response saturation plot and Scatchard plot analysis suggested that the kinetics of receptor-ligand binding were different in these cells. The dose-response saturation plot, which represents only specific binding, approached saturation at 31 ± 0.59 fmol/mg and 24 ± 3.5 fmol/mg ($P = 0.001$), of [^3H]1,25- D_3 for TDEC and MDEC, respectively. These results indicate that the vitamin D receptor level (as measured by maximum 1,25- D_3 binding capacity per milligram of protein) is higher in TDEC than in MDEC. By Scatchard plot analysis, linear regression analysis demonstrated a single class of specific, high affinity receptors with lower equilibrium binding disassociation constant (K_d) in TDEC (0.26 ± 0.001 nM) than in MDEC (0.65 ± 0.000005 nM) ($P = 0.0016$). Using 24-hydroxylase promoter-luciferase reporter assay, the VDR-RXR heterodimer complex in TDEC and MDEC translocated into the nucleus and transactivated the reporter gene in a dose dependent manner [32]. With increasing dose of the reporter gene, the endogenous VDR from both cells transactivated the exogenous DNA in a similar rate, when treated with 10 nM calcitriol. However, at a constant dose of the reporter gene and varying concentrations of calcitriol, VDR in TDEC transactivated the DNA at a higher rate than in MDEC. Taken together, these findings indicate that the VDR signaling was intact and functional in both TDEC and MDEC, but more efficient ligand binding and transactivation occurred in TDEC than MDEC. Studies using other types of endothelial cells also demonstrate VDR expression upon treatment with calcitriol, indicating possible role of VDR in the biology of these cells [33–34].

2.3. Anti-proliferative effects of calcitriol on TDEC/MDEC

We studied the effects of calcitriol on the growth of MDEC and TDEC [32,36]. Cells were treated with varying doses of calcitriol for 48 hours and cell viability was determined using the MTT assay. Calcitriol, at nanomolar concentrations, inhibited cell growth of TDEC. However, the growth of the MDEC was only minimally affected by the treatment with calcitriol. The resistance to calcitriol's effect was also observed in other types of normal endothelial cells, including those isolated from mouse brain, lung and yolk sac. The effects of calcitriol on cell cycle distribution in TDEC and MDEC were assessed by PI staining where calcitriol induced cell cycle arrest in TDEC and not in MDEC. The protein expression of p27 was induced and p21 was decreased in calcitriol-treated TDEC but not in MDEC. Further effects on DNA synthesis were examined and found a reduction of BrdU incorporation in TDEC and no change in MDEC. Using annexin V/7-AAD staining, treatment with calcitriol resulted in a significant increase in total cell death in TDEC and not MDEC. Using Western blot analysis, treatment of calcitriol caused a significant reduction of P-Erk and P-Akt protein in TDEC and no change in total Erk and total Akt; no changes in

these proteins were seen in MDEC. Furthermore, a reduction in Bcl-2 and full length PARP and an increase in cleavage of caspase-3, all markers of apoptosis, were observed in TDEC treated with calcitriol but no such effects was seen on MDEC.

To evaluate whether calcitriol treatment *in vivo* has an impact on tumor vasculature, we quantified the mean vessel density in SCC tumor and Matrigel plugs removed from animals treated daily $\times 3$ with calcitriol (0.625 μ g/mouse), a dose that resulted in a significant anti-tumor effect [6–8]. Three areas were evaluated from each tissue and mean vessel density calculated based on the number of CD31 stained microvessels in each field. As shown in Figure 1A, a significant reduction was observed in the number of vessels after calcitriol treatment in the tumor with little effect observed in the Matrigel. To quantitate apoptosis, we utilized CD31 and activated caspase-3 double staining to identify endothelial cells undergoing apoptosis. After treatment with calcitriol, an increase was observed in the number of apoptotic cells in the tumor and not in the Matrigel plug (Figure 1B). Additional studies demonstrated that treatment with calcitriol resulted in an increase of vessel permeability at 8hrs post injection and a decrease in blood volume compared to control tumors as measured by MR imaging. These studies suggest that calcitriol targets the vasculature and results in significant endothelial cell death and ultimately collapse of the vasculature in the tumor.

2.4. Calcitriol-induced CYP24 expression in TDEC and MDEC

CYP24 or 25-hydroxyvitamin D-24-hydroxylase is the key enzyme responsible for initiating the vitamin D degradation pathway by directing the side chain metabolism of 25-hydroxylated vitamin D metabolites, including calcitriol [37]. This enzyme is expressed in kidney and most vitamin D target tissues [38]. CYP24 can be induced by calcitriol via positive transcriptional regulation because of the presence of two VDREs in the CYP24 promoter region. This mechanism provides negative feedback regulation for attenuation of excessive vitamin D effects within the tissues.

Both TDEC and MDEC activate VDR signaling pathway upon treatment of calcitriol [22,32]. However, treatment of 10 nM calcitriol at various time points induced CYP24 mRNA and protein expression in MDEC but not in TDEC. The induction of CYP24 mRNA in MDEC was observed as early as 3 hours and was sustained for over 48 hours post treatment. Similarly, using antibody specific to CYP24 (a gift from Cytochroma, Inc), we observed that CYP24 protein was expressed in a time dependent manner in MDEC while no CYP24 protein expression was observed in TDEC. Briefly, [^{3} H]-25-D $_3$ oxidative metabolites were extracted by liquid/liquid partition using tetrahydrofuran and ethyl acetate, dried and analyzed by high performance liquid chromatography (HPLC) on Zorbax SIL column (Agilent Technologies Inc, Palo Alto, CA) using hexane/isopropanol (94:6 v/v) as the mobile phase [22]. We further confirmed the enzymatic activity of CYP24 protein induced by calcitriol in both cell types using high performance liquid chromatography (HPLC). In MDEC, treatment of 10 nM calcitriol for 24 and 48 hours resulted in approximately 4- and 7-fold increase of CYP24 enzymatic activity, respectively when compared to the vehicle ($P < 0.005$); no significant induction of CYP24 activity was observed in calcitriol-treated TDEC [22].

Differential induction of CYP24 by calcitriol in MDEC and TDEC could be a mechanism accounting for the lack of response in MDEC to calcitriol-mediated growth inhibition. To test this hypothesis, we used a siRNA-specific for CYP24 to treat MDEC concurrent with calcitriol. Treatment of 10 nM calcitriol for 48 hours resulted in a maximum of 15% cell kill of MDEC. MDEC were transfected with siRNA-CYP24, siRNA-NS or untransfected (mock) for 24 hrs before addition of 10 nM calcitriol for 48 hours. When MDEC was treated with siRNA-CYP24, there was a down-regulation of protein expression and enzymatic

activity and there was a 43% growth inhibition in MDEC with CYP24 siRNA as compared to 15% in the cells treated with siRNA-NS or mock control, indicating that CYP24 protects MDEC from the growth inhibitory effects of calcitriol.

2.5. Transcription gene silencing of CYP24 in TDEC

Since both TDEC and MDEC were isolated from the same strain background of mice (C3H/HeJ) but different microenvironments (tumor vs. non-tumor), the observation that calcitriol failed to induce CYP24 in TDEC was intriguing. CYP24 is a well known vitamin D target gene, which is ubiquitously expressed to regulate calcitriol homeostasis in the body [21]. The cytochrome P450 (CYP) family of enzymes is involved in numerous cellular functions including metabolism of eicosanoids, biosynthesis of cholesterol, synthesis and degradation of biogenic amines, hydroxylation of retinoic acid as well as the metabolism of vitamin D [39]. Using microarray (Figure 2A), we compared the overall expression of CYPs in TDEC and MDEC and found suppression across a number of CYPs, except CYP2C44, a CYP that metabolizes arachidonic acid [40]. These results were confirmed using qRT-PCR. Because CYP24 promoter hypermethylation in osteoblastic cells has been indicated in a previous report [41], we examined the promoter region of CYP24 gene in TDEC for CpG island hypermethylation and epigenetic gene silencing as a possible molecular mechanism for these effects. We characterized the methylation status of two proximal CpG islands in the 5' region of CYP24, region 1 (−1131 to −241) and region 2 (−214 to +800) in TDEC and MDEC (Figure 1B). Methylation-specific PCR analysis showed hypermethylation of region 1 in TDEC but not in MDEC. However, region 2 in both cells was found to be hypermethylated. Bisulfite sequencing from both cells revealed that the CpG dinucleotides found in the amplified sequence of region 1 and 2 were significantly hypermethylated in TDEC when compared to MDEC and these findings were also confirmed by MassArray, a quantitative method of DNA methylation analysis (Figure 1C). Whether VDREs at the distal promoter region are also heavily methylated in order for the CYP24 gene to be transcriptionally silenced, is yet to be determined.

Tumor-derived endothelial cells (TDEC) are in close contact with the tumor cells within the tumor microenvironment. Therefore, the extent of methylation for CpG islands in regions 1 and 2 was also determined for SCC (squamous cell carcinoma), RIF (radiation-induced fibrosarcoma) and TRAMPC2 (prostate) tumor cell lines and then compared to TDEC isolated from the same tumor type [22]. Both SCC and TDEC/SCC were equally hypermethylated (83/84% vs 79/87%, regions 1/2, respectively), however, RIF tumor cells were not as hypermethylated as TDEC/RIF (48/65% vs 74/82%, respectively) and TRAMP was not hypermethylated when compared to TDEC/TRAMP (26/28% vs 69/77%, respectively). Therefore, despite the consistent result that the CYP24 promoter was hypermethylated in TDEC regardless of the tumor model system examined, tumor cells from these same models differed in their extent of hypermethylation. It should be noted that these models differ in their sensitivity to the anti-proliferative effects of calcitriol *in vivo* with SCC the most sensitive, followed by RIF and TRAMP [4–6,42].

Based on the above data, we next tested whether 5-aza-dC, a classical DNA methyltransferase (DNMT) inhibitor, could reactivate calcitriol-induced CYP24 transcription in TDEC [22]. Treatment of TDEC with 0.25 μ M 5-aza-dC for 96 hours followed by 10 nM calcitriol resulted in a time dependent induction of CYP24 mRNA ($P < 0.005$). By Western blot analysis, TDEC had a higher expression of the DNA methyltransferase enzymes, DNMT1 and DNMT3b when compared to MDEC.

3. Conclusion

While calcitriol is anti-proliferative to a wide variety of tumor cells *in vitro* and *in vivo*, we demonstrate that calcitriol has significant anti-tumor effects on TDEC and these effects are not observed on MDEC or endothelial cells isolated from normal tissues. These differences in calcitriol sensitivity of TDEC and MDEC may be due to the epigenetic silencing of the CYP24 promoter that is hypermethylated in two CpG island regions of TDEC. In addition, when MDEC are incubated with conditioned media from tumor cells, these cells become sensitive to calcitriol concurrent with hypermethylation of the CYP24 promoter. These studies demonstrate epigenetic differences in “normal” cells that only differ in their microenvironment: tumor vs normal. Only recently has evidence suggested a role for epigenetic changes for normal cells that are in the tumor microenvironment. Future studies will examine the mechanisms of epigenetic silencing of CYP24 in endothelial cells in the normal and tumor microenvironments and to determine effects on calcitriol-mediated signaling pathways.

Calcitriol has anti-proliferative effects in pre-clinical prostate models and studies suggest that the endothelial cells are epigenetically silenced at the CYP24 promoter. The silencing of the CYP24 promoter has impact on calcitriol-mediated therapies. While clinical studies have demonstrated limited anti-tumor effects of calcitriol (alone or in combination with steroids/docetaxel) in the treatment of prostate cancer, these studies have been seriously hampered: 1) there has been limited availability of a drug formulation which allows consistent pharmacologic administration of high dose calcitriol; 2) the optimal biologic dose or maximum tolerated dose of calcitriol has not been delineated; and 3) studies have been limited to phase I and II trials and to small and/or poorly designed phase II trials. Studies that overcome these limitations are underway and we are optimistic that an appropriate dose and formulation of calcitriol will be available and able to be tested for its anti-angiogenic effects. In addition, if there is a defect in CYP24 signaling in tumor-derived endothelial cells that facilitates the growth of tumor derived endothelial cells, ways to modulate CYP24 expression or activity in the tumor microenvironment could lead to a more effective treatment plan. Therefore, the information gained through these studies will permit the design of more effective therapeutic approaches for cancer.

Acknowledgments

This work is supported in part by the National Institutes of Health/NCI grant CA95045.

References

1. Bikle DD, Pillai S. Vitamin D, calcium, and epidermal differentiation. *Endocr Rev* 1993;14:3–19. [PubMed: 8491153]
2. Reichel H, Koeffler HP, Norman AW. The role of the vitamin D endocrine system in health and disease. *N Engl J Med* 1989;320:980–991. [PubMed: 2648151]
3. McElwain MC, Dettelbach MA, Modzelewski RA, Russell DM, Uskokovic MR, Smith DC, Trump DL, Johnson CS. Antiproliferative effects *in vitro* and *in vivo* of 1,25-dihydroxyvitamin D3 and a vitamin D3 analog in a squamous cell carcinoma model system. *Mol Cell Differ* 1995;3:31–50.
4. Getzenberg RH, Light BW, Lapco PE, Konety BR, Nangia AK, Acierno JS, Dhir R, Shurin Z, Day RS, Trump DL, Johnson CS. Vitamin D inhibition of prostate adenocarcinoma growth and metastasis in the Dunning rat prostate model system. *Urology* 1997;50:999–1006. [PubMed: 9426741]
5. Modzelewski RA. Apoptotic effects of paclitaxel and calcitriol in rat dunning MLL and human PC-3 prostate tumor cells *in vitro*. *Proc Amer Assoc Cancer Res* 1999;40:580.

6. Hershberger PA, Yu WD, Modzelewski RA, Rueger RM, Johnson CS, Trump DL. Calcitriol (1,25-dihydroxycholecalciferol) enhances paclitaxel antitumor activity *in vitro* and *in vivo* and accelerates paclitaxel-induced apoptosis. *Clin Cancer Res* 2001;7:1043–1051. [PubMed: 11309356]
7. Light BW, Yu WD, McElwain MC, Russell DM, Trump DL, Johnson CS. Potentiation of cisplatin antitumor activity using a vitamin D analogue in a murine squamous cell carcinoma model system. *Cancer Res* 1997;57:3759–3764. [PubMed: 9288784]
8. Hershberger PA, Modzelewski RA, Shurin ZR, Rueger RM, Trump DL, Johnson CS. 1,25-Dihydroxycholecalciferol (1,25-D3) inhibits the growth of squamous cell carcinoma and down-modulates p21(Waf1/Cip1) *in vitro* and *in vivo*. *Cancer Res* 1999;59:2644–2649. [PubMed: 10363987]
9. McGuire TF, Trump DL, Johnson CS. Vitamin D(3)-induced apoptosis of murine squamous cell carcinoma cells. Selective induction of caspase-dependent MEK cleavage and up-regulation of MEKK-1. *J Biol Chem* 2001;276:26365–26373. [PubMed: 11331275]
10. McGuire TF, Trump DL, Johnson CS. 1,25-dihydroxyvitamin D3 induces cytosolic accumulation of MEKK-1 before onset of apoptosis in a p38 MAPK-regulated manner. *Proc Amer Assoc Cancer Res* 2002;2159:435.
11. Hershberger PA, McGuire TF, Yu WD, Zuhowski EG, Schellens JH, Egorin MJ, Trump DL, Johnson CS. Cisplatin potentiates 1,25-dihydroxyvitamin D3-induced apoptosis in association with increased mitogen-activated protein kinase kinase 1 (MEKK-1) expression. *Mol Cancer Ther* 2002;1:821–829. [PubMed: 12492115]
12. Johnson CS, Egorin MJ, Zuhowski R, Parise R, Cappozzoli M, Belani CP, Long GS, Muindi J, Trump DL. Effects of high dose calcitriol (1,25-dihydroxyvitamin D3) on the pharmacokinetics of paclitaxel or carboplatin: results of two phase I studies. *Amer Soc Clin Oncol* 2000;19:210a.
13. Muindi JR, Peng Y, Potter DM, Hershberger PA, Tauch JS, Capozzoli MJ, Egorin MJ, Johnson CS, Trump DL. Pharmacokinetics of high-dose oral calcitriol: results from a phase I trial of calcitriol and paclitaxel. *Clin Pharmacol Ther* 2002;72:648–659. [PubMed: 12496746]
14. Muindi JR, Modzelewski RA, Peng Y, Trump DL, Johnson CS. Pharmacokinetics of 1 α ,25-dihydroxyvitamin D3 in normal mice after systemic exposure to effective and safe antitumor doses. *Oncology* 2004;66:62–66. [PubMed: 15031600]
15. Yu WD, McElwain MC, Modzelewski RA, Russell DM, Smith DC, Trump DL, Johnson CS. Enhancement of 1,25-dihydroxyvitamin D3-mediated antitumor activity with dexamethasone. *J Natl Cancer Inst* 1998;90:134–141. [PubMed: 9450573]
16. Bernardi RJ, Trump DL, Yu WD, McGuire TF, Hershberger PA, Johnson CS. Combination of 1 α ,25-dihydroxyvitamin D(3) with dexamethasone enhances cell cycle arrest and apoptosis: role of nuclear receptor cross-talk and Erk/Akt signaling. *Clin Cancer Res* 2001;7:4164–4173. [PubMed: 11751517]
17. Ahmed S, Johnson CS, Rueger RM, Trump DL. Calcitriol (1,25-dihydroxycholecalciferol) potentiates activity of mitoxantrone/dexamethasone in an androgen independent prostate cancer model. *J Urol* 2002;168:756–761. [PubMed: 12131364]
18. Hershberger PA, Modzelewski RA, Rueger RM, Blum KE, Trump DL, Johnson CS. Enhanced anti-tumor efficacy with dexamethasone/calcitriol/cisplatin therapy: role of p21(waf1). *Proc Amer Assoc Cancer Res* 2000;41:15.
19. Yu WD, Rueger RM, Fuller RW, Johnson CS, Trump DL. 1,25-dihydroxycholecalciferol (calcitriol) enhancement of chemotherapeutic efficacy: synergistic effects by median dose effect. *Proc Amer Assoc Cancer Res* 2001;42:84.
20. Trump DL, Potter DM, Muindi J, Brufsky A, Johnson CS. Phase II trial of high-dose, intermittent calcitriol (1,25 dihydroxyvitamin D3) and dexamethasone in androgen-independent prostate cancer. *Cancer* 2006;106:2136–2142. [PubMed: 16598750]
21. Omdahl JL, Bobrovnikova EA, Choe S, Dwivedi PP, May BK. Overview of regulatory cytochrome P450 enzymes of the vitamin D pathway. *Steroids* 2001;66:381–389. [PubMed: 11179747]
22. Chung I, Karpf AR, Muindi JR, Conroy JM, Nowak NJ, Johnson CS, Trump DL. Epigenetic silencing of CYP24 in tumor-derived endothelial cells contributes to selective growth inhibition by calcitriol. *J Biol Chem* 2007;282(12):8704–8714. [PubMed: 17244627]

23. Jones PA, Baylin SB. The fundamental role of epigenetic events in cancer. *Nat Rev Genet* 2002;3:415–428. [PubMed: 12042769]
24. Fiegl H, Millinger S, Goebel G, Muller-Holzner E, Marth C, Laird PW, Widschwendter M. Breast cancer DNA methylation profiles in cancer cells and tumor stroma: association with HER-2/neu status in primary breast cancer. *Cancer Res* 2006;66:29–33. [PubMed: 16397211]
25. Hu M, Yao J, Cai L, Bachman KE, van den Brule F, Velculescu V, Polyak K. Distinct epigenetic changes in the stromal cells of breast cancers. *Nat Genet* 2005;37:899–905. [PubMed: 16007089]
26. Grover AC, Tangrea MA, Woodson KG, Wallis BS, Hanson JC, Chuaqui RF, Gillespie JW, Erickson HS, Bonner RF, Pohida TJ, Emmert-Buck MR, Libutti SK. Tumor-associated endothelial cells display GSTP1 and RARbeta2 promoter methylation in human prostate cancer. *J Transl Med* 2006;4:13. [PubMed: 16512911]
27. Rodriguez-Canales J, Hanson J, Tangrea M, Erickson H, Albert P, Wallis B, Richardson A, Pinto P, Linehan W, Gillespie J, Merino M, Libutti S, Woodson K, Emmert-Buck M, Chuaqui R. Identification of a unique epigenetic sub-microenvironment in prostate cancer. *J Pathol* 2007;211:410–419. [PubMed: 17278115]
28. Iseki K, Tatsuta M, Uehara H, Iishi H, Yano H, Sakai N, Ishiguro S. Inhibition of angiogenesis as a mechanism for inhibition by 1alpha-hydroxyvitamin D3 and 1,25-dihydroxyvitamin D3 of colon carcinogenesis induced by azoxymethane in Wistar rats. *Int J Cancer* 1999;81:730–733. [PubMed: 10328225]
29. Koli K, Keski-Oja J. 1alpha,25-dihydroxyvitamin D3 and its analogues down-regulate cell invasion-associated proteases in cultured malignant cells. *Cell Growth Differ* 2000;11:221–229. [PubMed: 10775039]
30. Majewski S, Skopinska M, Marczak M, Szmurlo A, Bollag W, Jablonska S. Vitamin D3 is a potent inhibitor of tumor cell-induced angiogenesis. *J Invest Dermatol Symp Proc* 1996;1:97–101.
31. Modzelewski RA, Davies P, Watkins SC, Auerbach R, Chang MJ, Johnson CS. Isolation and identification of fresh tumor-derived endothelial cells from a murine RIF-1 fibrosarcoma. *Cancer Res* 1994;54:336–339. [PubMed: 8275463]
32. Chung I, Wong MK, Flynn G, Yu WD, Johnson CS, Trump DL. Differential antiproliferative effects of calcitriol on tumor-derived and matrigel-derived endothelial cells. *Cancer Res* 2006;66:8565–8573. [PubMed: 16951169]
33. Merke J, Milde P, Lewicka S, Hugel U, Klaus G, Mangelsdorf DJ, Haussler MR, Rauterberg EW, Ritz E. Identification and regulation of 1,25-dihydroxyvitamin D3 receptor activity and biosynthesis of 1,25-dihydroxyvitamin D3. Studies in cultured bovine aortic endothelial cells and human dermal capillaries. *J Clin Invest* 1989;83(6):1903–1915. [PubMed: 2542376]
34. Zehnder D, Bland R, Chana RS, Wheeler DC, Howie AJ, Williams MC, Stewart PM, Hewison M. Synthesis of 1,25-dihydroxyvitamin D(3) by human endothelial cells is regulated by inflammatory cytokines: a novel autocrine determinant of vascular cell adhesion. *J Am Soc Nephrol* 2002;13(3):621–629. [PubMed: 11856765]
35. Bernardi RJ, Johnson CS, Modzelewski RA, Trump DL. Antiproliferative effects of 1alpha,25-dihydroxyvitamin D(3) and vitamin D analogs on tumor-derived endothelial cells. *Endocrinology* 2002;143:2508–2514. [PubMed: 12072382]
36. Flynn G, Chung I, Yu WD, Romano M, Modzelewski RA, Johnson CS, Trump DL. Calcitriol (1,25-Dihydroxycholecalciferol) Selectively Inhibits Proliferation of Freshly Isolated Tumor-Derived Endothelial Cells and Induces Apoptosis. *Oncology* 2007;70:447–457. [PubMed: 17237620]
37. Omdahl, JL.; May, BK. The 25-hydroxyvitamin D 24-hydroxylase. In: Feldman, D.; Pike, JW.; Glorieux, FH., editors. *Vitamin D*. 2. 2005. p. 85-104.
38. Prosser DE, Jones G. Enzymes involved in the activation and inactivation of vitamin D. *Trends Biochem Sci* 2004;29(12):664–673. [PubMed: 15544953]
39. Nebert DW, Dalton TP. The role of cytochrome P450 enzymes in endogenous signalling pathways and environmental carcinogenesis. *Nat Rev Cancer* 2006;6:947–960. [PubMed: 17128211]
40. DeLozier TC, Tsao CC, Coulter SJ, Foley J, Bradbury JA, Zeldin DC, Goldstein JA. CYP2C44, a new murine CYP2C that metabolizes arachidonic acid to unique stereospecific products. *J Pharmacol Exp Ther* 2004;310:845–854. [PubMed: 15084647]

41. Ohyama Y, Kusada T, Yamasaki T, Ide H. Extensive methylation of CpG island of CYP24 gene in osteoblastic ROS17/2.8 cells. *Nucleic Acids Res Suppl* 2002;249–250. [PubMed: 12903199]
42. Alagbala AA, Johnson CS, Trump DL, Posner GH, Foster BA. Antitumor effects of two less-calcemic vitamin D analogs (paricalcitol and QW-1624F2-2) in squamous cell carcinoma (SCC) cells. *Oncology* 2006;70:483–492. [PubMed: 17237623]

Figure 1.

A and B. Calcitriol effect on the vasculature *in vivo*. (A) The effects of calcitriol (daily $\times 3$ $0.625\mu\text{g}/\text{mouse}$) on mean vessel density (MVD) in both SCC tumors and Matrigel plugs where 15 fields and 3 animals in each group were counted. (B) Apoptotic endothelial cells (dark staining) in SCC tumors and Matrigel plugs as determined by double staining of CD31 (endothelial cells) and caspase-3 (apoptosis) on frozen sections from 5 animals/group treated with calcitriol as described above.

Figure 2.

A, B and C. CYP24 enzymes in TDEC and MDEC. (A) microarray screen of CYP enzymes in TDEC and MDEC, (B) schematic diagram showing the position of the two CpG islands at the promoter region of CYP24, (C) methylation analysis of CpG sites located at the CYP24 promoter using bisulfite sequencing. Each circle represents one CpG site.