Published in final edited form as:

Am J Addict. 2010 ; 19(4): 319–324. doi:10.1111/j.1521-0391.2010.00054.x.

Club Drug Use in Hispanic College Students

Michelle R. Resor, BA¹ and Theodore V. Cooper, PhD²

¹Department of Psychology, The University of North Carolina at Charlotte, Charlotte, North Carolina

²Department of Psychology, The University of Texas at El Paso, El Paso, Texas

Abstract

Club drug use and correlates were examined among 251 Hispanic college students on the Texas - México border. Participants completed questionnaires on substance use, club drug attitudes and beliefs, sexual risk-taking behaviors, depressive symptoms, and acculturation. One-quarter of participants reported club drug use. Regression analyses demonstrated that frequency and history of lifetime use were consistently associated with more permissive drug attitudes and other substance use but not sexual risk-taking, depression symptoms, or acculturation. Acculturation was negatively associated with frequency of club drug use, yet positively associated with use of other illicit substances. Avenues for future studies are suggested.

Introduction

Studies suggest that club drugs such as MDMA (ecstasy), ketamine, GHB, and methamphetamine have been used in the past primarily by Non Hispanic Whites, but club drug usage is on the rise among ethnic minorities, including Hispanics.¹⁻⁵ For example, high rates of using methamphetamine have been shown to occur in Hispanics,⁵⁻⁶ and Hispanics with methamphetamine use disorders have evidenced shortened time to relapse.⁷ In samples of Hispanic men who have sex with men (MSM), Fernández and colleagues have found relatively high rates of club drug use, with around half of study participants reporting recent club drug use.⁸⁻¹⁰

College students also may be at risk for increased rates of club drug use and related consequences. Overall rates of illicit substance use among college populations have remained the same or declined slightly within recent years, but rates of using several club drugs among youth and young adults appear on the rise.⁶⁻¹¹ For example, in a primarily Non Hispanic White college sample, 18% of participants reported ever using club drugs, and 11% reported use in the preceding year.¹² Further, one study assessing MDMA use among more than 14,000 college students found increases in MDMA use over time which was associated with more risky behaviors and less studying.¹³

Past studies have demonstrated club drug use may be associated with same-sex sexual activity, and increases in sexual risk-taking, depressive symptoms, and polysubstance use.⁸⁻¹¹⁻¹³⁻¹⁸ In addition to studies assessing club drug use in MSM samples,⁸⁻¹⁰ one study conducted among college students indicated that club drug use was associated with gay, lesbian, and bisexual identification.¹⁷ The positive correlation between club drug use and

Address correspondence to Dr. Cooper, Department of Psychology, 500 West University Ave, The University of Texas at El Paso, El Paso, TX 79968. tvcooper@utep.edu..

Declaration of Interest:

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

risky sexual behavior has also been demonstrated in a variety of samples.^{1-10,16-20} A relationship between depressive symptoms and club drug use has been found as well,²¹⁻²⁴ although evidence suggests that the mechanisms leading to this association are complex. In addition, individuals who use club drugs have reported more experience with other illicit substances and frequently report using multiple substances at once^{14,25-28} which can lead to exponential increases in negative consequences experienced as the result of drug using behaviors. Despite the growing number of Hispanics using club drugs, data assessing club drug use and related concerns in Hispanic populations are limited, particularly among college students and heterosexual individuals.¹⁻³

Studies of acculturation, ethnic identity, and other cultural beliefs and values in Hispanic youth and young adults have primarily focused on alcohol or licit/illicit substance use other than club drugs. Mixed findings were reported in a review of studies that have assessed the relationship between acculturation and substance use among young Hispanics, leading to the conclusion that acculturation studies have not thoroughly examined the complexity of acculturation to U.S. culture and related factors informed by theory in this population.²⁹ Other noteworthy findings associated with Hispanic substance use include that: religious affiliation has been negatively associated with substance use,³⁰ the deterioration of cultural values, such as familismo, has been associated with acculturation, acculturative stress, and heavy alcohol use in young Hispanic males,³¹ and ethnic identity has been positively associated with both heavy alcohol use in Mexican American college males³² and substance use in early adolescents.³³

From a theoretical perspective, according to the Problem Behavior Theory (PBT), risky behaviors often cluster together³⁴ and can be indicative of a single underlying behavioral factor.³⁵ This model has received moderate support in research with Hispanic youth;³⁶ however, cultural factors are thought to influence the way PBT functions in varying groups through unclear mechanisms.³⁶ Further, health behavior researchers have begun to explore the efficacy of multiple risk behavior change, with results often indicating that it is possible to successfully intervene upon several health behaviors simultaneously;³⁷⁻⁴⁰ thus, exploring how other risky behaviors correlate with club drug use is likely to promote the development and assessment of multi-component prevention and intervention efforts.

The current study aimed to assess rates of club drug use in Hispanic college students, measure knowledge and beliefs regarding club drug use, and explore the relationship of club drugs with past sexual experiences, risky sexual behavior, depression, polydrug use, attitudes about club drug use, and acculturation. It was hypothesized that same-sex sexual activity, risky sexual behavior, depressive symptoms, use of other substances, permissive attitudes toward club drug use, and acculturation would demonstrate significant positive relationships with club drug use.

Methods

Participants

Participants were 251 Hispanic students at a university located near the U.S.-México border. Most reported Mexican origin (76% identified themselves as Mexican American, 9% as Mexican National) with 15% identifying with other Latin or Hispanic origin. The mean age was 20.67 years ($SD = 3.76$); 61% were female, 39% were male. Participant characteristics are displayed in Table 1.

Measures

In addition to measuring age, gender and race/ethnicity, participants completed measures assessing substance use, attitudes and beliefs about club drugs, past sexual experiences,

depressive symptoms, and acculturation. Participants also reported their smoking status, which remained a categorical variable in inferential analyses (e.g., daily smoking, weekly smoking, monthly smoking, never smoking, quit smoking).

Substance Use—A club drug use questionnaire asked participants to report any history of using various club drugs. Use was measured for past month, past year, and lifetime. The term “club drugs” is defined in different ways in the literature; for this pilot study we chose to use guidelines suggested by the National Institute on Drug abuse⁴¹ to determine substances that would be categorized as club drugs. Additional literature reviews and internet searches were conducted to research lesser known substances identified by some as club drugs so that participants could be presented with a comprehensive list of club drugs and common slang terms. The list included the following substances: MDMA, GHB (gamma-hydroxybutyrate), GBL (gamma butyrolactone), BD (butanediol), Rohypnol, Ketamine, non-prescribed Viagra, LSD, Psilocybin Mushrooms, and Methamphetamine.

An alcohol, tobacco, and drug use frequency questionnaire asked participants to report past and present usage of other substances (past month, past year, lifetime). Some numerical response options for frequency of individual substance use were incorporated from the Adolescent Problem Severity Index⁴² to capture the range of usage frequency and patterns. For both questionnaires, use of each substance within the past month was chosen from 0 (coded as 0), 1-3 (coded as 1), 4-8 (coded as 2), or >8 (coded as 3). Use for the past year was chosen from 0 (coded as 0), 1-5 (coded as 1), 6-20 (coded as 2), or >20 (coded as 3); these categories were used for lifetime usage as well.

Attitudes and Beliefs about Club Drugs—In a survey measuring club drug attitudes and beliefs, participants were asked to indicate their level of agreement with ten statements (e.g., Using club drugs can be safe if proper precautions are taken) from a questionnaire previously used to assess attitudes about ecstasy and speed.⁴³ These items were internally consistent in the original study,⁴³ and the Cronbach's alpha for the current sample was .82. Two items were added to assess attitudes specific to the border region.

Sexual Experiences—The Kinsey Scale⁴⁴ asked participants to self-report their sexual behaviors on a Likert-type scale ranging from 0 (exclusively heterosexual experiences) to 6 (exclusively homosexual experiences). Most participants (81%) reported only having heterosexual experiences; 16 % reported at least some homosexual experiences and 3% of participants did not report the nature of their sexual experiences. The Sexual Risk subscale of the HIV Risk-taking Behavior Scale⁴⁵ was used to assess sexual risk taking behavior. This measure has demonstrated satisfactory reliability and validity in participants reporting opiate use.⁴⁵ For each item (e.g., How many people have you had sex with during each of the following time periods?), participants were asked to report how often they had engaged in the behavior during the past month, past year, and throughout their lifetime. In the current study this measure demonstrated good internal consistency ($\alpha = .88$).

Depressive Symptoms—The Primary Care Evaluation of Mental Disorders⁴⁶ patient questionnaire nine symptom checklist is composed of a list of depressive symptoms (e.g., suicidal ideations, concentration problems). Participants were asked to indicate how frequently they had experienced each problem in the last two weeks. This measure has shown reliability and validity in primary care samples,⁴⁷ and reliability in this study was adequate ($\alpha = .84$).

Acculturation—The Short Acculturation Scale for Hispanics⁴⁸ contains 12 items measuring the degree to which Hispanics have adopted American culture (e.g., In what language(s) are the TV programs you usually watch?). Scores are obtained by calculating

the scale mean. Adequate psychometric properties have been demonstrated for this measure, 48 and internal consistency was demonstrated in this sample ($\alpha = .92$).

Procedure

Prior to data collection, institutional review board approval was obtained at the university where this study was conducted on the Texas / México border. Participants were undergraduate psychology students aged 18 and older recruited online through the Psychology department participant pool; there were no other inclusion or exclusion criteria. They were informed that the survey contained personal questions regarding substance use and sexual behavior and that responses would remain anonymous. Participants were given a chance to discuss questions with the researchers and then were asked to indicate their consent to participate. Contact information for the university counseling center was provided in the event that students wanted to speak to a counselor about their survey responses. Students received course credit for participation; alternate options for obtaining course credit were available. Students were instructed not to write their names or any identifying marks on the questionnaires, and surveys were filed separately from consent forms.

Approach to Analyses

Descriptives were used to assess participant characteristics on variables of interest. Dependent variables included frequency of club drug use in one's lifetime and lifetime history of club drug use. Independent variables were selected to assess relevant demographic (e.g., gender), theoretically-based (e.g., other substance use), and empirically-derived (e.g., depressive symptoms) correlates. As frequency of club drug use is a continuous dependent variable, multiple linear regression was employed to assess the effects of gender, other substance use, smoking status, attitudes toward club drugs, past sexual experiences, risky sex, depressive symptoms, and acculturation on frequency of lifetime club drug use. As lifetime history of club drug use (vs. no reported history) is a dichotomous dependent variable, logistic regression was used to examine the relationship between potential correlates and reported lifetime history of club drug use.

Results

Participant characteristics are displayed in Table 1. Lifetime use of at least one club drug was reported by 61 participants (25%). Use of any club drug within the past year was reported by 10.7% of participants, and 5.6% reported club use in the preceding month. The most commonly used club drug was psilocybin mushrooms (14% of total sample), followed by MDMA (6% of sample). The most commonly used non club drug illicit substances were marijuana (41%), barbiturates/sedatives (18%), and cocaine (18%).

The multiple regression model was statistically significant, $F(8, 201) = 18.20, p < .001$ (see Table 2). The independent variables accounted for 42% of the variance in club drug use. Lifetime use of other substances was the strongest correlate of club drug use in this model. Other significant correlates were attitudes and beliefs about club drugs and acculturation scores. Club drug use was positively associated with more positive attitudes toward club drugs, and acculturation was negatively associated with club drug use. Interestingly, statistically significant positive relationships were observed between acculturation scores and use of other substances for each time period assessed (i.e., previous month [$r = .135, p = .036$], previous year [$r = .165, p = .010$], lifetime [$r = .190, p = .003$]), indicating a complex relationship between drug use and acculturation in this sample.

The overall logistic regression model was statistically significant ($\chi^2 = 70.779, df = 8, p < .001$; see Table 3). PRIME-MD scores, while not found to predict frequency of club drug use

in the multiple linear regression model, were significantly higher for participants who had ever used club drugs in the logistic regression model. For every unit increase in depression scores, students were 1.12 times more likely to have used club drugs. In addition to greater likelihood of reporting depressive symptomatology, for every unit increase in polysubstance use individuals were 1.84 times more likely to report club drug use. Finally, for every unit increase in more permissive attitudes toward club drugs, individuals were 1.09 times more likely to have used club drugs.

Discussion

This study sought to evaluate rates of club drug use and factors associated with club drug use in a Hispanic college population. Consistent with hypotheses, club drug use was found to be associated with use of other substances and attitudes and beliefs about club drugs. Inconsistent with hypotheses, sexual risk was not found to correlate with club drug use, which is contrary to findings of several previous studies.^{1·8·12·16·20} Frequency of club drug use, but not history of use, was negatively associated with acculturation, while lifetime history of club drug use, but not frequency of use, was positively associated with depressive symptoms. Particularly given the cross sectional nature of this study, interpretation of the latter relationship between club drug use and depression is limited; however, future studies may wish to prospectively assess this relationship, while considering the possibility that depression is implicated in experimentation with club drug use rather than frequency of use.

The finding of polydrug use is consistent with other studies demonstrating risky behaviors clustering (i.e., PBT) and those showing a relationship between club drug use and other substance use.^{14·25·28} These factors should be taken into account during the formation of strategies designed to help individuals reduce negative consequences associated with use of club drugs. For example, discussing the use of substances other than club drugs, including the possible effects of drug combinations, may be necessary to address in this population. Also, challenging beliefs and expectancies about the effects of club drugs may be a useful and promising approach to reducing club drug use.

Findings with regard to acculturation were mixed; higher levels of acculturation were associated with higher levels of non club drug use, while lower levels of acculturation were associated with higher frequency of club drug use. This inconsistency is also reflected in the acculturation and substance use literature, which indicates some studies find positive associations between acculturation and substance use, some find positive relationships between ethnic identity and substance use, while still others find no relationship between acculturation related constructs and substance use.^{29·32·33} Potential reasons for our findings include using club drugs for spiritual practices, lack of knowledge or misperceptions regarding club drug use, greater access and opportunity for use, and the potential for other constructs not measured in this study to inform the relationship. First, in one study of a predominantly Hispanic adolescent sample, using drugs for spiritual or religious purposes was associated with hallucinogen use, yet not with the use of alcohol, marijuana, or stimulants.⁴⁹ The present finding with acculturation is highly similar and may suggest spiritual / religious motives for use. Second, education, both formal and informal (e.g., familial influence), has likely focused on the hazards associated with non club drug substance use for some time, while the dangers of club drug use have been focused on far less. In fact, many club drugs have been made illegal in the U.S. relatively recently⁵⁰ and may still be available over the counter in Mexican farmacias. Third, it may be that club drug access and opportunities to use club drugs are more prominent in México. Not only are club drugs possibly more accessible in pharmacies but also they may be more available in club and/or rave settings. As an example, border region raves may be more prevalent on the Mexican side of the border; thus, lesser acculturated individuals may be more comfortable

and/or socially skilled in such venues. Further, in less acculturated individuals, accessibility (e.g., attending raves) may start at an earlier age than for more acculturated or U.S. residing Hispanics which may be related to club drug use frequency yet not history of use. Finally, it may be that the measurement of other theoretically and empirically relevant constructs not assessed in this study will help clarify the relationship between acculturation and substance use. For example, assessment of constructs such as stress and birthplace may enhance theoretical understandings,²⁹ while the measurement of constructs such as familismo and self esteem may further empirically-based mediational models interpreting this complex relationship.³¹⁻³³

Limitations

There are limitations of this study that should be noted. College students typically do not report high levels of substance use, so patterns observed in this study may not generalize to other young adult or treatment populations. An additional limitation is the nature of the data provided; interpretations of results should take into consideration the limitations associated with reliance on self report and correlational data.

Strengths and Future Directions

Strengths of this study include the assessment of a novel, often underserved population, as well as the assessment of the relationship between club drug use and acculturation on the U.S. – México border.

Future studies should employ longitudinal designs so that the temporal relationships between club drug use and correlates may be examined in depth. This may serve to increase our understanding of the connections between various problem behaviors and possible contributors to these behaviors. Further examination of ethnocultural factors is necessary as well. Specifically, acculturation and its relation to use of individual substances among Hispanic populations should be investigated, and factors that may mediate or moderate this association should be assessed. Such studies will enhance theory-based research into the use of club drugs in underserved populations and promote the empirical development, implementation, and assessment of future prevention and intervention efforts.

Acknowledgments

This research was supported by Grant No. T34 MH019978-05 from the National Institute of Mental Health -Career Opportunities in Research award to Michael A. Zárate, PhD, University of Texas at El Paso, El Paso, Texas.

The PHQ-9 is adapted from PRIMEMDTODAY, developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke, and colleagues, with an educational grant from Pfizer Inc. The names PRIME-MD(r) and PRIMEMDTODAY(tm) are trademarks of Pfizer Inc. Used with permission.

The authors would like to acknowledge the members of the Prevention and Treatment in Clinical Health research laboratory, particularly Hector I. Lopez, B.A., for support and work on this project.

References

1. Novoa RA, Ompad DC, Wu Y, Vlahov D, Galea S. Ecstasy use and its association with sexual behaviors among drug users in New York City. *J Community Health*. 2005; 30:331–343. [PubMed: 16175956]
2. Ompad DC, Galea S, Fuller CM, Phelan D, Vlahov D. Club drug use among minority substance users in New York City. *J Psychoactive Drugs*. 2004; 36:397–399. [PubMed: 15559686]
3. Rawson RA, Gonzales R, Brethen P. Treatment of methamphetamine use disorders: An update. *J Subst Abuse Treat*. 2002; 23:145–150. [PubMed: 12220612]

4. NDIC. National drug threat assessment 2009: MDMA. Available at: <http://www.usdoj.gov/ndic/pubs31/31379/mdma.htm#Overview>. Accessed September 14, 2009
5. Lucker, GW. Texas Department of state health services: Patterns and trends of drug abuse in El Paso, Texas. Available at: <http://www.dshs.state.tx.us/sa/research/currenttrends/1998/elpaso98.html>. Accessed September 15, 2009
6. Johnston, LD.; O'Malley, PM.; Bachman, JG.; Schulenberg, JE. Monitoring the Future National Survey Results on Drug Use, 1975-2004: Volume II, College Students and Adults Ages 19-45. National Institute on Drug Abuse; Bethesda, MD: 2005.
7. Brecht ML, von Mayrhauser C, Anglin MD. Predictors of relapse after treatment for methamphetamine use. *J Psychoactive Drugs*. 2000; 32:211–220. [PubMed: 10908010]
8. Fernández MI, Bowen GS, Varga LM, et al. High rates of club drug use and risky sexual practices among Hispanic men who have sex with men in Miami, Florida. *Subst Use Misuse*. 2005; 40:1347–1362. [PubMed: 16048821]
9. Fernández MI, Perrino T, Bowen GS, et al. Club drug use, sexual behavior, and HIV risk among community and internet samples of Hispanic MSM: Implications for clinicians. *J Soc Work Pract Addict*. 2005; 5:81–100.
10. Akin M, Fernández MI, Bowen GS, Warren JC. HIV risk behaviors of Latin American and Caribbean men who have sex with men in Miami, Florida, USA. *Pan Am J Public Health*. 2008; 23:341–348.
11. Kipke MD, Weiss G, Ramirez M, et al. Club drug use in Los Angeles among young men who have sex with men. *Subst Use Misuse*. 2007; 42:1723–1743. [PubMed: 17934992]
12. Simons JS, Gaher RM, Correia CJ, Bush JA. Club drug use among college students. *Addict Behav*. 2005; 30:1619–1624. [PubMed: 16122624]
13. Strote J, Lee JE, Wechsler H. Increasing MDMA use among college students: Results of a national survey. *J Adolesc Health*. 2002; 30:64–72. [PubMed: 11755802]
14. Kelly BC, Parsons JT. Predictors and comparisons of polydrug and non-polydrug cocaine use in club subcultures. *Am J Drug Alcohol Abuse*. 2008; 34:774–781. [PubMed: 19016183]
15. Substance Abuse and Mental Health Services Administration. Results from the 2008 National Survey on Drug Use and Health: National Findings. Rockville, MD: 2009. Office of Applied Studies, NSDUH Series H-36, HHS Publication No. SMA 09-4434
16. Banta-Green C, Goldbaum G, Kingston S, Golden M, Harruff R, Logan BK. Epidemiology of MDMA and associated club drugs in the Seattle Area. *Subst Use Misuse*. 2005; 40:1295–1315. [PubMed: 16048818]
17. Boyd CJ, McCabe SE, d'Arcy H. Ecstasy use among college undergraduates: Gender, race and sexual identity. *J Subst Abuse Treat*. 2003; 24:209–215. [PubMed: 12810141]
18. Clatts MC, Goldsamt LA, Yi H. Club drug use among young men who have sex with men in NYC: A preliminary epidemiological profile. *Subst Use Misuse*. 2006; 40:1317–1330. [PubMed: 16048819]
19. Eisman JM, Diamond S, Schensul JJ. 'Rollin' on E': A qualitative analysis of ecstasy use among inner city adolescents and young adults. *J Ethn Subst Abuse*. 2005; 4:9–38. [PubMed: 16275633]
20. McElrath K. MDMA and sexual behavior: Ecstasy users' perceptions about sexuality and sexual risk. *Subst Use Misuse*. 2005; 40:1461–1477. [PubMed: 16048828]
21. de Win MML, Reneman L, Reitsma JB, den Heeten GJ, Booij J, van den Brink W. Mood disorders and serotonin transporter density in ecstasy users—The influence of long-term abstinence, dose, and gender. *Psychopharmacology*. 2004; 173:376–382. [PubMed: 14726997]
22. Gold MS, Tabrah H, Frost-Pineda K. Psychopharmacology of MDMA (ecstasy). *Psychiatr Ann*. 2001; 31:675–681.
23. Semple SJ, Patterson TL, Rant I. Methamphetamine use and depressive symptoms among heterosexual men and women. *J Subst Use*. 2005; 10:31–47.
24. Travers KR, Lyvers M. Mood and impulsivity of recreational ecstasy users in the week following a "rave". *Addict Res Theory*. 2005; 13:43–52.
25. Fendrich M, Wislar JS, Johnson TP, Hubbell A. A contextual profile of club drug use among adults in Chicago. *Addiction*. 2003; 98:1693–1703. [PubMed: 14651501]

26. Gouzoulis-Mayfrank E, Daumann J. The confounding problem of polydrug use in recreational ecstasy/MDMA users: A brief overview. *J Psychopharmacol*. 2006; 20:188–193. [PubMed: 16510477]
27. Uys JDK, Niesink RJM. Pharmacological aspects of the combined use of 3,4-methylenedioxymethamphetamine (MDMA, ecstasy) and gamma-hydroxybutyric acid (GHB): A review of the literature. *Drug Alcohol Rev*. 2005; 24:359–368. [PubMed: 16234132]
28. Yacoubian GS Jr, Peters RJ, Palacios WR, Link TC. Identifying the perceived and diagnosed prevalence of ecstasy dependence among club rave attendees. *J Ethn Subst Abuse*. 2004; 3:33–45.
29. De La Rosa MR. Acculturation and Latino adolescents' substance use: A research agenda for the future. *Subst Use Misuse*. 2002; 37:429–456. [PubMed: 12064428]
30. Hodge DR, Andereck K, Montoya H. The protective influence of spiritual-religious lifestyle profiles on tobacco use, alcohol use, and gambling. *Soc Work Res*. 2007; 31:211–219.
31. Gil AG, Wagner EF, Vega WA. Acculturation, familism and alcohol use among Latino adolescent males: Longitudinal relations. *J Community Psychol*. 2000; 28:443–458.
32. Zamboanga BL, Raffaelli M, Horton NJ. Acculturation status and heavy alcohol use among Mexican American college students: Investigating the moderating role of gender. *Addict Behav*. 2006; 31:2188–2198. [PubMed: 16603321]
33. Zamboanga BL, Schwartz SJ, Jarvis LH, Van Tyne K. Acculturation and substance use among Hispanic early adolescents: Investigating the mediating roles of acculturative stress and self-esteem. *J Prim Prev*. 2009; 30:315–333. [PubMed: 19408121]
34. Jessor, R.; Jessor, SL. *Problem Behavior and Psychosocial Development: A Longitudinal Study of Youth*. Academic Press; New York, NY: 1977.
35. Zamboanga BL, Carlo G, Raffaelli M. Problem Behavior Theory: An examination of the behavior structure system in Latino and non-Latino college students. *Interam J Psychol*. 2005; 38:253–362.
36. Dinh KT, Roosa MW, Tein JY, Lopez VA. The relationship between acculturation and problem behavior proneness in a Hispanic youth sample: A longitudinal mediation model. *J Abnorm Child Psychol*. 2002; 30:295–309. [PubMed: 12041714]
37. Nigg CR, Allegrante JP, Ory M. Theory comparison and multiple behavior research: Common themes advancing health behavior research. *Health Educ Res*. 2002; 17:670–679. [PubMed: 12408211]
38. Persky I, Spring B, Vander Wal JS, Pagoto S, Hedeker D. Adherence across behavioral domains in treatment promoting smoking cessation plus weight control. *Health Psychol*. 2005; 24:153–160. [PubMed: 15755229]
39. Prochaska JO, Velicer WF, Rossi JS, et al. Multiple risk expert systems interventions: Impact of simultaneous stage-matched expert system interventions for smoking, high-fat diet, and sun exposure in a population of parents. *Health Psychol*. 2004; 23:503–516. [PubMed: 15367070]
40. Spring B, Pagoto S, Pingitore R, Doran N, Schneider K, Hedeker D. Randomized controlled trial for behavioral smoking and weight control treatment: Effect of concurrent versus sequential intervention. *J Consult Clin Psychol*. 2004; 72:785–796. [PubMed: 15482037]
41. NIDA. Club Drugs. 2009. Available at: <http://www.drugabuse.gov/drugpages/clubdrugs.html>. Accessed September 15, 2009
42. Metzger, DS.; Kushner, H.; McLellan, AT. *Adolescent Problem Severity Index Administration Manual*. Biomedical Computer Research Institute; Philadelphia, PA: 1991.
43. Brewer NT. The relation of internet searching to club drug knowledge and attitudes. *Psychol Health*. 2003; 18:387–401.
44. Kinsey, AC.; Pomeroy, WB.; Martin, CE. *Sexual Behavior in the Human Male*. WB Saunders; Philadelphia: 1948.
45. Darke S, Hall W, Heather N, Ward J, Wodak A. The reliability and validity of a scale to measure HIV risk-taking among intravenous drug users. *AIDS*. 1991; 5:181–185. [PubMed: 2031690]
46. Spitzer RL, Williams JB, Kroenke K, et al. Utility of a new procedure for diagnosing mental disorders in primary care: The PRIME-MD 1000 study. *JAMA*. 1994; 272:1749–1756. [PubMed: 7966923]
47. Kroenke K, Spitzer RL, Williams JBW. The PHQ-9: Validity of a brief depression severity measure. *J Gen Intern Med*. 2001; 16:606–613. [PubMed: 11556941]

48. Marín G, Sabogal F, VanOss MB, Otero-Sabogal R, Perez-Stable E. Development of a short acculturation scale for Hispanics. *Hisp J Behav Sci.* 1987; 9:183–205.
49. Sussman S, Skara S, Rodriguez Y, Pokhrel P. Non drug use- and drug use-specific spirituality as one-year predictors of drug use among high-risk youth. *Subst Use Misuse.* 2006; 41:1801–1816. [PubMed: 17118817]
50. NIDA. Infofacts: Methamphetamine. 2009. Available at:
<http://www.nida.nih.gov/Infofacts/methamphetamine.html>. Accessed September 15, 2009

Table 1

Participant Characteristics

Variable	Frequency		
Gender			
Male	39%		
Female	61%		
Hispanic Identification			
Mexican American	76%		
Mexican National	9%		
Other Hispanic	15%		
Variable	Mean	Standard Deviation	
Age (years)	20.67	3.76	
Kinsey (past sexual experiences)	.42	1.24	
Sexual risk	6.66	5.51	
Attitudes and beliefs about club drugs	17.24	5.52	
PRIME-MD (depressive symptoms)	6.85	5.15	
SASH (acculturation)	3.25	.77	

Table 2

Summary of Regression Analysis for Variables Predicting Frequency of Lifetime Club Drug Use

Variable	<i>B</i>	<i>SE B</i>	<i>B</i>
Gender	.247	.169	.080
Lifetime use of other substances	.465	.050	.636**
Attitudes and beliefs about club drugs	.039	.017	.144*
Sexual risk	−.022	.016	−.079
PRIME-MD (depressive symptoms)	.014	.016	.048
SASH (acculturation)	−.269	.110	−.139*
Smoking status	.054	.043	.078
Kinsey (past sexual experiences)	−.007	.067	−.006

Note. $R^2 = .42$;*
 $p < .05$;**
 $p < .001$

Table 3

Summary of Logistic Regression Analysis for Variables Predicting Reported Lifetime History of Club Drug Use

Variable	Odds ratio	95% Confidence interval
Gender	.767	[.336, 1.747]
Lifetime use of other substances	1.843***	[1.422, 2.388]
Attitudes and beliefs about club drugs	1.090*	[1.005, 1.181]
Sexual risk	1.040	[.964, 1.122]
PRIME-MD (depressive symptoms)	1.116**	[1.038, 1.200]
SASH (acculturation)	.698	[.288, 1.254]
Smoking status	1.127	[.911, 1.396]
Kinsey (past sexual experiences)	.934	[.676, 1.292]

Note. Model $X^2 = 70.779$, $df = 8$, $p < .001$,

Nagelkerke R-square = .432.

*
 $p < .05$;

**
 $p < .01$;

 $p < .001$.