Published in final edited form as:

Gen Comp Endocrinol. 2010 September 1; 168(2): 209–219. doi:10.1016/j.ygcen.2010.03.014.

Molecular mechanisms of corticosteroid synergy with thyroid hormone during tadpole metamorphosis

Ronald M. Bonett^{*}, Eric D. Hoopfer[†], and Robert J. Denver

Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, Michigan 48109-1048

Abstract

Corticosteroids (CS) act synergistically with thyroid hormone (TH) to accelerate amphibian metamorphosis. Earlier studies showed that CS increase nuclear 3,5,3'-triiodothyronine (T₃) binding capacity in tadpole tail, and 5' deiodinase activity in tadpole tissues, increasing the generation of T₃ from thyroxine (T₄). In the present study we investigated CS synergy with TH by analyzing expression of key genes involved in TH and CS signaling using tadpole tail explant cultures, prometamorphic tadpoles, and frog tissue culture cells (XTC-2 and XLT-15). Treatment of tail explants with T₃ at 100 nM, but not at 10 nM caused tail regression. Corticosterone (CORT) at three doses (100, 500, 3400 nM) had no effect or increased tail size. T₃ at 10 nM plus CORT caused tails to regress similar to 100 nM T₃. Thyroid hormone receptor beta (TR β) mRNA was synergistically upregulated by T₃ plus CORT in tail explants, tail and brain *in vivo*, and tissue culture cells. The activating 5' deiodinase type 2 (D2) mRNA was induced by T₃ and CORT in tail explants and tail *in vivo*. Thyroid hormone increased expression of glucocorticoid (GR) and mineralocorticoid receptor (MR) mRNAs. Our findings support that the synergistic actions of TH and CS in metamorphosis occur at the level of expression of genes for TR β and D2, enhancing tissue sensitivity to TH. Concurrently, TH enhances tissue sensitivity to CS by upregulating GR and MR. Environmental stressors can modulate the timing of tadpole metamorphosis in part by CS enhancing the response of tadpole tissues to the actions of TH.

Keywords

Tadpole metamorphosis; thyroid hormone; corticosteroid; stress; nuclear receptors

Introduction

Amphibian metamorphosis is dependent on thyroid hormone (TH), which induces the suite of molecular and cellular changes that cause a tadpole to transform into a frog. Thyroid hormone actions are mediated by TH receptors (TRs; TR α and TR β) that are members of the nuclear receptor (NR) superfamily and function as ligand-activated transcription factors (Mangelsdorf et al., 1995). In *Xenopus laevis*, TR α is expressed shortly after hatching and is

Address all correspondence to: Dr. Robert J. Denver, 3065C Natural Science Building, Department of Molecular, Cellular and Developmental Biology, The University of Michigan, Ann Arbor, MI 48109-1048, USA, Tel. 734-936-6625, FAX 734-647-0884, rdenver@umich.edu.

^{*}Current address: Department of Biological Sciences, University of Tulsa, Tulsa, OK, 74104.

[†]Current address: Division of Biology, California Institute of Technology, Pasadena, CA 91125.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

maintained at a relatively constant level throughout tadpole life and metamorphosis (reviewed by Shi, 2000). The TR α may establish tissue competence to respond to TH once the thyroid gland matures and starts to secrete hormone, and may mediate the actions of TH on cell proliferation (Denver et al., 2009; Furlow and Neff, 2006). By contrast, TR β is expressed at a low level throughout tadpole life, and increases dramatically at the onset of metamorphosis when the plasma TH concentration rises (Shi, 2000). The expression of TR β is dependent on the rise in plasma TH; i.e., it is autoinduced, and the level of TR β expression in target cells is predicted to play a central role in determining the responsiveness of the cell to TH (reviewed by Furlow and Neff, 2006; Tata, 2006).

In addition to the amount of TR made by a cell, an important determinant of TH action during metamorphosis is the activity of intracellular enzymes (deiodinases) that establish the concentration of bioactive TH within the cell (St Germain et al., 2009). The major product of the amphibian thyroid gland is thyroxine (T₄) with minor amounts of 3,5,3'-triiodothyronine (T₃) produced (Buscaglia, 1985; Rosenkilde, 1978). The biologically active form of TH is T₃, which has up to ten times greater activity than T₄ (Frieden, 1981; Lindsay et al., 1967; Rosenkilde, 1978; Wahlborg et al., 1964; White and Nicoll, 1981). Of the three deiodinases expressed in amphibian tissues, two appear to play dominant roles during metamorphosis: the 5' deiodinase type 2 (D2), which converts T₄ to T₃, and the 5 deiodinase type 3 (D3), which converts T₄ to reverse T₃ (3,3',5'-triiodothyronine; rT₃), and T₃ to 3,5-diiodo-L-thyronine (T₂), both of which are inactive in that they do not bind to TRs (reviewed by Denver, 2009a). The balance of D2 and D3 activity within a cell is hypothesized to establish the intracellular level of bioactive hormone, although precisely how this balance is achieved is not well understood (St Germain et al., 2009).

Hormones other than TH play important roles in amphibian metamorphosis, in part by modifying the production and actions of TH. For example, pituitary prolactin may reduce tissue responsiveness to TH by blocking TR β autoinduction. This action of PRL may be important during metamorphic climax when PRL expression increases, possibly to control the morphogenic actions of TH (reviewed by Denver, 2009a). Corticosteroids (CS), hormones produced by adrenocortical cells (interrenal glands in frogs), synergize with TH at target tissues to promote morphogenesis (Denver, 2009b; Kikuyama et al., 1993). The production of CS changes with development, rising throughout metamorphosis and reaching a peak at metamorphic climax (Kikuyama et al., 1993). Also, because CS are stress hormones, their production is increased by exposure to environmental stressors such as habitat dessication, competition for resources, predation risk, etc. (Denver, 2009b). Various environmental stressors have been shown to accelerate tadpole metamorphosis, and this developmental acceleration may be mediated by stress hormones produced both centrally (central nervous system – CNS - and pituitary gland) and peripherally (interrenal glands; Denver, 2009b). At the level of the CNS and pituitary gland, hypothalamic corticotropin-releasing factor (CRF) has a dual hypophysiotropic role in tadpoles, stimulating the secretion of pituitary adrenocorticotrophic hormone (ACTH), which increases CS production by adrenocortical cells, and thyroid stimulating hormone (TSH), which increases hormone secretion by the thyroid gland. At the level of hormone target cells in peripheral tissues, CS have been shown to synergize with TH to promote morphogenesis (Denver, 2009b; Kikuyama et al., 1993).

Like TH, CS actions are mediated by NRs encoded by two different genes: the type I (mineralocorticoid receptor; MR) and the type II (glucocorticoid receptor; GR). Two mechanisms have been proposed to explain CS synergy with TH during tadpole metamorphosis. Work from Kikuyama's group showed that CS increase nuclear T₃ binding capacity in tadpole tail (reviewed by Kikuyama et al., 1993). Galton et al. (1990) found that CS influence deiodinase activity in tadpole tissues, increasing 5' deiodinase and decreasing 5

deiodinase. These findings suggest that CS act to enhance cellular responsiveness by upregulating TRs and at the same time increasing conversion of the precursor T_4 to the biologically active hormone T_3 . Thus, stress hormones mediate environmental effects on development, and they interact with the thyroid axis at both central and peripheral levels.

In the current study we examined morphological and gene expression changes in tadpole tissues and cell lines caused by TH, CS, or combined treatment with the two hormones. We investigated whether the synergy between TH and CS occurs at physiological doses of the hormones, and is seen at the level of TR and deiodinase gene expression. We also investigated the expression of CS receptors, to determine if they are autoregulated, or crossregulated by TH. Our findings provide a molecular basis for understanding the enhanced sensitivity of tadpole tissues to TH caused by increased CS, and support a mechanism for accelerated metamorphosis via interactions among the thyroid and stress hormone axes at the hormone target tissues.

Materials & Methods

Animal care

We raised *Xenopus laevis* and *Silurana tropicalis* tadpoles obtained from in-house breeding in dechlorinated tap water (water temperature 21–23°C - *X. laevis*; 24–25°C - *S. tropicalis*) under a 12L:12D photoperiod and fed frog brittle (*X. laevis*; NASCO, Fort Atkinson, WI) or Seramicon (Sera North America, Inc., Montgomeryville, PA) plus boiled spinach (*S. tropicalis*). All procedures involving animals were conducted in accordance with the guidelines of the University Committee on the Care and Use of Animals of the University of Michigan.

Tissue explant culture and hormone treatment

We initiated tadpole tail explants cultures to investigate the actions of T_3 and corticosterone (CORT) on tail regression and gene expression. We treated Nieuwkoop-Faber stage 52–54 (Nieuwkoop and Faber, 1956) *X. laevis* tadpoles with oxytetracycline (100µg/ml in aquarium water) for 24 h prior to dissection. Dissections were carried out under semi-sterile conditions. Tadpoles were anesthetized in 0.001% benzocaine and dipped in 70% ethanol to sterilize the epidermis. The tails were dissected into sterile 6-well tissue culture dishes containing 2 ml ice-cold tissue culture medium (n=7/treatment for morphological measurements, n=6/treatment for RNA analyses). Tail explants were cultured in Dulbecco's Modified Eagle's Medium (DMEM; Gibco BRL) diluted 1:1.5 for amphibian tissues at 25°C in an atmosphere of 5% CO_2 and 95% O_2 with gentle rotation (50 rpm). The T_3 was dissolved in 0.01 N NaOH and CORT was dissolved in 100% ethanol. All treatments received an equivalent amount of ethanol vehicle (0.001%). Tail explants were treated with T_3 (10 or 100 nM), CORT (100, 500, or 3400 nM) or 10 nM T_3 plus CORT (100, 500 or 3400 nM). The medium was replaced every 12 h. For RNA analysis, three tails were harvested per treatment 2 days after initiation of hormone treatments; the remaining three tails per treatment were harvested at 5 days. Tails were rinsed twice in ice-cold Dulbecco's Phosphate-Buffered Saline (DPBS; Sigma Chemical Co., St Louis, MO), snap frozen in a dry ice-ethanol bath and stored at –80°C until RNA extraction.

We conducted morphometric analyses of tail explants by measuring tail area and final dry weight. For tail area, we captured images every day for seven days with a digital camera and then used Scion Image software (v. 3.0 Scion Corporation, Frederick, MD, USA) to trace the perimeter of each tail and calculate the area. At the end of the experiment we dried the tails in a drying oven and recorded the final dry weight.

Treatment of tadpoles with hormones in vivo

To investigate gene expression responses to T_3 and CORT *in vivo* we treated NF stage 52 *S. silurana* tadpoles with T_3 and CORT for 24 hr before sacrifice. We maintained tadpoles in 4 L aquaria (6 tadpoles per aquarium) and hormones were added to the aquarium water. All treatments received an equivalent amount of ethanol vehicle (0.001%). Tadpoles were treated with T_3 (1, 5 or 50 nM), CORT (100 nM) or 100 nM CORT plus T_3 (1 or 5 nM) and the aquarium water was changed and hormones replenished at 12 hr. At sacrifice tail and brain (a region encompassing the preoptic area/diencephalon) were harvested, snap frozen in a dry ice-ethanol bath and stored at -80°C for subsequent RNA isolation.

X. laevis tissue culture cells

To further investigate gene expression responses to T_3 and CORT, and to establish a cell culture model system for future detailed molecular analyses of hormone synergy we treated the *X. laevis* embryonic fibroblast-derived cell line XTC-2 and the tadpole tail myoblast-derived cell line XLT-15 with hormones and measured gene expression by reverse transcriptase-polymerase chain reaction (RT-PCR). We cultured cells in Leibovitz L-15 medium (diluted 1:1.5 for amphibian tissues) plus 10% Fetal Bovine Serum (Invitrogen, Carlsbad, CA; growth medium) that was stripped of TH and steroids (Samuels et al., 1979; Yao et al., 2008) at 25°C in an atmosphere of 5% CO_2 . All cell culture experiments were performed in 6-well plates. We plated cells overnight in growth medium at a density of 2×10^5 cells/well (XTC-2) and 5×10^4 cells/well (XLT-15). We replaced the growth medium with serum-free L-15 and continued the cultures for 24 h before adding hormones. Cells reached approximately 75% confluency at the start of hormone treatments. We treated cells with hormones or vehicle (ethanol; 0.001%) for 4 hr (XTC-2) or 14 and 24 hr (XLT-15). At the end of the experiment we rinsed cells twice with DPBS then harvested them into 1 ml Trizol (Invitrogen) and proceeded with RNA extraction (described below).

RNA isolation and gene expression analyses

We isolated total RNA from tail explants, freshly dissected tadpole tissues or cell lines using the Trizol reagent (Invitrogen) following the manufacturer's instructions. We analyzed gene expression by either Northern blotting (tail explants), semi-quantitative RT-PCR (XLT-15 cells) or quantitative real time RT-PCR (RTqPCR; tadpole tail and brain, XTC-2 cells) using Taqman assays.

We prepared Northern blots by electrophoresing 10 μg of total tail RNA per lane in a 1% agarose-formaldehyde gel and transferring the RNAs to nylon membranes. Blots were prehybridized in Ultrahyb (Ambion) for 30 min and hybridized for 16 h at 42°C with cDNAs labeled with [^{32}P]-dCTP by random priming (Amersham Corp.). Blots were then washed with 2 X saline sodium citrate (SSC) buffer (0.3 M NaCl, 30 mM trisodium citrate, pH 7.0), 0.5% SDS at room temperature for 10 min and then with 0.25 X SSC, 0.1% SDS at 65°C for 1 h before exposure to x-ray film for 1–14 days. Blots were consecutively stripped and reprobed with [^{32}P]-labeled cDNA probes for *X. laevis* genes. The genes analyzed were TR α A1 (Genbank Accession # M35343), TR β A1 (M35359), GR (X72211), MR (NM001090605), D2 (AF354707) and D3 (L28111). To normalize for RNA loading blots were probed for ribosomal protein L8 (rpL8; NM001086996). X-ray images were digitally captured and densitometric quantitation of the mRNA expression was done using Scion Image software.

For semi-quantitative RT-PCR we treated 1 μg total RNA with 1.5 U RNase-free DNase I (Roche, Indianapolis, IN, USA) to digest contaminating genomic DNA, then reverse-transcribed the RNA into cDNA using 250 ng random hexamers and Superscript II reverse transcriptase following the manufacturer's instructions (see also Bagamasbad et al., 2008;

Manzon and Denver, 2004). Minus RT controls were included to verify no genomic DNA contamination. For PCR we used HotStar Taq DNA polymerase following the manufacturer's instructions (Qiagen, Valencia, CA, USA). Deoxyoligonucleotide primer sequences used for PCR are given in Table 1. Reaction volumes were 50 μ l, and the reaction included an initial 15 min denaturation and activation step at 95 °C. The PCR cycling conditions were 94°C for 45 s, 57 to 63°C for 45 s, and 72 °C for 60 s. The number of cycles used for each gene was determined empirically using a linear amplification range (Table 1). PCR products were electrophoresed in a 1.5% agarose gel and the bands stained with ethidium bromide. Stained bands were captured digitally using a closed-circuit television camera and densitometric analyses were conducted using Scion Image software.

For quantitative real time RT-PCR (RTqPCR) we synthesized cDNAs from 1 μ g DNase-treated total RNA as described above. Custom Taqman assays (Applied Biosystems Inc.) were designed for TR α , TR β , GR, MR, D2, D3, and rpL8 for *X. laevis* and *S. tropicalis* (genome version 4.1; Table 1). Quantitative PCR was conducted using ABsoluteTM qPCR low Rox Mix (ABgene) and reactions were run on either an ABI 3500 or an ABI StepONE Plus qPCR machine. Relative quantities were determined using standard curves generated with pooled cDNAs.

Data analysis and statistics

We normalized all gene expression data to the level of expression of the housekeeping gene rpL8. Northern blot data shown on the graphs are expressed as a percentage of maximum expression normalized to rpL8, but statistics were calculated from band densities for each gene normalized to rpL8. Semi-quantitative and quantitative RT-PCR data are given as ratios of gene expression to rpL8; the normalized gene expression data were log₁₀-transformed before statistical analysis. We analyzed morphometric and gene expression data by one-way analysis of variance (ANOVA) using SYSTAT statistical software (v. 10). Fisher's least significant difference (Fisher's LSD) test was used to test for significant differences among individual treatments ($\alpha=0.05$). We define synergy between T₃ and CORT as: 1) no effect of either hormone alone at the dose tested, but a significant effect with combined treatment, or 2) an effect of combined hormone treatment that is greater than the estimated additive effect by one standard deviation of the combined hormone treatment mean.

Results

Corticosterone synergizes with T₃ to cause tadpole tail regression in vitro

Several groups have reported regression of tadpole tail explants by treatment with TH or TH plus CS (Gray and Janssens, 1990; Kikuyama et al., 1993). Here, we first report confirmation of these findings as a basis for investigating hormone-dependent gene expression changes using this experimental paradigm. We analyzed the effects of hormone treatments on tail area daily over a seven day period, and tail dry weight at the end of the experiment. We treated tails from NF stage 52–54 tadpoles with a dose of T₃ that was previously found not to affect tail regression over a 7 day period (10 nM) and a dose that caused significant tail regression as a positive control (100 nM; Gray and Janssens, 1990; Robinson et al., 1977). We chose three doses of CORT for the experiment: 100 nM, 500 nM and 3400 nM. The high dose, 3400 nM, which would generate supraphysiological tissue content of CORT, was used previously by Gray and Janssens (Gray and Janssens, 1990), and we chose this dose as a positive control in an attempt to replicate their findings. We also chose lower doses of CORT, 100 nM and 500 nM, to generate changes in tissue CORT content within the physiological range for tadpoles. We chose the 100 nM dose because it elevated whole body CORT within the physiological range in tadpoles, and also caused

phenotypic and gene expression responses in tadpoles, frogs and frog cells (Bonett et al., 2009; Glennemeier and Denver, 2002; Hu et al., 2008; Yao et al., 2008). We chose the 500 nM dose in order to generate tissue CORT content in the high physiological range that is achieved during a stress response (Krain and Denver, 2004; Yao et al., 2008).

We saw a significant effect of treatment ($F_{(8, 413)}=43.919$, $P<0.0001$) and day ($F_{(6, 413)}=32.918$, $P<0.0001$) on tail area (Figure 1A), and a significant interaction between treatment and time ($F_{(48, 413)}=3.795$, $P<0.0001$). Tail area diverged among treatments by day 4 ($F_{(8, 60)}=5.194$, $P<0.0001$). At day 7, the areas of tails treated with 100 nM T_3 , or T_3 plus all three doses of CORT were not different from each other, but all were significantly smaller than the control, CORT only treatments, or 10 nM T_3 ($F_{(8, 55)}=31.833$, $P<0.0001$). The final dry weight of tails treated with 100 nM T_3 , or T_3 + CORT (all doses) was significantly less than the control, CORT only, or 10 nM T_3 treatments (Figure 1B; $F_{(8, 54)}=9.699$, $P<0.0001$). Treatment with CORT alone tended to increase final tail dry weight, and this was significantly different from controls for the 3400 nM dose.

Corticosterone synergizes with a low dose of T_3 to induce gene expression in tadpole tail explants

Hormone treatments caused significant elevations in TR β mRNA in tadpole tail explants (48 hr $F_{(8, 17)}=4.854$, $P<0.005$; 120 hr $F_{(8, 17)}=5.103$, $P<0.004$). Thyroid hormone receptor β mRNA was autoinduced, a well known phenomenon that occurs in tadpole tissues during metamorphosis (Fig. 2; reviewed by Tata, 2006). Treatment with CORT alone elevated TR β mRNA at 48 hr but not at 120 hr. Corticosterone plus 10 nM T_3 elevated TR β mRNA that was greater than additive at all doses and timepoints analyzed. The effects of the hormones were synergistic (synergy as defined earlier - see Materials and Methods) at 48 hr for the 100 nM CORT dose, and at 120 hr for the 500 nM and 3400 nM CORT doses. Although there were trends in TR α mRNA levels caused by hormone treatments that were similar to those observed for TR β , these changes were not statistically significant (data not shown).

Deiodinase type 2 mRNA in control tail explants was undetectable in our assay (even after 2 week exposure of the Northern blots), which agrees with the findings of Shintani et al. (2002). Because the controls were undetectable we could not conduct a comparative densitometric analysis. Representative Northern blots are shown in Fig. 3. The 100 nM dose of T_3 caused a small increase in D2 mRNA at 48 hr, and this appeared greater at 120 hr. Neither the 10 nM dose of T_3 nor any of the doses of CORT alone affected D2 mRNA at any time point. Combined treatment with 10 nM T_3 plus CORT at each of the doses tested increased D2 mRNA, and this effect appeared greater at the 120 hr time point. Thus, combined treatment with a low dose of T_3 plus CORT caused synergistic upregulation of D2 mRNA.

Deiodinase type 3 mRNA was detectable in control tail explants, similar to what was found previously in tadpole tail by Brown et al. (1996). Only the 100 nM dose of T_3 caused a statistically significant ($P<0.05$) increase that was comparable in magnitude at both time points measured (data not shown). Treatment with CORT with or without T_3 tended to reduce D3 mRNA in tail explants at both time points but these effects were not statistically significant. GR mRNA was upregulated by 100 nM T_3 at both time points ($P<0.05$) but unaffected by CORT alone or in combination with T_3 (data not shown; see also Krain and Denver, 2004). Although we detected MR mRNA in tail by Northern blotting we were unable to quantify its expression.

Corticosterone synergizes with T₃ to induce gene expression in tadpole tail in vivo

We treated NF stage 52 *S. tropicalis* tadpoles with hormones (1, 5 or 50 nM T₃, 100 nM CORT, 1 nM T₃ + 100 nM CORT, 5 nM T₃ + 100 nM CORT) by addition to the aquarium water for 24 hr before sacrifice and tissue harvest. Hormone treatments caused significant changes compared with controls in expression of all genes analyzed in the tadpole tail: TR α ($F_{(6, 35)} = 11.1$, $P = 0.0001$), TR β ($F_{(6, 35)} = 38.22$, $P < 0.0001$), GR ($F_{(6, 35)} = 10.6$, $P < 0.0001$), MR ($F_{(6, 35)} = 3.2$, $P < 0.012$), D2 ($F_{(6, 29)} = 13.07$, $P < 0.0001$), and D3 ($F_{(6, 35)} = 6.4$, $P < 0.0001$). TR α mRNA was increased by T₃ (5 and 50 nM) or CORT (100 nM) alone, and the increase was greater than additive for the combined hormone treatments; it was synergistic for the 5 nM dose of T₃ (Fig. 4A). TR β mRNA was increased by all doses of T₃ tested, but not by CORT alone. The increase in TR β mRNA was synergistic for the 5 nM dose of T₃ plus CORT (Fig. 4B).

Deiodinase type 2 mRNA was increased by treatment with 50 nM T₃ or 100 nM CORT (Fig. 5A). Combined treatment with 100 nM CORT plus 1 nM or 5 nM T₃ also caused significant increases in D2 mRNA, but these actions were not additive nor synergistic. Like D2, D3 mRNA was also increased by 50 nM T₃ or 100 nM CORT (Fig. 5B). Combined treatment with 100 nM CORT plus 1 or 5 nM T₃ produced responses that were greater than additive, and this effect was synergistic for the 5 nM T₃ dose.

Glucocorticoid receptor mRNA was increased by 50 nM T₃, but was not affected by the lower doses of T₃ or by CORT, nor by any of the combined hormone treatments (Fig. 6A). Similar to GR, MR mRNA was increased by 50 nM T₃, but not by 1 nM T₃ or by CORT; there was a trend towards increased MR mRNA by 5 nM T₃ and the combined treatment of 5 nM T₃ plus CORT (Fig. 6B).

In the same experiment we analyzed TR β and D2 mRNA expression in the brain (preoptic area/diencephalon; Suppl. Fig. S1). Hormone treatments caused statistically significant changes in TR β mRNA ($F_{(6, 35)} = 62.24$, $P < 0.0001$); TR β was strongly autoinduced by all doses of T₃ tested (Fig. S1A). Treatment with CORT caused a small but statistically significant increase in TR β mRNA. Combined treatment of 5 nM T₃ plus CORT caused synergistic upregulation of TR β mRNA. Hormone treatments caused statistically significant changes in D2 mRNA ($F_{(6, 35)} = 4.94$, $P = 0.001$). D2 was increased by 50 nM T₃, but decreased by 100 nM CORT; other hormone treatments had no statistically significant effects (Fig. S1B).

Corticosterone and T₃ influence gene expression in *X. laevis* tissue culture cells

The embryonic fibroblast-derived *X. laevis* cell line XTC-2 responds to T₃ in a dose and time-dependent manner by strongly upregulating TR β mRNA (Iwamuro and Tata, 1995; Kanamori and Brown, 1992) and it expresses both GR and MR (Bonett et al., 2009). We therefore tested for effects of CORT with or without T₃ on gene expression in XTC-2 cells. We treated cells with 1 nM T₃ or 1 nM T₃ + 100 nM CORT for 4 hr to analyze early gene expression responses. Hormone treatments caused statistically significant changes in expression of mRNAs for TR α ($F_{(3, 17)} = 16.12$, $P < 0.0001$), TR β ($F_{(3, 20)} = 139.335$, $P < 0.0001$), D3 ($F_{(3, 17)} = 224.29$, $P < 0.0001$), GR ($F_{(3, 16)} = 25.35$, $P < 0.0001$) and MR ($F_{(3, 17)} = 125.32$, $P < 0.0001$) but not for D2 (Fig. 7). TR α mRNA was increased by treatment with T₃ but not by CORT; no synergistic action was observed. Similar to TR α , TR β mRNA was increased by T₃ but not by CORT; the response of TR β mRNA to T₃ was considerably greater than that of TR α (TR β >130 fold; TR α 1.5 fold). Combined treatment of T₃ with CORT caused synergistic upregulation of TR β mRNA (Fig. 7).

The expression of D3 mRNA was strongly increased by 1 nM T₃, and although CORT alone had no effect, CORT reduced the T₃-dependent increase by one half. GR mRNA was

reduced by CORT, but induced by T_3 . Cotreatment with CORT blocked the T_3 -induced increase in GR mRNA. The expression of MR mRNA was strongly induced by T_3 ; CORT had no effect alone or in combination with T_3 .

We also investigated the effects of T_3 and CORT on TR β and GR mRNA expression in the tadpole tail myoblast-derived cell line XLT-15 (Fig. 8). We found statistically significant effects of hormone treatments on TR β (14 hr: $F_{(4,25)}=31.44$, $P<0.0001$; 24 hr: $F_{(4,25)}=24.26$, $P<0.0001$) and GR mRNA ($F_{(4,25)}=3.74$, $P=0.016$; $F_{(4,25)}=7.18$, $P=0.001$). Expression of TR β mRNA was increased by both doses of T_3 at each of the two time points investigated (Fig. 8A). Corticosterone alone had no effect, but CORT synergized with T_3 to increase TR β mRNA. GR mRNA was induced by T_3 , and the largest increase was seen at 24 hr (Fig. 8B). There were no effects of CORT on GR mRNA in XLT-15 cells. We could not detect D2 or D3 mRNA in this cell line; TR α and MR were not investigated.

Discussion

Earlier studies showed that CS can synergize with TH to promote tadpole tail resorption, and biochemical analyses supported that CS can enhance TH action by upregulating TRs and 5' deiodinase (reviewed by Kikuyama et al., 1993). Many of these studies used doses of CS that would be expected to produce supraphysiological tissue hormone content (e.g., Gray and Janssens, 1990), used the superactive glucocorticoid analog dexamethasone (DEX), or were not replicated, and therefore precluded quantitative or statistical analysis (e.g., Iwamuro and Tata, 1995). Here we provide a molecular basis for synergy among CS and TH in tadpole metamorphosis through analysis of gene expression using several experimental paradigms and doses of CORT that we previously showed produce tissue hormone content within the physiological range. A major effect of CS is at the level of the TR β gene, whose mRNA was increased in a synergistic way by combined treatment with T_3 and CORT. This effect on TR β was seen in tail explants, in tadpole tail and brain *in vivo*, and in two *X. laevis* tissue culture cell lines, XTC-2 and XLT-15. Furthermore, the kinetics in XTC-2 cells was rapid, with significant potentiation of mRNA levels by 4 hr of hormone treatment. A second basis for the synergy is at the level of D2 gene expression, where CORT induced D2 mRNA, or acted in synergy with T_3 . Thus, CS may enhance tissue sensitivity to TH by upregulating expression of TR β , increasing cellular responsiveness to the hormone, and by upregulating D2, thereby promoting generation of the more biologically active form of the hormone T_3 .

The potentiating actions of CS on TH-induced tadpole metamorphosis have been known since the mid-1950s, and were first described by Freiden and Naile (1955) who showed that hydrocortisone accelerated both T_4 and T_3 -induced metamorphosis in three anuran species. Similar findings were obtained by others with tadpoles of different anuran species including *X. laevis* (Gray and Janssens, 1990; reviewed by Denver, 2009a; Kikuyama et al., 1993). A physiological role for CS in tadpole metamorphosis is supported by the findings of Kikuyama et al. (1982) who showed that treatment of tadpoles with the CS synthesis inhibitor amphenone B retarded T_4 -induced tail resorption, and that this could be reversed by treatment with CS. Several investigators reported synergistic actions between CS and TH in tadpole tail explants (Kikuyama et al., 1993; Kikuyama et al., 1983). Gray and Janssens (Gray and Janssens, 1990) found synergy between T_3 (50 nM) and CORT (3400 nM) in *X. laevis* tadpole tail explants; lower doses of CORT were not tested. We found that doses of CORT (100 and 500 nM) that were previously shown to produce tissue content within the physiological range for amphibians caused significant synergistic actions with T_3 on tadpole tail regression *in vitro*, thus supporting a physiological role for CS in tadpole tail regression. Furthermore, these actions were seen with a dose of T_3 (10 nM) that by itself had no effect on tail regression, which provides convincing evidence for hormone synergy.

Work from Kikuyama's laboratory showed that CS increase nuclear binding capacity for T_3 in tadpole tail (Niki et al., 1981; Suzuki and Kikuyama, 1983). The frog TR β gene is autoinduced during metamorphosis (reviewed by Tata, 2006). The synergy that we saw among T_3 and CORT on regression of tadpole tail explants was paralleled by a simultaneous, synergistic upregulation of TR β mRNA. This synergy was also seen in the tail and brain of tadpoles treated with hormones *in vivo*, and in both XTC-2 and XLT-15 cells. Iwamuro and Tata (1995) reported that the potent glucocorticoid DEX (they used a dose roughly equivalent to 1000 nM CORT) enhanced T_3 -dependent TR β expression in tadpole tail explants and XTC-2 cells. There are other examples where CS potentiates TH actions on gene expression. For example, in rodents, TH and CS act synergistically to induce growth hormone (GH) gene expression in pituitary and pituitary cell lines (Dobner et al., 1981; Martial et al., 1977; Nogami et al., 1995; Samuels et al., 1977; Shapiro et al., 1978). Glucocorticoids also act synergistically to increase somatotrope abundance in the rat (Hemming et al., 1988) and chick embryonic pituitary gland, and in the chick this may be attributed to an increase in 5' deiodinase activity (Liu et al., 2003). Similar synergy between TH and CS was observed on several hepatic genes in mammals (Menjo et al., 1993; Molero et al., 1993; Yamaguchi et al., 1999).

While the autoinduction of TR β is explained by the presence of TH response elements (TREs) located in the proximal promoter region of the gene (Ranjan et al., 1994; Machuca et al., 1995), we do not yet understand the molecular basis for synergy between T_3 and CORT on TR β . One hypothesis is that CS responsive elements are present in the TR β promoter region perhaps adjacent to or overlapping with the TREs. However, we found no evidence that CS could modulate TR β promoter activity in transient transfection assays using XTC-2 or XLT-15 cells (E.D. Hooper, C. Kholdani and R.J. Denver, unpublished data; see also Iwamuro and Tata, 1995). This suggests that 1) CS responsive elements are located elsewhere in the TR β locus, or 2) CS induce the expression of a transcription factor that enhances TR β autoinduction. It is also possible that CS act postranscriptionally to stabilize TR β mRNA. A direct transcriptional mechanism of action may occur in mammals where CS increased nuclear T_3 binding capacity and TR β 1 transcription in rat liver (Montesinos et al., 2006). Montesinos et al. (2006) also provided some evidence that CS action on TR β 1 transcription was due to direct binding and transactivation of the TR β promoter by GR. Synergy between TR and GR was reported in rat pituitary tumor cells and required the GR DNA-binding domain and was mediated by the GR transactivation domains (Leers et al., 1994).

The kinetics of synergistic TR β upregulation by T_3 plus CORT was rapid in XTC-2 cells suggesting direct actions of CS on TR β transcription in tadpoles. However, we cannot rule out that the actions of CS were due to upregulation of a transcription factor that regulates the TR β gene rather than direct modulation of TR β by GR or MR. A candidate for this transcription factor is the immediate early gene Krüppel-like factor 9 (KLF9; also known as basic transcription element binding protein 1; BTEB1). Krüppel-like factor 9 is the most rapidly responding TH-induced gene thus far identified in tadpole tissues, and its expression parallels the increase in plasma TH that occurs during tadpole metamorphosis (Denver et al., 1997; Furlow and Kanamori, 2002; Hooper et al., 2002; Shi, 2000; Wang and Brown, 1993). The TH responsiveness of the frog gene can be attributed to a TRE located ~6.5 kb upstream of the transcription start site of the *X. laevis* KLF9 gene (Furlow and Kanamori, 2002). Our previous work showed that KLF9 functions as an accessory transcription factor for TR β autoinduction (Bagamasbad et al., 2008). We also found that KLF9 is strongly induced by CORT in juvenile frog (Bonett et al., 2009) and tadpole brain (F. Hu and R.J. Denver, unpublished data). Indeed, KLF9 is synergistically upregulated by TR and GR via a highly conserved 'synergy module' located in the 5' flanking region of frog and mouse

genes (P. Bagamasbad and R.J. Denver, unpublished data). Therefore, rapid CS upregulation of KLF9 could be responsible for the synergistic activation of TR β by T₃ and CORT.

Another possibility for a CS-regulated transcription factor is retinoid X receptor (RXR) which heterodimerizes with TR on positive TREs. Iwamuro and Tata (Iwamuro and Tata, 1995) reported that while T₃ alone downregulated RXR γ mRNA in *X. laevis* tadpole tail explants, DEX alone increased expression (but T₃ suppressed the DEX effect when the two hormones were combined). Treatment of rat hepatocytes with DEX potentiated T₃ effects on type 1 deiodinase, Spot 14 and malic enzyme mRNAs (Menjo et al., 1993; Yamaguchi et al., 1999). DEX alone increased RXR α mRNA, and these authors proposed that the enhancement of T₃ action was due to the upregulation of RXR α .

Although TR α exhibited a similar pattern of response to TR β in tadpole tail explants, these effects were subtle and did not reach statistical significance. However, we found statistically significant responses to hormone treatment of TR α mRNA in tail of tadpoles treated *in vivo*, and in XTC-2 cells. In these experiments TR α mRNA was induced by CORT or by T₃, and was synergistically induced by the higher dose of T₃ (5 nM) in tadpole tail *in vivo*, but not in XTC-2 cells at 4 hr. Taken together, our results suggest that the more significant synergistic regulation by T₃ and CORT occurs on TR β as compared with TR α .

The intracellular concentration of T₃ is modulated by the activities of D2 and D3 (St Germain et al., 2009). Our findings support that a secondary mechanism for CORT to enhance tissue responsiveness to TH is through upregulation of D2, either alone or in synergy with T₃. We consider this to be secondary because the hormonal response of D2 may be delayed as compared with the immediate and strong upregulation of TR β . We found that CORT alone could induce D2 mRNA in tadpole tail *in vivo* but there was no additive or synergistic effects of combined treatment with T₃ plus CORT at the single time point measured (24 hr; see Fig. 4A). In tadpole tail explants the expression of D2 mRNA was induced by combined treatment with T₃ (10 nM) plus CORT (500 and 3400 nM) at both time points investigated (48 and 120 hr). Although we detected D2 mRNA in XTC-2 cells it was not induced by T₃ by 4 hr, which may reflect that D2 is a delayed response gene (discussed below). Thus, our data suggest that T₃ and CORT can promote expression of D2 in tadpole tail. The action of T₃ is consistent with earlier findings that showed that TH positively regulates 5' deiodinase activity, and D2 mRNA in tadpoles (Brown, 2005; Buscaglia, 1985; Hogan et al., 2007). Our finding that CORT induced D2 mRNA in tadpole tail *in vivo* agrees with the findings of Galton (1990) who reported that treatment of bullfrog tadpoles with 1 μ M CORT in their aquarium water increased 5' deiodinase activity in skin.

The 5 and 5' deiodinase activities, and D2 and D3 mRNAs show tissue-specific and developmental stage-specific expression patterns in tadpoles that reflect the asynchronous tissue morphogenesis that occurs during metamorphosis, and the roles that the deiodinases play in modulating TH action (Brown, 2005; St Germain et al., 2009). Both enzymes may be co-expressed in cells, the relative expression levels providing a push-pull mechanism to regulate intracellular T₃ concentrations (Becker et al., 1997), or they may be expressed at different times of development. In the tadpole tail D3 mRNA is found in several cell types including tail fibroblasts, but not muscle cells (Berry et al., 1998), and D2 is expressed primarily in fibroblasts (Cai and Brown, 2004). D3 mRNA (and 5 deiodinase activity) is expressed in the tail throughout tadpole life, increases during late prometamorphosis (NF stage 59–61) but then declines sharply at metamorphic climax (Brown et al., 1996; St Germain et al., 1994). The tadpole tail is an essential locomotory organ, and this pattern of D3 expression is hypothesized to protect the tail from the actions of T₃ before the animal reaches an appropriate stage of development to initiate tail regression (Brown, 2005). By contrast, D2 expression is low or nondetectable before metamorphic climax (NF stage 62)

when its expression increases strongly and continues through NF stage 64 (Cai and Brown, 2004). Our failure to detect D2 or D3 mRNA in the tail myoblast-derived XLT-15 cells is consistent with *in situ* hybridization studies that showed that these genes are not expressed in tail muscle cells. The expression of D2 and D3 in non-muscle tail cells suggests that these cells either destroy bioactive TH to protect tail muscle cells from apoptosis (when D3 is expressed), or generate a high local concentration of T₃ to act on tail muscle cells to cause tail regression during metamorphic climax (when D2 is expressed). This situation is analogous to that in mammalian brain, where astrocytes, but not neurons, express deiodinases and therefore control exposure of neurons to bioactive TH (Trentin, 2006).

The increase in D3 expression is coincident with a rise in circulating T₃ during prometamorphosis (Leloup and Buscaglia, 1977). The 5 deiodinase activity and D3 mRNA are induced by TH in tail (Brown et al., 1996; St Germain et al., 1994; Wang and Brown, 1993), brain (Denver et al., 1997; Hogan et al., 2007) and in XTC-2 cells (St Germain et al., 1994). We also observed this regulation of D3 in tadpole tail and brain, and the early response kinetics in XTC-2 cells supports that D3 is a direct TH target gene (Das et al., 2006; St Germain et al., 1994). We observed a small increase in D3 mRNA in the tails from tadpoles treated with CORT for 24 hr, and also synergy with T₃. However, in XTC-2 cells CORT alone had no effect, but CORT reduced the response to T₃ in D3 mRNA. Galton (1990) found that treatment of bullfrog tadpoles with CORT for 48 hr reduced 5 deiodinase activity in tadpole liver and gut, and Lorenz et al. (2009) reported that treatment of *X. laevis* tadpoles with CORT or dexamethasone for 21 days decreased D3 mRNA. We observed a trend towards decreasing D3 mRNA with CORT in tadpole tail explants at both time points. In the chicken, glucocorticoids have been shown to downregulate 5 deiodinase activity and D3 mRNA (reviewed by De Groef et al., 2008). Taken together, the findings suggest that the actions of CS on D3 expression are tissue-specific and time-dependent. CORT may synergize with TH to upregulate D3 mRNA in tail as an early response (by contrast to its inhibitory effect on T₃-induced upregulation of D3 mRNA in XTC-2 cells). However, over a longer period, D3 may be repressed by CS in most tissues. For reasons that are unknown, the expression of the D3 gene is turned down at metamorphic climax when TH concentrations are highest. Expression of other direct TR target genes (e.g., KLF9) continues to increase in the tadpole tail through metamorphic climax (Brown et al., 1996). This is a time when plasma CS are also rising in tadpoles (reviewed by Denver, 2009a), which suggests the hypothesis that the late decrease in D3 is related to the elevated CS.

Unlike D3 which shows early response kinetics, D2 appears to be a delayed response gene. Brown (2005) commented that the upregulation of D2 mRNA caused by T₃ required several days of hormone treatment, and Das et al. (2006) did not find D2 to be a direct T₃ target in microarray studies. Our results in tadpole tail explants are consistent with these findings in that we saw only a small effect of T₃ on D2 mRNA at 48 hr but clear upregulation by 120 hr. However, when we treated tadpoles with T₃ (50 nM) we saw upregulation of D2 mRNA by 24 hr in the tail and in the brain; we did not examine earlier timepoints. More study is needed to understand the mechanisms by which D2 gene expression is regulated by TH in *X. laevis* and *S. tropicalis*.

Another important point of regulation in this system is the induction of GR and MR expression by T₃, which would be expected to increase tissue sensitivity to further actions of CS. We found that treatment of tadpoles with T₃ induced GR and MR mRNA in tail. We obtained similar results in XTC-2 cells for both genes, and in tail explants and XLT-15 cells for GR. The positive effect of T₃ on GR in tail in *S. tropicalis* is consistent with our previous findings in *X. laevis* (Krain and Denver, 2004). CORT treatment did not affect GR or MR mRNA in tadpole tail *in vivo* or in tail explants. However, GR was negatively regulated by

CORT in XTC-2 cells; GR is negatively regulated by T_3 and CORT in tadpole brain (Krain and Denver, 2004; Yao et al., 2008).

By contrast to the synergistic effect of combined treatment with T_3 plus CORT, treatment of tadpole tail explants with CORT alone tended to increase final tail dry weight which was statistically significant for the highest CORT dose. We do not know whether this effect is of physiological significance. However, it is worth noting that CORT treatment *in vivo* increased tail size of *Rana pipiens* (Glennemeier and Denver, 2002) and *Rana sylvatica* tadpoles (J. Middlemis-Maher, E.E. Werner, and R.J. Denver, unpublished data). The mechanism for this effect is unknown, but it may have functional importance with regard to predation, where chronic predator presence elevates CORT in premetamorphic tadpoles (J. Middlemis-Maher, E.E. Werner, and R.J. Denver, unpublished data). Tadpoles of many amphibian species exposed to predation develop larger tails (Relyea, 2007), and this increased tail size could be blocked with the CS synthesis inhibitor metyrapone (J. Middlemis-Maher, E.E. Werner, and R.J. Denver, unpublished data).

In summary, TH and CS act synergistically to cause regression of the tadpole tail. This synergy may be explained by the cooperative upregulation of TR β and D2 expression which should increase tissue sensitivity to T_3 , and enhance conversion of T_4 to T_3 , the more biologically active form of the hormone. Concurrently, CS may cause early upregulation of D3, an adaptive response to protect the tail from the apoptotic actions of T_3 , but with prolonged exposure cause downregulation of D3 to promote tail regression. Thus, the early response to hormones is to upregulate TR β and D3, the delayed response is to upregulate D2 and to downregulate D3. The TR β gene expression response is sustained and amplified due to autoinduction (which is favored by the synergistic upregulation of the immediate early gene KLF9), synergy with CS, and enhanced generation of T_3 in target tissues. The CS response is also enhanced by the actions of T_3 on the expression of GR and MR.

The ecological/organismal significance of this hormonal gene regulation during development is that activation of the stress axis in a tadpole by environmental stressors would 1) lead to increased production of TH and CS (through increased CRF secretion), and 2) enhance tissue sensitivity of target tissues to the TH signal through the actions of CS. Together, these two hormonal systems may cooperate to accelerate metamorphosis under conditions of ecological stress (e.g., pond drying, competition for resources, predation, habitat degradation, etc.)

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to Cyrus Kholdani for providing technical assistance. We thank Jamshed Tata for providing the XTC-2 cells, and Yoshio Yaoita for providing the XLT-15 cells. Yun-Bo Shi provided plasmids with cDNAs for *X. laevis* TR α A1, TR β A1 and rPL8, and Olivier Destree provided the plasmid with a cDNA for the *X. laevis* GR. This research was supported by grants from the National Science Foundation (IBN 0235401) and the National Institutes of Health (1 R01 NS046690) to R.J.D.

Literature Cited

Bagamasbad P, Howdeshell KL, Sachs LM, Demeneix BA, Denver RJ. A role for basic transcription element-binding protein 1 (BTEB1) in the autoinduction of thyroid hormone receptor beta. *Journal of Biological Chemistry* 2008;283:2275–2285. [PubMed: 18045867]

- Becker KB, Stephens KC, Davey JC, Schneider MJ, Galton VA. The type 2 and type 3 iodothyronine deiodinases play important roles in coordinating development in *Rana catesbeiana* tadpoles. *Endocrinology* 1997;138:2989–2997. [PubMed: 9202244]
- Berry DL, Schwartzman RA, Brown DD. The expression pattern of thyroid hormone response genes in the tadpole tail identifies multiple resorption programs. *Developmental Biology* 1998;203:12–23. [PubMed: 9806769]
- Bonett RM, Hu F, Bagamasbad P, Denver RJ. Stressor and glucocorticoid-dependent induction of the immediate early gene Krüppel-like factor 9: implications for neural development and plasticity. *Endocrinology* 2009;150:1757–1765. [PubMed: 19036875]
- Brown DD. The role of deiodinases in amphibian metamorphosis. *Thyroid* 2005;15:815–821. [PubMed: 16131324]
- Brown DD, Wang Z, Furlow JD, Kanamori A, Schwartzman RA, Remo BF, Pinder A. The thyroid hormone-induced tail resorption program during *Xenopus laevis* metamorphosis. *Proceedings of the National Academy of Sciences of the United States of America* 1996;93:1924–1929. [PubMed: 8700860]
- Buscaglia, M.; Leloup, J.; De Luze, A. The role and regulation of monodeiodination of thyroxine to 3,5,3'-triiodothyronine during amphibian metamorphosis. In: Bownes, MBaM, editor. *Metamorphosis*. Clarendon Press; Oxford: 1985. p. 273–293.
- Cai LQ, Brown DD. Expression of type II iodothyronine deiodinase marks the time that a tissue responds to thyroid hormone-induced metamorphosis in *Xenopus laevis*. *Developmental Biology* 2004;266:87–95. [PubMed: 14729480]
- Das B, Cai LQ, Carter MG, Piao YL, Sharov AA, Ko MSH, Brown DD. Gene expression changes at metamorphosis induced by thyroid hormone in *Xenopus laevis* tadpoles. *Developmental Biology* 2006;291:342–355. [PubMed: 16458881]
- De Groef B, Grommen SVH, Darras V. The chicken embryo as a model for developmental endocrinology: Development of the thyrotropic, corticotropic, and somatotrophic axes. *Molecular and Cellular Endocrinology* 2008;293:17–24. [PubMed: 18619516]
- Denver, RJ. Endocrinology of complex life cycles: Amphibians. In: Pfaff, AAD.; Etgen, A.; Fahrbach, S.; Moss, R.; Rubin, R., editors. *Hormones, Brain and Behavior*. Academic Press, Inc; San Diego: 2009a.
- Denver RJ. Stress hormones mediate environment-genotype interactions during amphibian development. *General & Comparative Endocrinology* 2009b;164:20–31. [PubMed: 19393659]
- Denver RJ, Hu F, Scanlan TS, Furlow JD. Thyroid hormone receptor subtype specificity for hormone-dependent neurogenesis in *Xenopus laevis*. *Developmental Biology* 2009;326:155–168. [PubMed: 19056375]
- Denver RJ, Pavgi S, Shi YB. Thyroid hormone-dependent gene expression program for *Xenopus* neural development. *Journal of Biological Chemistry* 1997;272:8179–8188. [PubMed: 9079635]
- Dobner PR, Kawasaki ES, Yu LY, Bancroft FC. Thyroid or glucocorticoid hormone induces pre-growth hormone messenger RNA and its probable nuclear precursor in rat pituitary cells. *Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences* 1981;78:2230–2234.
- Frieden, E. The dual role of thyroid hormones in vertebrate development and calorogenesis. In: Gilbert, L.; Frieden, E., editors. *Metamorphosis: A problem in developmental biology*. Plenum Press; New York: 1981. p. 545–564.
- Frieden E, Naile B. Biochemistry of amphibian metamorphosis. 1. Enhancement of induced metamorphosis by glucocorticoids. *Science* 1955;121:37–38. [PubMed: 13225738]
- Furlow JD, Kanamori A. The transcription factor basic transcription element-binding protein 1 is a direct thyroid hormone response gene in the frog *Xenopus laevis*. *Endocrinology* 2002;143:3295–3305. [PubMed: 12193541]
- Furlow JD, Neff ES. A developmental switch induced by thyroid hormone: *Xenopus laevis* metamorphosis. *Trends in Endocrinology and Metabolism* 2006;17:38–45.
- Galton VA. Mechanisms underlying the acceleration of thyroid hormone-induced tadpole metamorphosis by corticosterone. *Endocrinology* 1990;127:2997–3002. [PubMed: 2249638]

- Glennemeier KA, Denver RJ. Small changes in whole-body corticosterone content affect larval *Rana pipiens* fitness components. *General and Comparative Endocrinology* 2002;127:16–25. [PubMed: 12161197]
- Gray KM, Janssens PA. Gonadal hormones inhibit the induction of metamorphosis by thyroid hormones in *Xenopus laevis* tadpoles *in vivo*, but not *in vitro*. *General and Comparative Endocrinology* 1990;77:202–211. [PubMed: 2307343]
- Hemming FJ, Aubert ML, Dubois PM. Differentiation of fetal rat somatotropes in vitro - Effects of cortisol, 3,5,3'-triiodothyronine, and glucagon, a light microscopic and radioimmunological study. *Endocrinology* 1988;123:1230–1236. [PubMed: 3042369]
- Hogan NS, Crump KL, Duarte P, Lean DRS, Trudeau VL. Hormone cross-regulation in the tadpole brain: Developmental expression profiles and effect of T3 exposure on thyroid hormone- and estrogen-responsive genes in *Rana pipiens*. *General and Comparative Endocrinology* 2007;154:5–15. [PubMed: 17400220]
- Hoopfer ED, Huang LY, Denver RJ. Basic transcription element binding protein is a thyroid hormone-regulated transcription factor expressed during metamorphosis in *Xenopus laevis*. *Development Growth & Differentiation* 2002;44:365–381.
- Hu F, Crespi EJ, Denver RJ. Programming neuroendocrine stress axis activity by exposure to glucocorticoids during postembryonic development of the frog *Xenopus laevis*. *Endocrinology* 2008;149:5470–5481. [PubMed: 18653715]
- Iwamuro S, Tata JR. Contrasting patterns of expression of thyroid hormone and retinoid X receptor genes during hormonal manipulation of *Xenopus* tadpole tail regression in culture. *Mol Cell Endocrinol* 1995;113:235–43. [PubMed: 8674831]
- Kanamori A, Brown DD. The regulation of thyroid hormone receptor beta genes by thyroid hormone in *Xenopus laevis*. *J Biol Chem* 1992;267:739–45. [PubMed: 1730665]
- Kikuyama S, Kawamura K, Tanaka S, Yamamoto K. Aspects of amphibian metamorphosis: hormonal control. *International Review of Cytology* 1993;145:105–148. [PubMed: 8500980]
- Kikuyama S, Niki K, Mayumi M, Kawamura K. Retardation of thyroxine-induced metamorphosis by amphenone B in toad tadpoles. *Endocrinologia Japonica* 1982;29:659–662. [PubMed: 6221924]
- Kikuyama S, Niki K, Mayumi M, Shibayama R, Nishikawa M, Shintake N. Studies on corticoid action on the toad tadpole tail *in vitro*. *General and Comparative Endocrinology* 1983;52:395–399. [PubMed: 6421652]
- Krain LP, Denver RJ. Developmental expression and hormonal regulation of glucocorticoid and thyroid hormone receptors during metamorphosis in *Xenopus laevis*. *Journal of Endocrinology* 2004;181:91–104. [PubMed: 15072570]
- Leers J, Steiner C, Renkawitz R, Muller M. A thyroid hormone receptor-dependent glucocorticoid induction. *Mol Endocrinol* 1994;8:440–7. [PubMed: 8052265]
- Leloup J, Buscaglia M. Triiodothyronine, hormone of amphibian metamorphosis. *Comptes Rendus Hebdomadaires Des Seances De L Academie Des Sciences Serie D* 1977;284:2261–2263.
- Lindsay RH, Buettner L, Wimberly N, Pittman JA. Effects of thyroxine analogs on isolated tadpole tail tips. *General and Comparative Endocrinology* 1967;9:416–421. [PubMed: 4170365]
- Liu LX, Dean CE, Porter TE. Thyroid hormones interact with glucocorticoids to affect somatotroph abundance in chicken embryonic pituitary cells *in vitro*. *Endocrinology* 2003;144:3836–3841. [PubMed: 12933655]
- Lorenz C, Opitz R, Lutz I, Kloas W. Corticosteroids disrupt amphibian metamorphosis by complex modes of action including increased prolactin expression. *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 2009;150:314–321.
- Mangelsdorf DJ, Thummel C, Beato M, Herrlick P, Schutz G, Umesono K, Blumberg B, Kastner P, Mark M, Chambon P, Evans RM. The nuclear receptor superfamily. The second decade. *Cell* 1995;83:835–839. [PubMed: 8521507]
- Manzon RG, Denver RJ. Regulation of pituitary thyrotropin gene expression during *Xenopus* metamorphosis: negative feedback is functional throughout metamorphosis. *Journal of Endocrinology* 2004;182:273–285. [PubMed: 15283688]

- Martial JA, Baxter JD, Goodman HM, Seeburg PH. Regulation of growth hormone messenger RNA by thyroid and glucocorticoid hormones. *Proceedings of the National Academy of Sciences of the United States of America* 1977;74:1816–1820. [PubMed: 266704]
- Menjo M, Murata Y, Fujii T, Nimura Y, Seo H. Effects of thyroid and glucocorticoid on the level of messenger ribonucleic acid for iodothyronine type-I 5'-deiodinase in rat primary hepatocyte cultures grown as spheroids. *Endocrinology* 1993;133:2984–2990. [PubMed: 8243326]
- Molero C, Benito M, Lorenzo M. Regulation of malic enzyme gene expression by nutrients, hormones, and growth factors in fetal hepatocyte primary cultures. *Journal of Cellular Physiology* 1993;155:197–203. [PubMed: 8468366]
- Montesinos MM, Pellizas CG, Velez ML, Susperreguy S, Masini-Repiso AM, Coleoni AH. Thyroid hormone receptor beta(1) gene expression is increased by Dexamethasone at transcriptional level in rat liver. *Life Sciences* 2006;78:2584–2594. [PubMed: 16330054]
- Nieuwkoop, PD.; Faber, J. *Normal Table of *Xenopus laevis** Daudin. North Holland Publishers; Amsterdam: 1956.
- Niki K, Yoshizato K, Kikuyama S. Augmentation of nuclear binding capacity for triiodothyronine by aldosterone in tadpole tail. *Proceedings of the Japan Academy Series B-Physical and Biological Sciences* 1981;57:271–275.
- Nogami H, Yokose T, Tachibana T. Regulation of growth hormone expression in fetal rat pituitary gland by thyroid or glucocorticoid hormone. *American Journal of Physiology-Endocrinology and Metabolism* 1995;268:E262–E267.
- Ranjan M, Wong J, Shi Y-B. Transcriptional repression of *Xenopus* TR β gene is mediated by a thyroid hormone response element located near the start site. *J Biol Chem* 1994;269:24699–24705. [PubMed: 7929143]
- Relyea RA. Getting out alive: how predators affect the decision to metamorphose. *Oecologia* 2007;152:389–400. [PubMed: 17356812]
- Robinson H, Chaffee S, Galton VA. Sensitivity of *Xenopus laevis* tadpole tail tissue to action of thyroid hormones. *General and Comparative Endocrinology* 1977;32:179–186. [PubMed: 892410]
- Rosenkilde P. Thyroid hormone synthesis in metamorphosing and adult *Xenopus laevis* Gen. *Comp Endocrinol* 1978;34:95–96.
- Samuels HH, Horwitz ZD, Stanley F, Casanova J, Shapiro LE. Thyroid hormone controls glucocorticoid action in cultured GH1 cells. *Nature* 1977;268:254–257. [PubMed: 196206]
- Samuels HH, Stanley F, Casanova J. Depletion of L-3,5,3'-triiodothyronine and L-thyroxine in euthyroid calf serum for use in cell-culture studies of thyroid hormone. *Endocrinology* 1979;105:80–85. [PubMed: 446419]
- Shapiro LE, Samuels HH, Yaffe BM. Thyroid and glucocorticoid hormones synergistically control growth hormone messenger RNA in cultured GH1 cells. *Proceedings of the National Academy of Sciences of the United States of America* 1978;75:45–49. [PubMed: 272661]
- Shi, YB. *From Morphology to Molecular Biology*. Wiley-Liss; New York: 2000. *Amphibian Metamorphosis*.
- Shintani N, Nohira T, Hikosaka A, Kawahara A. Tissue-specific regulation of type III iodothyronine 5-deiodinase gene expression mediates the effects of prolactin and growth hormone in *Xenopus* metamorphosis. *Development Growth & Differentiation* 2002;44:327–335.
- St Germain DL, Galton VA, Hernandez A. Minireview: Defining the roles of the iodothyronine deiodinases: current concepts and challenges. *Endocrinology* 2009;150:1097–107. [PubMed: 19179439]
- St Germain DL, Schwartzman RA, Croteau W, Kanamori A, Wang Z, Brown DD, Galton VA. A thyroid hormone-regulated gene in *Xenopus laevis* encodes a type III iodothyronine 5-deiodinase [published erratum appears in *Proc Natl Acad Sci U S A* 1994 Nov 8;91(23):11282]. *Proceedings of the National Academy of Sciences of the United States of America* 1994;91:7767–71. [PubMed: 8052658]
- Suzuki RM, Kikuyama S. Corticoids augment nuclear binding capacity for triiodothyronine in bullfrog tadpole tail fins. *General and Comparative Endocrinology* 1983;52:272–278. [PubMed: 6197340]
- Tata JR. *Amphibian metamorphosis as a model for the developmental actions of thyroid hormone*. *Molecular and Cellular Endocrinology* 2006;246:10–20. [PubMed: 16413959]

- Trentin AG. Thyroid hormone and astrocyte morphogenesis. *Journal of Endocrinology* 2006;189:189–197. [PubMed: 16648287]
- Wahlborg A, Bright C, Frieden E. Activity of some new triiodothyronine analogs in tadpole. *Endocrinology* 1964;75:561–564. [PubMed: 14211899]
- Wang Z, Brown DD. Thyroid hormone-induced gene expression program for amphibian tail resorption. *Journal of Biological Chemistry* 1993;268:16270–8. [PubMed: 8344914]
- White, BA.; Nicoll, CS. Hormonal control of amphibian metamorphosis. In: Frieden, LIGaE, editor. *Metamorphosis: A problem in developmental biology*. Plenum; New York: 1981. p. 363-396.
- Yamaguchi S, Murata Y, Nagaya T, Hayashi Y, Ohmori S, Nimura Y, Seo H. Glucocorticoids increase retinoid-X receptor alpha (RXR alpha) expression and enhance thyroid hormone action in primary cultured rat hepatocytes. *Journal of Molecular Endocrinology* 1999;22:81–90. [PubMed: 9924183]
- Yao M, Hu F, Denver RJ. Distribution and corticosteroid regulation of glucocorticoid receptor in the brain of *Xenopus laevis*. *Journal of Comparative Neurology* 2008;508:967–982. [PubMed: 18399546]

Fig. 1.

Corticosterone (CORT) synergizes with 3,5,3' triiodothyronine (T₃) to cause regression of tadpole tail explants. Tails were harvested from NF stage 52–54 *X. laevis* tadpoles and cultured for one week (see Materials and Methods). The medium was replenished and hormones were added at the indicated doses every 12 hr. Tails were imaged and area calculated daily, and tails were dried in an oven and final dry weight measured at the end of the experiment. **A.** Percent initial tail area over the 7 day culture period. Statistical analyses were based on the actual tail areas (ANOVA; n=7/treatment). **B.** Final dry weight of tadpole tail explants. Bars represent means+SEM. Data were analyzed by one-way ANOVA and letters indicate significant differences between the group means based on Fisher's LSD

($P < 0.05$; $n = 7$ /treatment). Representative images of tails from each treatment group at the 7 day time point are shown below the graphs.

Fig. 2.

Corticosterone (CORT) synergizes with 3,5,3' triiodothyronine (T₃) to induce thyroid hormone receptor beta (TR β) mRNA in tadpole tail explants. Tails were harvested from NF stage 52–54 *X. laevis* tadpoles and cultured for one week (see Materials and Methods). The medium was replenished and hormones were added at the indicated doses every 12 hr. Three tails were harvested from each group at 48 hr (**A**) and at 120 hr (**B**) after initiation of hormone treatments and analyzed for TR β mRNA levels by Northern blotting and densitometry of autoradiograms. The data shown in the graph are the percent of maximal expression, but statistical analyses were calculated from band densities for each gene normalized to rpL8. The data were first log₁₀-transformed before analysis by ANOVA, and letters indicate significant differences between the group means based on Fisher's LSD

($P < 0.05$). The dashed lines indicate the control level of expression. The black bars represent the estimated additive responses to T_3 or CORT alone at the doses indicated.

Fig. 3.

Corticosterone (CORT) synergizes with 3,5,3' triiodothyronine (T₃) to induce deiodinase type 2 (D2) mRNA in tadpole tail explants. Tails were harvested from NF stage 52–54 *X. laevis* tadpoles and cultured for one week (see Materials and Methods). The medium was replenished and hormones were added at the indicated doses every 12 hr. Three tails were harvested from each group at 48 hr (**A**) and at 120 hr (**B**) after initiation of hormone treatments and analyzed for TR β mRNA levels by Northern blotting and densitometry of autoradiograms. Shown are representative Northern blots for both time points. Densitometry could not be conducted because the controls were undetectable.

Fig. 4.

Corticosterone (CORT) synergizes with 3,5,3' triiodothyronine (T₃) to induce thyroid hormone receptor alpha (TR α ; **A**) and TR β (**B**) mRNAs in tail from tadpoles treated with hormones *in vivo*. Nieuwkoop-Faber stage 52 *S. silurana* tadpoles were treated with T₃ and CORT at the doses indicated for 24 hr before sacrifice as described in the Materials and Methods. Gene expression was analyzed by RTqPCR using Taqman assays. The data are given as ratios of gene expression to rpL8; the normalized gene expression data were log₁₀-transformed before statistical analysis using ANOVA. The letters indicate significant differences between the group means based on Fisher's LSD ($P < 0.05$; $n = 5$ /treatment). The black bars represent the estimated additive responses to T₃ or CORT alone at the doses indicated.

Fig. 5. Effects of corticosterone (CORT) and 3,5,3' triiodothyronine (T₃) alone or in combination on expression of deiodinase type 2 (D2; **A**) and type 3 (D3; **B**) mRNAs in tail from tadpoles treated with hormones *in vivo*. Nieuwkoop-Faber stage 52 *S. silurana* tadpoles were treated with T₃ and CORT at the doses indicated for 24 hr before sacrifice as described in the Materials and Methods. Gene expression was analyzed by RTqPCR using Taqman assays. The data are given as ratios of gene expression to rpl8; the normalized gene expression data were log₁₀-transformed before statistical analysis using ANOVA. The letters indicate significant differences between the group means based on Fisher's LSD ($P < 0.05$; $n = 5$).

treatment). The black bars represent the estimated additive responses to T₃ or CORT alone at the doses indicated.

Fig. 6.

Effects of corticosterone (CORT) and 3,5,3' triiodothyronine (T₃) alone or in combination on expression of glucocorticoid receptor (GR; **A**) and mineralocorticoid receptor (MR; **B**) mRNAs in tail from tadpoles treated with hormones *in vivo*. Nieuwkoop-Faber stage 52 *S. silurana* tadpoles were treated with T₃ and CORT at the doses indicated for 24 hr before sacrifice as described in the Materials and Methods. Gene expression was analyzed by RTqPCR using Taqman assays. The data are given as ratios of gene expression to rpL8; the normalized gene expression data were log₁₀-transformed before statistical analysis using ANOVA. The letters indicate significant differences between the group means based on

Fisher's LSD ($P < 0.05$; $n = 5/\text{treatment}$). The black bars represent the estimated additive responses to T_3 or CORT alone at the doses indicated.

Fig. 7. Effects of corticosterone (CORT) and 3,5,3' triiodothyronine (T₃) alone or in combination on expression of thyroid hormone receptor alpha (TR α), TR β , deiodinase type 2 (D2), D3, glucocorticoid receptor (GR) and mineralocorticoid receptor (MR) mRNAs in the fibroblast-derived *X. laevis* cell line XTC-2. Cells were treated with or without hormones for 4 hr before harvest and RNA isolation as described in the Materials and Methods. Gene expression was analyzed by RTqPCR using Taqman assays. The data are given as ratios of gene expression to rpL8; the normalized gene expression data were log₁₀-transformed before statistical analysis using ANOVA. The letters indicate significant differences between the group means based on Fisher's LSD (P < 0.05; n = 5/treatment). The black bar on the TR β graph represents the estimated additive responses to T₃ or CORT alone at the doses indicated.

Fig. 8.

Effects of corticosterone (CORT) and 3,5,3' triiodothyronine (T₃) alone or in combination on expression of thyroid hormone receptor beta (TRβ) and glucocorticoid receptor (GR) mRNAs in the *X. laevis* tadpole tail myoblast-derived cell line XLT-15. Cells were treated with or without hormones for 14 or 24 hr before harvest and RNA isolation as described in the Materials and Methods. Gene expression was analyzed by semi-quantitative RTPCR. The data are given as ratios of gene expression to rpl8; the normalized gene expression data were log₁₀-transformed before statistical analysis using ANOVA. The letters indicate significant differences between the group means based on Fisher's LSD ($P < 0.05$; $n = 6$ /treatment). The black bars represent the estimated additive responses to T₃ or CORT alone at the doses indicated.

Table 1

Primers and probes used for semi-quantitative and quantitative real time RT-PCR analysis of gene expression in frog tissues and cells.

Primers for semi-quantitative RT-PCR – <i>X. laevis</i> genes						
Gene	Primer (sense above, antisense below; 5' → 3')	PCR product (bp)	Annealing temp (°C)	# cycles	GenBank accession #	
<i>TrβA</i>	GGCAACAGACTTGGTTTGG AAGTCCACTTTTCCACCCTC	275	57	36	M35359	
<i>D2</i>	AGCCATAAATAGCTTGCCCTGTG GGAACCTTAACCTTGGGACTGC	278	57	34	AF354707	
<i>GR</i>	TCA CCT GGG ACT AGC ATT GCA GAA AGC TGC CAC AAG TCA GTA CTC CAT	256	59.9	34	X72211	
<i>ηL8</i>	CACAGAAAGGTGCTGCTAAG CAGGATGGGTTTGTCAATACG	477	63	34	U00920	

Primers and probes for quantitative RT-PCR – <i>X. laevis</i> genes		
Gene	Primer (sense above, antisense below; 5' → 3')	TaqMan probe
<i>TrαA</i>	GGCAGCCACTGGAACA GGGAGACTGCCCGATATCTTC	CGTAGGAAGTTTCTGCC
<i>TrβA</i>	GGAAGCCACTGGAACACAGAAAA CAITTAACATGGGAGCTTGTCCTCAA	AAAAATTTTGGCCAGAGGAC
<i>D2</i>	AAGGGCTGCGCTGTGTGT GTCCCAGCTTAACCTGCTGTAG	AACAGCTTCTCTATTGGATG
<i>D3</i>	ACCGTGGTTGTCAATTTCG TGCAGCGGAGCCATGAA	CTGCACCTGACCCC
<i>GR</i>	CTGGCAGCGCTTTTACCAA ATTCTCAGCCACCTCATGCAAT	TGACAAAGCTATTGGACTC
<i>MR</i>	ATGGTTCAAGGTGGTGAATGG GGTCTCCAGAGGCAAAATTCT	CAAAAGTGATTCCAGGATT
<i>ηL8</i>	TTTGCTGAAAGAAATGGCTACATC CACGGCTGGATCATGGA	AGGGTATTGTGAAAGACA

Primers and probes for quantitative RT-PCR – <i>S. tropicalis</i> genes		
Gene	Primer (sense above, antisense below; 5' → 3')	TaqMan probe
<i>TrαA</i>	GGCAGCCACTGGAACA GGGAGACTGCCCGATATCTTC	CGTAGGAAGTTTCTGCC
<i>TrβA</i>	GGAAGCCACTGGAACACAGAAAA CAITTAACATGGGAGCTTGTCCTCAA	AAAAATTTTGGCCAGAGGAC

Primers and probes for quantitative RT-PCR – <i>S. tropicalis</i> genes		
Gene	Primer (sense above, antisense below; 5' → 3')	TaqMan probe
<i>D2</i>	AAGGGCTGCGCTGTGTGT GTCCCAAGCTTAACTGCTTGTAG	AACAGCTTCCTATTGGATG
<i>D3</i>	ACCGCTGGTGTCAATTTCG TGCAGGCGAGCCATGAA	CTGCACCTGACCCCC
<i>GR</i>	CTGGCAGGGTTTTTATCAA ATTTTCAGCCACCTCATGCAT	TGACAAAAGCTATTGGACTC
<i>MR</i>	ATGGTTCAGGTGGTGAAATGG GGTCCTCTAGAGGCAAAATTCCT	CAAAAGTGATTCACAGGATT
<i>rpL8</i>	TTTGCTGAAAGAAAATGGGTACATC CACGGCCTGGATCATGGA	AGGGTATTGTGAAAAGACA