Published in final edited form as:

Org Lett. 2010 June 18; 12(12): 2884–2887. doi:10.1021/ol101040p.

Intramolecular Fe(II)-Catalyzed N–O or N–N Bond Formation from Aryl Azides

Benjamin J. Stokes, Carl V. Vogel, Linda K. Urnezis, Minjie Pan, and Tom G. Driver*

Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, Illinois, 60607-7061

Abstract

Iron(II) bromide catalyzes the transformation of aryl- and vinyl azides with ketone- or methyl oxime substituents into 2,1-benzisoxazoles, indazoles or pyrazoles through the formation of an N–O or N–N bond. This transformation tolerates a variety of different functional groups to facilitate access to a range of benzisoxazoles or indazoles. The unreactivity of the *Z*-methyloxime indicates that *N*-heterocycle formation occurs through a nucleophilic attack of the ketone or oxime onto an activated planar iron azide complex.

Despite the prevalence of nitrogen–heteroatom bonds in biologically active *N*-heterocycles, methods that form these bonds remain rare. 2,1-Benzisoxazoles^{1,2} or indazoles^{3,4} typically originate from starting materials with pre-existing N–O or N–N bonds such as oximes, nitrile oxides, or hydrazines. Direct construction of the N–O⁵ or N–N⁶ bond provides an attractive alternative approach to these *N*-heterocycles. Surprisingly, there are no prior examples of transition metal-catalyzed routes to these bonds even though such complexes promote the oxidation^{7,8} or imination⁹ of sulfides. Since we established that benzimidazoles could be produced from *N*-aryl imines with *ortho*-azido substituents upon exposure to FeBr₂ (Scheme 1),¹⁰ we were curious if nitrogen–heteroatom bond formation could be achieved by transposing the *ortho*-heteroatom substituent from the α-position to the β-position. While thermolysis of **3** affords the new nitrogen–heteroatom bond, ^{5a–b}6a–c the high temperature limits the functional group tolerated in the reaction. Herein, we report that iron(II) bromide catalyzes the formation of N–O or N–N bonds to transform azides **1** into 2,1-benzisoxazoles **4** or pyrazoles **5** under markedly benign conditions.

To achieve transition metal-catalyzed nitrogen–heteroatom bond formation, the reactivity of 2-azidobenzophenone toward a variety of transition metal complexes was investigated (Table 1). This aryl azide is available in one step from commercially available 2-aminobenzophenone.¹¹ While a plethora of methods have been reported to generate an electrophilic nitrogen atom from aryl azides,¹² we found that only iron salts promoted 2,1-benzisoxazole **7a** formation at significantly reduced temperatures. Screening other metal complexes known to decompose azides—including Cu(I)- and Cu(II) salts—did not result in nitrogen–oxygen bond formation.¹³ Our data suggests that the oxidation state of iron

tgd@uic.edu.

Supporting Information Available Complete experimental procedures, spectroscopic and analytical data for the products (PDF) is available free of charge via the Internet at <http://pubs.acs.org>.

affected the efficiency of the reaction because lower conversions were observed when more Lewis acidic iron(III) salts were used. If the reaction temperature was increased to 40 °C, the catalyst loading could be reduced to 5 mol %. The low catalyst loading contrasted with benzimidazole formation, which required 30 mol % FeBr₂ for satisfactory conversions.¹⁰ Molecular sieves, while not required, were added to eliminate any adventitious water.

With these optimized conditions, the scope and limitations of iron(II)-catalyzed N–O bond formation was examined (Table 2).¹⁴ 2,1-Benzisoxazole formation depended on the electronic nature of the azide. While the reaction tolerated R¹- or R²-substituents, strong electron-withdrawing groups are incompatible with the reaction conditions (entries 1 – 7). The effect of the R³-substituent on the reaction was briefly surveyed. 2,1-Benzisoxazoles were made from substrates bearing aryl- or alkyl R³-groups (entries 8 – 14). Our data suggests that a carbon R³-substituent is necessary to achieve N–O bond formation because aldehydes, such as **6n**, did not react. Varying the electronic nature of *o*-azidobenzaldehyde **6n** (R¹ = OMe or CF₃) did not result in benzisoxazole formation.

(1)

To determine if our method could be used to form N–N bonds, aryl azide **8**, bearing a β-nitrogen atom in the tether, was screened (eq 1). In addition to testing the possibility of N–N bond formation, this substrate would also allow an examination of the stereochemical requirements of the reaction. When a mixture of **8** (25:75, *E:Z*) was exposed to FeBr₂, the *E*-isomer reacted quantitatively to form 2*H*-indazole **9** while *Z*-**8** was entirely recovered from the reaction mixture.¹⁵ Isomerization of *Z*-**8** was never observed, even when the reaction conditions were substantially altered.¹⁶

Iron(II)-catalyzed N–N bond formation from *E*-methyl oximes **10** exhibited a broader substrate scope than our benzisoxazole study (Table 3).¹⁷ Biaryl ketoximes (**10a** – **10f**) bearing either electron-donating or electron-withdrawing R¹- and R²-substituents were cleanly transformed into **11** (entries 1 – 6). The reaction also tolerated aryl-, alkyl-, and hydrogen R⁴-substituents (entries 7 – 11). As with **7m**, indazole **11j**, which contained a secondary alkyl substituent, isomerized to the 1*H*-aromatic product upon exposure to SiO₂ (entry 10). The identity of the imine nitrogen substituent affected the reaction outcome. While methyl ketoximes were competent substrates, no reaction was observed for ketoximes. The reaction was not limited to aryl azide substrates: trisubstituted pyrazoles **11m** – **11p** could be accessed from vinyl azides upon exposure to 5 mol % of FeBr₂ (entries 12 – 16). Vinyl azides **10m** – **10p** are readily synthesized from the cycloalkanone via a three-step procedure (eq 2).¹⁸

(2)

The reactivity of diazides **10q** and **10r** showcases the mildness of our Fe(II) process (Scheme 2). Whereas thermolysis of either diazide produced a thick tar containing trace

amounts indazole (<5%) and aldoximine, exposure of diazide **10q** or **10r** to FeBr₂ cleanly afforded the 4- or 6-azido-substituted indazole. In contrast to the thermal process, the second azide group did not decompose in our Fe(II)-catalyzed process.

N-Heterocycles **7** or **11** can be elaborated into other synthetically valuable products (Scheme 3). SmI₂-mediated reduction of the N–O bond in benzisoxazole **7a** in the presence of acetophenone or acetone formed quinolines **12a** and **12b**.¹⁹ Palladium-catalyzed hydrogenolysis of the N–OMe bond present in **11q** and **11p** produced 1*H*-indazole **13** and 2*H*-pyrazole **14**.²⁰ The vanilloid receptor (TRPV1) antagonist, ABT-102,²¹ contains 1*H*-indazole **13**. Abbott Pharmaceuticals developed ABT-102 as a clinical candidate for the treatment of chronic pain.

While several mechanisms could account for *N*-heterocycle formation,²² the reactivity differences between the *E*- and *Z*-oxime isomers of **10** suggest that N–X bond formation occurs through a planar reactive intermediate (Scheme 4).²³ Because the corresponding aniline was unreactive, we assign this reactive intermediate to be an iron(II) azide complex. This intermediate (**15**) could be produced from *E*-**10** by coordination of the iron(II) catalyst to the terminal *N*-atom of the azide.²⁴ Planarization then triggers a nucleophilic attack of the oxime on the activated azide to form the N–N bond.^{22,25} Extrusion of N₂ from **16** followed by dissociation of the iron catalyst would provide the 2*H*-indazole product. The origin of the *Z*-isomer's inertia can be inferred from examination of the iron azide complex **18**: in addition to the destabilizing steric interactions produced upon planarization of **18**, the lone pair of electrons on the oxime are oriented in the wrong direction for N–N bond formation.

In conclusion, we have demonstrated that iron(II) bromide readily forms N–O or N–N bonds from a range of *ortho*-azide-substituted aryl ketones or *E*-methyl oximes. Future experiments will determine if a contiguous π -system is required for Fe(II)-catalyzed azide decomposition and if any of the reactive intermediates can be trapped by dipolarophiles to produce additional C–C or C–N bonds.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the National Institutes of Health NIGMS (R01GM084945), Petroleum Research Fund administered by the American Chemical Society (46850-G), and the University of Illinois at Chicago for their generous support. We also thank Prof. L. Anderson (UIC) for insightful discussions and Mr. Furong Sun (UIUC) for mass spectrometry data.

References

1. Recent reviews, see: (a) Elguero, J.; Katrizky, AR.; Rees, CW., editors. Comprehensive Heterocyclic Chemistry. Vol. 4. New York: Pergamon Press; 1984. p. 167(indazoles) (b) Stadlbauer W. Sci. Synth. 2002; 12:243. (indazoles). (c) Smalley RK. Sci. Synth. 2002; 11:337. (benzisoxazoles).
2. (a) Davis RB, Pizzini LC. J. Org. Chem. 1960; 25:1884. (b) Musso H, Schröder H. Chem. Ber. 1965; 98:1562. (b) Wróbel Z. Synthesis. 1997:753. (c) Boduszek B, Halama A, Zon J. Tetrahedron. 1997; 53:11399.
3. Indazoles: (a) Jacobsen P, Huber L. Ber. Chem. Ges. 1908; 41:660. (b) Rüchardt C, Hassmann V. Liebigs Ann. Chem. 1980:908.
4. For recent examples, see: (a) Song JJ, Yee YK. Org. Lett. 2000; 2:519. [PubMed: 10814366] (b) Lukin K, Hsu MC, Fernando D, Leanna MR. J. Org. Chem. 2006; 71:8166. [PubMed: 17025307] (c) Vina D, del Olmo E, Lopez-Perez JL, San Feliciano A. Org. Lett. 2007; 9:525. [PubMed: 17025307]

- 17249803] (d) Jin T, Yamamoto Y. *Angew. Chem., Int. Ed.* 2007; 46:3323. (e) Liu Z, Shi F, Martinez PDG, Raminelli C, Larock RC. *J. Org. Chem.* 2008; 73:219. [PubMed: 18067316]
5. (a) Smith PAS, Brown BB, Putney RK, Reinisch PF. *J. Am. Chem. Soc.* 1953; 75:6335. (b) Boulton AJ, Ghosh PB, Katritzky AR. *Angew. Chem., Int. Ed. Engl.* 1964; 3:693. (c) Prakash O, Saini RK, Singh SP, Varma RS. *Tetrahedron Lett.* 1997; 38:3147.
6. (a) Krbeček L, Takimoto H. *J. Org. Chem.* 1964; 29:1150. (b) Cadogan JIG, Cameron-Wood M, Mackie RK, Searle RJG. *J. Chem. Soc.* 1965:4831. (c) Takada K, Thoe KW, Boulton AJ. *J. Org. Chem.* 1982; 47:4323. (d) Counciller CM, Eichman CC, Wray BC, Stambuli JP. *Org. Lett.* 2008; 10:1021. [PubMed: 18229936]
7. Reviews on asymmetric sulfoxidation: (a) Bolm, C.; Muñoz, K.; Hildebrand, JP. *Comprehensive Asymmetric Catalysis*. Jacobsen, EN.; Pfaltz, A.; Yamamoto, H., editors. Berlin: Springer; 1999. p. 697 (b) Kagan, HB. *Catalytic Asymmetric Synthesis*. Ogima, I., editor. New York: Wiley-VCH; 2000. p. 327 (c) Bolm C. *Coord. Chem. Rev.* 2003; 237:245.
8. For recent reports, see: (a) Legros J, Bolm C. *Angew. Chem., Int. Ed.* 2004; 43:4225. (Fe). (b) Drago C, Caggiano L, Jackson RFW. *Angew. Chem., Int. Ed.* 2005; 44:7221. (V). (c) Yamaguchi T, Matsumoto K, Saito B, Katsuki T. *Angew. Chem., Int. Ed.* 2007; 46:4729. (Al). (d) Egami H, Katsuki T. *J. Am. Chem. Soc.* 2007; 129:8940. (Fe). [PubMed: 17602480]
9. (a) Takada H, Ohe K, Uemura S. *Angew. Chem., Int. Ed.* 1999; 38:1288. (Cu). (b) Bach T, Körber C. *J. Org. Chem.* 2000; 65:2358. (Fe). [PubMed: 10789447] (c) Murakami M, Uchida T, Katsuki T. *Tetrahedron Lett.* 2001; 42:7071. (Ru). (d) Tamura Y, Uchida T, Katsuki T. *Tetrahedron Lett.* 2003; 44:3301. (Ru). (e) Okamura H, Bolm C. *Org. Lett.* 2004; 6:1305. (Rh). [PubMed: 15070323] (f) Mancheño OG, Bolm C. *Chem.—Eur. J.* 2007; 13:6674.
10. Shen M, Driver TG. *Org. Lett.* 2008; 10:3367. [PubMed: 18597473]
11. Many 2-amino-substituted aryl ketones are commercially available. Alternatively, the substrates were synthesized from 2-aminobenzaldehyde derivatives. See the Supporting Information for the details of their synthesis.
12. For reviews, see: (a) Katsuki T. *Chem. Lett.* 2005; 34:1304. (b) Bräse S, Gil C, Knepper K, Zimmermann V. *Angew. Chem., Int. Ed.* 2005; 44:5188. (c) Cenini S, Gallo E, Caselli A, Ragaini F, Fantauzzi S, Piangiolino C. *Coord. Chem. Rev.* 2006; 250:1234.
13. The use of other transition metal complexes resulted in attenuated conversions. Please refer to the Supporting Information for more details.
14. To a mixture of aryl azide **6** (0.1 mmol), 100 % w/w of crushed 4 Å molecular sieves, and metal salt (5 – 10 mol %) in a conical vial with teflon septum was added 0.250 mL of solvent. The resulting mixture was heated and, after 16 h, the heterogeneous mixture was filtered through SiO₂. The filtrate was concentrated *in vacuo*. Purification by MPLC afforded the desired product.
15. Z-Oximes were reported to be unreactive in the metal-free thermal reaction. See: ref. 6c.
16. For a complete description of the isomerization conditions screened, refer to the Supporting Information.
17. *E*-**10** (or a mixture of *E*-, *Z*-**10**), 100 % w/w of crushed 4 Å molecular sieves, and metal salt (5 – 10 mol %) were added to a conical vial, which was sealed with a PTFE septum, followed by the addition of CH₂Cl₂ as solvent. The resulting mixture was heated to 40 °C for 16 h. The heterogeneous mixture was purified by MPLC to afford the product.
18. Tabyaoui B, Aubert T, Farnier M, Guillard R. *Synth. Commun.* 1988; 18:1475. Refer to the Supporting Information for the synthesis of **10m** – **10p**.
19. For a related SmI₂-mediated pyridine synthesis, see: Fan X, Zhang X, Zhang Y. *Heteroatom Chem.* 2005; 16:637.
20. Lower yields of 1*H*-indazoles were obtained when alternative reduction protocols were attempted.
21. (a) Drizin I, Gomtsyan A, Bayburt EK, Schmidt RG, Zheng GZ, Perner RJ, DiDomenico S, Koenig JR, Turner SC, Jinkerson TK, Brown BS, Keddy RG, McDonald HA, Honore P, Wismer CT, Marsh KC, Wetter JM, Polakowski JS, Segreti JA, Jarvis MF, Faltynek CR, Lee C-H. *Bioorg. Med. Chem.* 2006; 14:4740. [PubMed: 16621571] (b) Lukin K, Hsu MC, Chambournier G, Kotecki B, Venkatramani CJ, Leanna MR. *Org. Proc. Res. Dev.* 2007; 11:578.
22. The lack of aniline byproducts suggests that an alternative mechanism involving the formation of iron nitrenoid intermediates is not occurring. For leading references on iron-mediated nitrene

- transfer, see: (a) ref. 9b (b) Bacci JP, Greenman KL, Van Vranken DL. *J. Org. Chem.* 2003; 68:4955. [PubMed: 12790609] (c) Cowley RE, Eckert NA, Elhaik J, Holland PL. *Chem. Commun.* 2009:1760.
23. For leading mechanistic studies of the thermolysis reaction, see: (a) Smith PAS, Budde GF, Chou S-SP. *J. Org. Chem.* 1985; 50:2062. (b) Dyllal LK, Karpa GJ. *Aust. J. Chem.* 1988; 41:1231. (c) Dyllal LK, Holmes AL. *Aust. J. Chem.* 1988; 41:1677.
24. For crystal structures of metal azide complexes, see: (a) Proulx GR, Bergman RG. *J. Am. Chem. Soc.* 1995; 117:6382. (N_γ -coordination). (b) Fickes MG, Davis WM, Cummins CC. *J. Am. Chem. Soc.* 1995; 117:6384. (N_γ -coordination). (c) Dias HVR, Polach SA, Goh S-K, Archibong EF, Marynick DS. *Inorg. Chem.* 2000; 39:3894. (N_α^- and N_γ -coordination). [PubMed: 11196786] (d) Waterman R, Hillhouse GL. *J. Am. Chem. Soc.* 2008; 130:12628. (η^2 -coordination). [PubMed: 18729364]
25. For an alternative mechanism of N–N bond formation through an intramolecular 1,3-dipolar cyclization, see: Hall JH, Behr FE, Reed RL. *J. Am. Chem. Soc.* 1972; 94:4952.

Scheme 1.
Iron(II) Bromide-Catalyzed *N*-Heterocycle Formation

Scheme 2.
Comparison of the Reactivity of Diazides **10**

Sm(II)-Mediated Reduction of N–O Bond ***Hydrogenolysis of N–O Bond***

Scheme 3.
Functionalization of *N*-Heterocyclic Products

Scheme 4.
Potential Mechanism for 2H-Indazole Formation

Table 1

Optimization of Metal-Catalyzed N–O Bond Formation

entry	catalyst	mol %	temp (°C)	yield (%) ^a
1 ^b	none	n.a.	120	92
2	$\text{Rh}_2(\text{O}_2\text{CC}_3\text{F}_7)_4$	5	25	trace
3	CuCl	10	25	no reaction
4	$\text{Cu}(\text{OTf})_2$	10	25	trace
5 ^b	$\text{RuCl}_3 \cdot \text{H}_2\text{O}$	10	25	trace
6	ZnI_2	10	25	no reaction
7	$\text{Fe}(\text{acac})_2$	10	25	8
8	FeCl_3	10	25	20
9 ^e	FeCl_2	10	25	97
10	FeBr_2	10	25	97
11	FeBr_2	5	40	99
12 ^b	FeBr_2	5	40	90

^a As determined using ^1H NMR spectroscopy.^b No molecular sieves were added.

^a After SiO₂ chromatography.

^b 10 mol % of FeBr₂ used.^c 10% aniline formed.

d Isomerization to the 1*H*-benzisoxazole tautomer occurred upon SiO₂ purification.

NIH-PA Author Manuscript

NIH-PA Author Manuscript

NIH-PA Author Manuscript

entry	10	R ¹	R ²	R ³	yield (%) ^a
<div> </div>					
12 ^b	1				99 ^c
13	m				91
14	n				86
15	o				86

entry	10	R ¹	R ²	R ³	yield (%) ^a
16	p				79

^a After SiO₂ chromatography.

^b Reaction performed using an *E*-/*Z*-mixture of **10**.

^c The remainder of the reaction mixture was unreacted **8**.

^d 10 mol % of FeBr₂ used.

^e Isomerization to the 1*H*-benzisoxazole tautomer occurred upon SiO₂ purification.

^f 20 mol % FeBr₂, 25 °C.