Published in final edited form as:

Org Lett. 2010 August 6; 12(15): 3548–3551. doi:10.1021/ol101410g.

Synthetic Access to Bent Polycycles by Cation- π Cyclization

Ryan A. Shenvi and E. J. Corey*

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138

Abstract

The presence of an ether oxygen within a chain undergoing cation–polyene cyclization has a profound influence on the stereochemistry of this important construction, apparently due to nucleophilic participation of oxygen in the cyclization process and formation of an oxonium intermediate, leading to bent fused ring systems.

The carbocation-initiated polycyclization of appropriate polyunsaturated substrates is one of the most powerful molecular constructions in synthetic chemistry. It is also an extremely important biosynthetic process, for instance, in the one-step tetracyclization of (*S*)-2,3-oxidosqualene **1** to a sterol (Figure 1, **1** → **2**). The carbocation- π -cyclization process is also the dominating pathway for the biosynthesis of cyclic terpenoids. Cationic cyclization has served well for the chemical syntheses of many complex terpenoids, e.g., pentacyclosqualene **2a** (an early example), dammarenediol, **2b** lanostenol, **2c** onocerin, **2d** serratenediol, **2e** β -amyrins, and lupeol. **2f,g** However, there is still a large gap in efficiency between biosynthetic polycyclizations and present chemical syntheses. Unlike the enzymic cyclizations which proceed with an efficiency of ca. 99% per ring formed, the chemical cyclizations are, at best, only 70–80% efficient per ring formed. This difference arises partly because the cyclization substrate is held by the enzyme in a prefolded conformational arrangement which selects the proper π -face of a double bond for attack by the propagating carbocation, minimizing the decrease in the entropy of activation of enzymic cyclization.

Although there is no known way to mimic this template-guided cyclization, the entropic problem may be diminished by conducting cyclization at the lowest possible temperature (to minimize the (positive) $T\Delta S^\ddagger$ component of the free energy of activation ($\Delta G^\ddagger$)) in CH_2Cl_2 with MeAlCl_2 as catalyst. Functional groups more basic than epoxide are not tolerated. The studies described herein were undertaken to gain a better understanding of the relationship between substrate structure and π -facial selectivity and to extend the scope of cyclization. One motivation for this study was the observation that the common stereochemical pathway for cyclization, exemplified by the conversion of **3** to the A/B/C/*trans*-*anti*-*trans* product **4**

(Figure 2),³ is not universal since cases such as **5** → **6** are known (Figure 3)⁴ which would seem to involve cations **7** and **8** and a different π -facial selectivity at the second olefinic linkage. This difference in π -facial selectivity at the double bond involved in closure of the second ring is of great interest since it is clearly a branch point in the biosynthetic cyclizations which lead to sterols or plant triterpenes.⁵

Our initial research of the relationship between the structure of the substrate and the stereochemistry of closure of the second ring in epoxide-initiated cation–olefin cyclizations was conducted with simple epoxyfarnesol derivatives. In the discussion which follows, the two modes of bicyclization will be referred to as the 9α -H or 9β -H pathway, as structurally indicated with formulas **9**–**11** (Figure 4). This phase of our work was carried out with CH_2Cl_2 as solvent, and RAlCl_2 or R_2AlCl (3 equiv, e.g., EtAlCl_2) as catalytic Lewis acid at -78°C , conditions which generally are most favorable for epoxide-initiated cationic polycyclization reactions.

When the benzoate of 10,11-epoxyfarnesol was examined under standard conditions for cyclization only monocyclic reaction products were obtained. The most obvious explanation for this result is that the Lewis acid coordinates more strongly with the benzoate carbonyl, and essentially completely (with 3 equiv of EtAlCl_2), and that coordination removes electron density from the terminating double bond which prevents formation of the second ring. Because of this result, we next examined the two bulky silyl ethers of 10,11-epoxyfarnesol **12a** and **12b** (Scheme 1), the assumption being that Lewis acid complexation with the sterically hindered silyloxy oxygen would not be a complication. Indeed, in each case bicyclization was the principal reaction pathway (70–85% isolated yield of **13** and **14** after desilylation and column chromatography on silica gel). For each of these substrates there was a near balance between 9α -H and 9β -H pathways for cyclization, as indicated in Scheme 1. The 9α -H and 9β -H products each consisted of a mixture of one endo- and one exocyclic olefinic species.⁶

The close similarity of ratios for the 9α -H and 9β -H cyclization pathways came as a surprise, given the considerable number of reported examples in which the 9α -H product is strongly preferred. One possible reason for the exceptional behavior of the substrates **12a** and **12b** becomes apparent when the reported sequence **5** → **6** (presumably via **7** and **8**) is recalled. If it were generally true that the 9β -H pathway is favored for reactions that proceed to bicyclic 6/6-fused product via a bicyclic 6/5-fused intermediate, it is logical to explain the formation of the 9β -H product **14** via the oxonium intermediate **15**, which then converts to **16** (Figure 5). To test whether the 9β -H pathway is made more favorable if the incipient ring B is generated via a 5-membered-like precursor, cyclization of the corresponding allylsilane **17** was examined (Scheme 2).

The substrate **17** was synthesized from 9,10-epoxyfarnesyl benzoate by coupling with the reagent derived from dimethylphenylsilyllithium and copper(I) cyanide. Cyclization of **17** under the conditions developed for **12a** and **12b** and subsequent desilylation with TBAF afforded an equimolar mixture of two products which were chromatographically separated after conversion to their 3,5-dibromobenzoates **20** and **21** (chromatography on AgNO_3 -impregnated silica gel). The stereochemistry of the decalin product **19** corresponds to the usual 9α -H cyclization (confirmed by X-ray crystallographic analysis of **21**, mp 144–146 $^\circ\text{C}$), whereas that of the hydrindane **18** corresponds to the π -facial selection of the 9β -H geometry (confirmed by NOE studies of **20**). This result for **18** is consistent with the proposal that cyclization via a 5-membered B-ring structure can involve the opposite π -face selectivity to that which is generally found for 6-membered B-ring formation.

Further studies uncovered a surprising effect of π -bond basicity on the stereochemistry of B-ring formation. The more π -basic tri-*n*-butylstannyl analogue **23** of the allylic silane **17** underwent cyclization under the standard conditions to produce a 1:1 mixture of the two diastereomeric bicyclic structures **24a** and **24b** (Scheme 3). This complete lack of π -face selectivity in the formation of ring B may well be a consequence of a much earlier transition state for ring B closure, i.e., a much longer incipient C–C bond in the transition state.

Behavior similar to that just described for **23** was observed in the cyclization of epoxide **25** which leads to a 1:1 mixture of **26a** and **26b** (Scheme 4). In this case also, there is no π -face selectivity in the formation of ring B, probably for the same reason: a longer C–C bond in the transition state for B-ring closure and a small energy difference between the diastereomeric transition states.⁷

It was found that 9,10-epoxyfarnesyl phenyl ether **27** underwent efficient conversion to the tetracyclic product **28a** in a single step (Scheme 5). Evidently, the phenolic oxygen is sufficiently nonbasic to allow monocoordination of EtAlCl₂ to the more basic epoxide function, in contrast to the benzyl and cinnamyl ethers, which gave essentially no polycyclic products under identical reaction conditions. Of great interest is the fact that cyclization of **27** favored the 9 β -H pathway in preference to the 9 α -H path, since **28a** predominated over the C9 diastereomer **28b** by 2:1 (see the Supporting Information). The major tetracyclic product was isolated by chromatography after conversion to the 3,5-dibromobenzoate (51% overall yield from **27**). The assignment of stereochemistry to **28a** and **28b** was made unambiguously by ¹H NMR measurements at 500 MHz and NOE correlations. It should be noted that decalin cation **29** must undergo a conformational change⁸ from initially formed boatlike conformer **29** to the chair form **30** before closure of ring C to produce tetracycle **28a**.

The contrast between the cyclization pathways **27** $\rightarrow$ **28a** and **3** $\rightarrow$ **4** (Figure 6) provides another indication of the importance of the lone pairs of oxygen in influencing the balance between the 9 α -H and 9 β -H modes of reaction.

Cyclization of the analogous 1-naphthyl ether **31** also proceeded cleanly and afforded the pentacycle **32** as the major product (Scheme 6). A slight increase in the diastereoselectivity of cyclization (9 β -H/9 α -H, 2.4:1) was observed relative to the cyclization of phenyl ether **27** (9 β -H/9 α -H, 2:1).

Additional oxygen substitution on the aryl moiety is also tolerated in the cationic polycyclization process and as such can provide useful reactive centers for elaboration to more complex structures (Scheme 7). The farnesyl ether derived from hydroquinone (**33**) underwent efficient cyclization to tetracycle **35** in good yield and in 2.3:1 excess over the 9 α -H epimer (70% combined yield of diastereomers). Similarly, the phloroglucinol-derived bis-triisopropylsilyl (TIPS) ether **34** underwent cyclization to the corresponding tetracycle **36** with a 9 β -H to 9 α -H ratio of 3:1 (total yield 62%).

We also investigated the cyclization of the difarnesyl 1,5-naphthyl ether **37** (Scheme 8) which was synthesized in one step from 1,5-dihydroxynaphthalene and (*S*)-10,11-epoxyfarnesyl bromide. Treatment of **37** with EtAlCl₂ (6 equiv) in CH₂Cl₂ at –78 °C led to a mixture of diols, which were converted to the corresponding 3,5-dibromobenzoates and purified by chromatography to give octacycle **38** (31% from **37**), the structure of which was confirmed by NMR and single-crystal X-ray analysis.⁸

The unique twisted, rigid octacyclic structure of **38**, prepared in just a few steps from farnesol and commercially available 1,5-dihydroxynaphthalene, would be difficult to make by

other methods.⁹ It further demonstrates the power of cationic polycyclization methodology to produce a wide range of interesting structures.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the NIH for a Postdoctoral Fellowship to R.A.S. and Dr. Shao-Liang Zheng (Harvard) for X-ray diffraction analysis.

References

- (1). Wendt KU, Schulz GE, Corey EJ, Liu DR. *Angew. Chem., Int. Ed.* 2000; 39:2812–2833.
- (2). Corey EJ, Sauers RR. *J. Am. Chem. Soc.* 1957; 79:3925–3926. Corey EJ, Lin SA. *J. Am. Chem. Soc.* 1996; 118:8765–8766. Corey EJ, Lee J, Liu DR. *Tetrahedron Lett.* 1994; 35:9149–9152. Mi Y, Schreiber JV, Corey EJ. *J. Am. Chem. Soc.* 2002; 124:11290–11291. [PubMed: 12236742] Zhang J, Corey EJ. *Org. Lett.* 2001; 3:3215–3216. [PubMed: 11574034] Surendra K, Corey EJ. *J. Am. Chem. Soc.* 2009; 131:13928–13929. [PubMed: 19788328] For a review, see: Yoder RA, Johnston JN. *Chem. Rev.* 2005; 105:4730–4756. [PubMed: 16351060]
- (3). Surendra K, Corey EJ. *J. Am. Chem. Soc.* 2008; 130:8865–8869. [PubMed: 18597440]
- (4). Corey EJ, Roberts BE. *Tetrahedron Lett.* 1997; 38:8921–8924.
- (5). Dev, S.; Nagasampagi, BA. *Triterpenoids*. Vol. 1. CRC Press; Boca Raton: 1986.
- (6). For details of analyses and quantitative data, see the Supporting Information.
- (7). A less likely extreme case can also be visualized in which electron transfer occurs from the π -electron-rich terminating C=C subunit to the monocyclic carbocation (without stereoselectivity) to give a diradical that collapses to the 1:1 mixture of **24a** and **24b**.
- (8). Corey EJ, Wood HB. *J. Am. Chem. Soc.* 1996; 118:11982–11983.
- (9). Corey EJ, Luo G, Lin LS. *Angew. Chem., Int. Ed.* 1998; 37:1126–1128.

Figure 1.
Biosynthetic conversion of (*S*)-2,3-oxidosqualene (1) to protosteryl cation (2).

Figure 2.

Trans-anti-trans cyclization of **3** to **4** producing 9 α -H stereochemistry.

Figure 3.
Cyclization of **5** to **6** through cation **8**, which possesses 9β -H stereochemistry.

Figure 4.

Two modes of cyclization for the formation of ring B: the $9\alpha\text{-H}$ or $9\beta\text{-H}$ pathways.

Scheme 1.

Cyclization of Silyl Ethers **12a** and **12b** To Produce Equimolar Mixtures of 9α-H and 9β-H Products^a

^a Conditions: (1) EtAlCl₂ (3 equiv added slowly), CH₂Cl₂ (0.01 M), -78 °C, 3 h; (2) TBAF (1.5 equiv), THF (0.5 M), 25 °C, 6 h.

Figure 5.

Intermediacy of an oxygen-stabilized secondary cation **15** may explain the formation of 9 β -H products.

Scheme 2.

Cyclization of Silane **17** Giving Rise To Hydrindane **18**, Which Possesses 9 β -H Stereochemistry^a

^a Conditions: (1) EtAlCl_2 (3 equiv added slowly), CH_2Cl_2 (0.01 M), -78°C , 3 h; (2) TBAF (1.5 equiv), THF (0.5 M), 25°C , 6 h; (3) DCC (1.1 equiv), 3,5-dibromobenzoic acid (**22**) (1.1 equiv), DMAP (0.05 equiv), CH_2Cl_2 (0.1 M), 25°C , 1 h.

Scheme 3.
Cyclization of Allylstannane **23** To Form a Mixture of Diastereomers **24a** and **24b**

Scheme 4. Cyclization of Allylsilane **25** To Form a Mixture of Diastereomers **26a** and **26b**

Scheme 5.
Cyclization of Phenyl Ether **27** To Produce Predominantly 9 β -H Product **28a**

Figure 6.
Cyclization of **27** $\rightarrow$ **28a** compared to the cyclization of **3** $\rightarrow$ **4**.

Scheme 6.
Cyclization of **31** To Produce Pentacycle **32**

Scheme 7.

Oxygen Substitution on the Arene Is Tolerated under Cyclization Conditions

Scheme 8.
Bidirectional Cyclization of **37** To Octacycle **38**