Published in final edited form as:

J Abnorm Psychol. 2010 August ; 119(3): 502–512. doi:10.1037/a0019196.

Resting Frontal EEG Asymmetry as an Endophenotype for Depression Risk: Sex-specific Patterns of Frontal Brain Asymmetry

Jennifer L. Stewart¹, Andrew W. Bismark¹, David N. Towers², James A. Coan³, and John J.B. Allen¹

¹University of Arizona

²University of Illinois at Urbana-Champaign

³University of Virginia

Abstract

Resting frontal electroencephalographic (EEG) asymmetry has been hypothesized as a marker of risk for major depressive disorder (MDD) but the extant literature is based predominately on female samples. Resting frontal asymmetry was assessed on 4 occasions within a two-week period in 306 individuals age 18–34 (31% male) with ($n = 143$) and without ($n = 163$) DSM-IV defined lifetime MDD. Lifetime MDD was linked to relatively less left frontal activity for both sexes using a current source density (CSD) reference, findings that were not accounted for solely by current MDD status or current depression severity, suggesting that CSD-referenced EEG asymmetry is a possible endophenotype for depression. In contrast, results for average and linked mastoid references were less consistent, but demonstrated a link between less left frontal activity and current depression severity in women.

Keywords

EEG asymmetry; depression; sex differences; endophenotype

Resting Frontal EEG Asymmetry as an Endophenotype for Depression Risk: Sex-specific Patterns of Frontal Brain Asymmetry

The economic and personal burden of depression is overwhelming, contributing to pervasive and chronic disability in occupational and interpersonal functioning (e.g., Greenberg et al., 2003; Judd et al., 2000), and substantially increased risk for suicide (Kessler, DuPont, Berglund, & Wittchen, 1999). In the quest to reduce this burden, *endophenotypes* (Gottesman & Gould, 2003; Iacono, 1998), measurable endogenous characteristics of an individual that are related to underlying mechanisms conferring risk, may prove useful in identifying a subset of the population at risk for developing depression, and may ultimately

Address correspondence to Jennifer L. Stewart or John J.B. Allen, Department of Psychology, University of Arizona, 1503 E. University Ave., room 312, Tucson, AZ 85721-0068; jlstewar@email.arizona.edu or John.JB.Allen@Arizona.edu..

Publisher's Disclaimer: The following manuscript is the final accepted manuscript. It has not been subjected to the final copyediting, fact-checking, and proofreading required for formal publication. It is not the definitive, publisher-authenticated version. The American Psychological Association and its Council of Editors disclaim any responsibility or liabilities for errors or omissions of this manuscript version, any version derived from this manuscript by NIH, or other third parties. The published version is available at www.apa.org/pubs/journals/abn.

assist in identifying functional mechanisms that may point to new treatments and preventions. A potential promising endophenotype is resting frontal electroencephalographic (EEG) asymmetry.

Depressed individuals appear to be characterized by a pattern of relatively less left than right resting frontal activity (inferred by relatively greater left than right alpha band activity; see Allen, Coan, & Nazarian, 2004), which is thought to reflect reduced approach motivation and decreased responsiveness to reward. Moreover, this pattern distinguishes both symptomatic and remitted depressed individuals from never-depressed individuals, suggesting that frontal EEG asymmetry may be a trait-like liability marker for development of depression (e.g., Allen, Urry, Hitt, & Coan, 2004b; Coan & Allen, 2003, Gotlib, Ranganath, & Rosenfeld, 1998; Henriques & Davidson, 1990, 1991). Consistent with this hypothesis, research has also demonstrated relatively less left frontal activity in children of depressed mothers (e.g., Dawson, Fray, Panagiotides, Osterling, & Hessel, 1997). Studies examining the heritability of frontal EEG asymmetry have determined that this pattern of frontal regional brain activity has a significant genetic component (Anokhin, Heath, & Myers, 2006; Smit, Posthuma, Boomsma, & DeGeus, 2007) but EEG asymmetry may only relate to anxiety and depression risk in young women, not men (Smit et al., 2007), suggesting that further research is needed to explore sex differences in this potential endophenotype for depression.

Psychometric characteristics of resting frontal EEG asymmetry also support a trait-level individual difference, with excellent internal consistency (Towers & Allen, 2009) and modest test-retest reliability in both depressed and non-psychiatric samples (Allen et al., 2004b; Hagemann, Naumann, Thayer, & Bartussek, 2002; Tomarken, Davidson, Wheeler, & Kinney, 1992), and approximately 60% of the variance representing stable trait variance (e.g., Hagemann et al., 2002). Several EEG studies, however, have failed to confirm an association between left frontal hypoactivity and depression (e.g., Bruder et al., 1997; Pizzagalli et al., 2002; Reid, Duke & Allen, 1998), and although some evidence suggests that this pattern of asymmetrical frontal activity remains stable in the midst of fluctuating symptom severity (Allen et al., 2004b), there exists some evidence to the contrary (Debener et al., 2000). Before resting frontal EEG asymmetry can be considered a viable endophenotype for depression risk in a particular subset of individuals, these inconsistencies must be addressed.

Several methodological and sample-specific factors might explain the inconsistent findings, including small patient samples, diagnostic heterogeneity and comorbidity, sexual dimorphism in EEG asymmetry and/or depressive illness, choice of EEG reference, the measure of depression used (e.g., DSM-IV diagnosis versus questionnaire indexing current symptom severity) and the reliability and stability of EEG asymmetry within and across sessions (e.g., Allen et al., 2004a, 2004b; Coan, Allen, & McKnight, 2006; Davidson, 1998; Hagemann, 2004). To date, however, the role of these factors has not been systematically examined due primarily to relatively small patient samples. The present study utilized a large sample of depressed and non-depressed individuals to address these and other factors in order to evaluate the utility of frontal EEG asymmetry as an endophenotype of risk for depression.

Although a recent meta-analysis suggests a moderately robust association exists between depression (and related constructs) and relatively less left frontal activity (Thibodeau, Jorgensen, & Kim, 2006), participants in EEG asymmetry investigations have been predominantly female (Coan & Allen, 2004; Thibodeau et al., 2006). It thus remains to be determined how reliable and stable this pattern is for depressed men, especially given that studies including substantial male samples (not included in the meta-analysis) are

inconsistent, one indicating that male depression is associated with relatively more left frontal activity (Miller et al., 2002), and one demonstrating the opposite pattern (Jacobs & Snyder, 1996). Small patient samples have also prevented within-study comparisons of EEG activity between currently depressed individuals and euthymic individuals with a history of depression (with the exception of Gotlib et al., 1998), a contrast needed to test the assertion that EEG asymmetry is a trait-like marker of depression risk. Conflicting results across studies may also be due to heterogeneity of depressed samples, potentially attributable to types of comorbid anxiety that are associated with different patterns of brain asymmetry than those displayed by non-anxious depressed individuals (e.g., Heller & Nitschke, 1998).

In addition to differences in sample selection and recruitment, EEG-specific methodological issues such as choice of electrode reference may account for differences across studies. Cz has been the most-often utilized reference for asymmetry studies (Coan & Allen, 2003), particularly in infant and child samples, but data from the Cz reference do not correlate highly with that from other references thought to be better suited to capturing regionally specific brain activity (cf. Reid et al., 1998; see Allen et al., 2004a, Hagemann, Naumann, & Thayer, 2001, and Hagemann, 2004 for discussion of additional reference issues). A current source density (CSD) derivation has more recently been advocated as an alternative that reduces the influence of cross-hemisphere and distal volume conduction on the asymmetry score with greater specificity for local electrical sources and sinks (Hagemann, 2004). In simpler terms, the CSD algorithm for a particular electrode location estimates the amount of the brain's electrical current flowing in and out of superficial scalp regions adjacent to that electrode. The CSD-reference might thus be preferred as the reference most likely to link surface recorded frontal EEG asymmetry to activity generated in frontal systems (as opposed to activity from distal non-frontal sources that can be reflected at frontal recording sites under other reference montages such as average or linked-mastoids). No EEG asymmetry studies of clinical depression using a CSD re-reference appear in the literature.

The present study addressed these methodological issues with a substantially larger sample than previous EEG asymmetry studies of depression, obtaining adequate power to examine whether relatively less left frontal activity would characterize individuals with any history of depression, and whether this effect might vary for men versus women. To reduce potential heterogeneity of depression, depressed participants included in the study met DSM-IV criteria for lifetime major depressive disorder (MDD) and endorsed no comorbid Axis I disorders with the exception of current dysthymia. Current depression was examined categorically (e.g., current MDD status) and dimensionally (e.g., by level of depressive questionnaire symptoms) to determine whether EEG asymmetry findings were simply due to elevated current symptomatology (indicating a state rather than a trait effect). To enhance the ability to identify trait-related variance associated with EEG asymmetry, resting EEG was recorded eight times, twice per day on four separate days within a two-week period to obtain a reliable measure of trait asymmetry for each participant. Moreover, asymmetry scores were calculated for four reference derivations (average, CSD, Cz, and linked mastoid) to examine whether depression-asymmetry relationships would be found for derivations that are more sensitive to localized potentials versus those that include potentials that are widely distributed or that arise from distant electrical sources.

Method

Participants

Prospective participants were identified using Beck Depression Inventory (BDI; Beck, Ward, Mendelson, Mock, & Erbaugh, 1961) scores obtained either from pre-testing in an introductory psychology course at the University of Arizona or an online survey. The sampling strategy was to recruit not only those with extreme high and low BDI scores, but to

recruit subjects with a broad range of symptoms that would include individuals ranging from no symptoms, to few symptoms, to several symptoms, to full clinical severity. All individuals with BDI pre-testing scores over 20 were invited for screening, with other participants drawn from among those individuals with scores in the ranges of 0-5, 6-10, 11-15, and 16-20, with the goal to obtain a sample with scores across the entire range of severity. Prospective participants were initially telephoned by a post-baccalaureate project manager, who determined if they met exclusionary criteria such as left handedness, history of head injury with loss of consciousness >10 minutes, concussion, epilepsy, electroshock therapy, use of current psychotropic medications, and active suicidal potential necessitating immediate treatment. Individuals receiving current psychotherapy were not excluded. Participants not meeting exclusionary criteria by this screen were invited for an intake interview. All participants accepted into the study were required to be strongly right handed (a score of greater than 35 on the 39 point scale of Chapman & Chapman, 1987). Figure 1 provides a detailed flow chart summarizing study recruitment across a four year period.

During the intake interview administered by a graduate-level trained clinical rater on a separate day before any EEG evaluations, participants were again screened for the same exclusionary criteria covered in the phone screen, and then further screened for Axis I psychopathology using the Structured Clinical Interview for DSM-IV (SCID, First, Spitzer, Gibbon & Williams, 1997). Participants were excluded if they met criteria for any current comorbid DSM-IV Axis I disorder other than lifetime MDD or current dysthymia. Inter-rater reliability analyses (performed by clinical interviewers and the first and last authors) for a randomly selected 10% of SCIDs demonstrated high inter-rater agreement for current and past MDD diagnoses (Kappa = .81 and .91, respectively). The final sample consisted of 306 participants (95 male), with an age range of 17 to 34 years ($M = 19.1$, $SE = 0.1$). A total of 143 participants met criteria for the lifetime MDD+ group (Table 1 lists DSM-IV diagnoses for this sample), whereas the remaining 163 participants did not meet criteria for any Axis I disorder and were considered the lifetime MDD- group.

Depression severity was assessed during the intake interview with the Beck Depression Inventory II (Beck, Steer, & Brown, 1996) and the 17-item Hamilton Rating Scale for Depression (HRSD; Hamilton, 1960). The intra-class correlation of inter-rater agreement was .95 for a randomly selected sample of 10% of HRSD interviews in the present sample. Because epidemiological studies have indicated an inverse relation between social class and rates of unipolar depression (e.g., Leventhal & Brooks-Gunn, 2000), and because SES has been linked to frontal EEG asymmetry (Tomarken, Dichter, Garber & Simien, 2004), socioeconomic status (SES) was assessed using the Four Factor Index of Social Status (Hollingshead, 1975).

Table 2 displays participant demographic information as a function of lifetime MDD status and biological sex. Univariate ANOVAs (lifetime MDD status by biological sex) examined these demographic variables. A main effect of lifetime MDD status emerged for BDI-II ($F(1, 302) = 112.0$, $p < .001$, $d = 1.30$) and HRSD ($F(1, 302) = 93.4$, $p < .001$, $d = 1.07$), indicating that the MDD+ group had higher scores on current depression measures than the MDD- group (see Table 2). No effects of sex or lifetime MDD status by sex emerged for BDI-II or HRSD (all $p > .12$). In addition, no effects emerged for SES (all $p > .22$).

EEG Data Collection and Reduction

Two resting EEG sessions were completed each day, on four separate days with no fewer than 24 hours between visits, and with all four visits completed within a 14 day period (such that the fourth day is not more than 14 days after the first day).¹ Nine participants attended fewer than all four EEG assessment days ($N = 5$ three days, $N = 2$ two days, $N = 2$ one day),

but all subjects were included in mixed linear model analyses that successfully accommodated missing data (see Bagiella, Sloan, and Heitjan, 2000).

Resting EEG at each session was recorded for eight one-minute baselines of eyes-open (O) and eyes-closed (C), in one of two counterbalanced orders (OCCOCOOC or COOCOCCO). Sessions within day were separated by approximately 20 minutes. All EEG data were collected using a 64-channel NeuroScan Synamps2 amplifier (Charlotte, NC) and acquisition system, utilizing the international 10-20 system for electrode placement. Two electrooculogram (EOG) channels (vertical: superior and inferior orbit of the left eye; lateral: outer canthi) were collected for ocular artifact rejection. All impedances were kept under 10K Ohms. Data for each resting session were digitized continuously at 1000 Hz, amplified 2816 times, and filtered with 200Hz low pass filter prior to digitization. EEG data were acquired with an online reference site immediately posterior to Cz and subsequently re-referenced offline to four different references: 1) average of all EEG leads = AVG, 2) CSD (using algorithms from Kayser & Tenke, 2006 and based on the spherical spline approach summarized by Perrin, Pernier, Bertrand, & Echallier, 1989, 1990), 3) Cz, and 4) averaged ("linked") mastoids = LM.

After acquisition, visual inspection removed epochs with movement and muscle artifacts. Data reduction was implemented using custom scripts in Matlab. A blink rejection algorithm rejected data segments where ocular activity exceeded ± 75 microvolts in the vertical ocular channel, and an artifact rejection algorithm rejected segments with large fast deviations in amplitude in any channel (e.g., DC shifts and spikes) that may have eluded human inspection. Data were segmented into one-minute EEG blocks and further epoched into 117 epochs of 2.048 sec per block, overlapping by 1.5 s. This overlapping compensates for the minimal weight applied to the end of the epoch by the use of the Hamming window function. Following windowing, a Fast Fourier Transform (FFT) was applied to all artifact-free epochs. The power spectra from all artifact free epochs across all eight minutes were averaged to provide a summary spectrum for each resting session.² Total alpha power (8-13 Hz) was then extracted from the spectrum for each resting session and site. An asymmetry score for each resting session for each reference montage was calculated by subtracting the natural log transformed scores (i.e., $\ln[\text{Right}] - \ln[\text{Left}]$) for each homologous left and right pair; e.g., F7 & F8, F5 & F6, F3 & F4, F1 & F2). Higher values on this index putatively reflect relatively greater left activity (i.e., greater right than left alpha; cf. Allen et al., 2004). Asymmetry scores for the two resting sessions within each day were then averaged together to create a measurement of regional brain activity per day. Thus, separate asymmetry scores for four days for each of four reference montages were utilized in analyses, resulting in 16 asymmetry scores per participant at each homologous pair. Although asymmetry scores were computed for all homologous site pairs, analyses for the present study focused on a specific subset of those pairs (F2-F1, F4-F3, F6-F5, F8-F7) that correspond to regions commonly studied throughout the asymmetry literature (pairs F4-F3 and F8-F7; see review by Coan & Allen, 2004) as well as pairs of channels that neighbor these pairs to provide better resolution of the lateral and medial extent of depression-related frontal asymmetry.³ Intraclass correlations indicated that these frontal EEG asymmetry scores were highly stable

¹Of the 21 participants who did not complete their sessions within a 14-day period, 15 completed all sessions within 16 days, whereas the remaining 6 completed all sessions within 18-20 days.

²Although several studies have demonstrated that eyes-open and eyes-closed alpha power do not differ from each other (e.g., Henriques & Davidson, 1990; Henriques & Davidson, 1991; Reid, Duke, & Allen, 1998), total alpha power was also calculated separately for usable eyes-open and eyes-closed epochs, and EEG asymmetry analyses performed for both types of data indicated that results generally replicated those of alpha power averaged across all eight minutes of each resting session.

³Since parietal EEG asymmetry has also been suggested as a risk marker for depression and therefore deserves its own in-depth examination, results for parietal channels are presented in an additional paper (Stewart, Towers, Coan, & Allen, 2010). To summarize the relationship between frontal and parietal channels, effects of frontal EEG asymmetry tended to reverse direction in the parietal region, particularly for AVG and LM.

across sessions for each reference montage (AVG range = .71-.77; CSD range = .67-.72; Cz range = .67-.72; LM range = .62-.66).

Results

Frontal EEG Asymmetry as a Function of Lifetime DSM-IV-Defined Depression

Lifetime MDD status—To examine the relationship between lifetime MDD status and frontal EEG asymmetry, a full factorial mixed linear model (SAS 9.2) was performed. Lifetime MDD status (past and/or current MDD = lifetime MDD+, never depressed = lifetime MDD-) and biological sex (male, female) were centered between-subjects variables, whereas visit day (4), reference (4: AVG, CSD, Cz, and LM), and channel (4: F2-F1, F4-F3, F6-F5, F8-F7) were within-subjects variables. EEG asymmetry score based on total 8-13 Hz alpha power was the dependent variable.⁴ Cohen's *d* is reported for significant differences between lifetime MDD groups.

Several effects not involving lifetime MDD status emerged. Results indicated that main effects of reference ($F(3, 906) = 122.3, p < .001$) and channel ($F(3, 906) = 39.2, p < .001$) were qualified by a reference by channel interaction ($F(9, 2718) = 5.8, p < .001$), and supplemental mixed models for each reference separately indicated that main effects of channel emerged for AVG, CSD, and LM (all $p < .001$) but not Cz ($p > .09$). For AVG and LM, F2-F1 was associated with the largest relative left frontal activity, followed by F4-F3, F6-F5, then F8-F7 (all channels were significantly different from each other at $p < .05$ with the exception of F2-F1 and F4-F3 for AVG). For CSD, F2-F1 and F4-F3 were associated with higher relative left frontal activity than F6-F5 and F8-F7 (all $p < .05$) but frontal asymmetry did not differ within medial channels (F2-F1 and F4-F3) or within lateral channels (F6-F5 and F8-F7).

Of key interest were effects involving lifetime MDD status. Figure 2 illustrates that a main effect of lifetime MDD emerged ($F(1, 302) = 4.7, p = .03$) that was qualified by a lifetime MDD by reference interaction ($F(3, 906) = 22.2, p < .001$), and supplemental full factorial mixed models performed separately for each reference montage indicated that a main effect of lifetime MDD emerged for CSD ($p < .001$ and $d = .93$), demonstrating that the lifetime MDD+ group displayed relatively less left frontal activity than the MDD- group. In contrast, no effects of lifetime MDD emerged for AVG ($p > .06$), Cz ($p > .26$), or LM ($p > .43$). No lifetime MDD by sex or lifetime MDD by sex by reference interactions emerged for the omnibus analysis (both $p > .29$). In addition, no lifetime MDD by channel or lifetime MDD by channel by reference interactions emerged (both $p > .92$), indicating that the lifetime MDD effect was not differentially manifest across the four frontal sites.

Follow-up analyses: Current MDD status and depression severity—Two approaches were used to assess whether the apparent link between lifetime MDD status and frontal EEG asymmetry for the CSD-referenced data was due to current levels of depressive symptoms. The first approach reran the full factorial linear mixed model with CSD referenced data but instead of lifetime MDD status, current MDD status was used (current MDD+ = all participants with current MDD, regardless of past MDD status; past MDD+ = participants with past MDD but not current MDD or current dysthymia; MDD- = participants without current or past MDD or dysthymia; six participants with past MDD but

⁴Asymmetry scores were also computed for lower alpha power (8.5-10.0 Hz) because differential effects associated with psychopathology have been observed in this band (e.g., Davidson, Marshall, Tomarken, & Henriques, 2000; Tomarken et al., 2004), and findings for lower alpha power largely replicated total alpha power results. In order to follow up on asymmetry metric results involving lifetime MDD and current depression severity, analyses examining hemispheric differences in total alpha power (as opposed to differences in asymmetry scores) are included in online supplementary analyses.

current dysthymia were not included in these analyses). Current MDD status was the between-subjects variable, day and channel were within-subject variables, and EEG asymmetry score was again the dependent variable. Cohen's d is reported for significant differences between current MDD+, past MDD+, and MDD- groups. Based on EEG asymmetry results, the main effect of current MDD status was the effect of importance. Figure 2 illustrates that a main effect of current MDD status emerged ($F(2, 297) = 14.1, p < .001$), indicating that the current MDD+ group and past MDD+ group displayed relatively less left frontal activity than the MDD- group (both $p < .001$ and $d = .57$ and $.58$, respectively), but did not differ significantly from each other. These findings suggest that the previously-reported effects of lifetime MDD were not accounted for solely by those with current MDD, as individuals with past MDD appear indistinguishable from those with current MDD in terms of frontal EEG asymmetry.

The second approach utilized continuous measures of depression severity. The between-group analysis above necessarily resulted in reduced sample sizes per cell, and additionally those diagnosed with past MDD still had some level of depressive symptoms, so continuous measures of symptomatology (as indexed by BDI-II and HRSD) were utilized to assess whether current severity was responsible for the lifetime MDD effects observed in the CSD referenced-data. Hierarchical linear mixed models using Type 1 (rather than Type 3) sums of squares were used to examine whether significant lifetime MDD results could be accounted for by depressive symptoms measured dimensionally in a model that included either BDI-II or HRSD scores as a main effect. Session day and channel were entered first, followed by BDI-II or HRSD (z-scored), then lifetime MDD was added to the model. EEG asymmetry score based on total 8-13 Hz alpha power was again the dependent variable. If EEG asymmetry results represent trait effects that are not due to current depressive symptomatology, a lifetime MDD effect should still emerge. A main effect emerged for BDI-II ($F(1, 303) = 11.0, p = .001$), indicating that higher BDI-II scores were associated with relatively less left frontal activity (estimated -1 SD on BDI-II: $M = .069$ and $SE = .01$ vs. estimated +1 SD on BDI-II: $M = .063$, $SE = .01$, $d = .05$). A main effect also emerged for HRSD ($F(1, 303) = 20.2, p < .001$), demonstrating that higher HRSD scores were associated with relatively less left frontal activity (estimated -1 SD on HRSD: $M = .077$ and $SE = .01$ vs. estimated +1 SD on HRSD: $M = .054$, $SE = .01$, $d = .21$). Most importantly, however, the main effect of lifetime MDD remained significant when entered after BDI-II ($F(1, 303) = 12.7, p < .001, d = .50$) or HRSD ($F(1, 303) = 18.6, p < .001, d = .43$), indicating that relationship of lifetime MDD status to CSD-referenced frontal asymmetry could not be accounted for by current levels of depression symptoms.

Supplementary Asymmetry Analyses for Other Reference Montages

In an attempt to reconcile nonsignificant AVG, Cz, and LM results of the present study with prior research that has been based on these reference montages and that has found significant relationships between frontal EEG asymmetry and BDI-defined depressive groups (e.g., Reid et al., 1998; Schaffer et al., 1983), supplementary analyses were undertaken by creating depression groups based on a measure of current depression symptom severity, the BDI-II. Asymmetry differences as a function of BDI-II defined depression status were examined for each reference separately. This conceptualization of depression may replicate several studies that have used questionnaire indices to define depression groups in conjunction with, or instead of, DSM-IV measures of depression (see Thibodeau et al., 2006 for several examples). Three groups were established after examination of the mean and standard deviation of BDI-II intake scores across the entire sample ($M = 11.1, SD = 9.6$): low (BDI-II scores from 0-10; males $n = 58$, females $n = 104$), moderate (BDI-II scores from 11-20; males $n = 23$, females $n = 65$), and high (BDI-II scores 21 or greater; males $n = 14$, females $n = 42$).

Full factorial mixed models were run for AVG, Cz, and LM separately with BDI-II group and biological sex as centered between-subjects variables, whereas day and channel were within-subjects variables. EEG asymmetry score was again the dependent variable. Although no main effect of BDI group emerged for AVG, Cz, or LM (all $p > .22$), a BDI group by sex interaction emerged for AVG ($F(2, 300) = 4.5, p = .01$), Cz ($F(2, 300) = 3.8, p = .02$), and LM ($F(2, 300) = 8.4, p < .001$), so follow-up mixed models were run for each sex separately to examine BDI-II group differences. Cohen's d is reported for significant differences between BDI-II depression groups.⁵

Results for women (see Figure 3, upper panel) indicated that main effects of BDI-II group emerged for AVG ($F(2, 208) = 7.5, p < .001$), wherein women with high-BDI-II scores displayed lower relative left frontal activity than women with moderate BDI-II scores ($p = .04$ and $d = .42$) and women with low BDI-II scores ($p < .001$ and $d = .70$). In addition, a main effect of BDI-II group emerged for LM ($F(2, 208) = 6.8, p < .01$), demonstrating that women with high and moderate BDI-II scores displayed lower relative left frontal activity than women with low levels of depression (both $p < .01$, and $d = .58$ and $.45$, respectively).

In contrast, findings for men (see Figure 3, lower panel) demonstrated that main effects of BDI-II group emerged for Cz ($F(2, 92) = 3.5, p = .03$), wherein men with moderate BDI-II scores exhibited lower relative left frontal activity than men with low BDI-II scores ($p = .01$ and $d = .63$) but those with high BDI-II scores did not differ from either group (both $p > .18$). A main effect of BDI-II group also emerged for LM ($F(2, 92) = 3.8, p = .03$), indicating that men with moderate and high BDI-II scores displayed higher relative left frontal activity than men with low scores ($p < .05$ and $d = .50$, and $p = .02$ and $d = .68$, respectively). No main effect of BDI-II group emerged for AVG-referenced data ($p > .36$).

Finally, BDI-II group analyses were also performed for the CSD reference to examine the concordance of results with previously reported CSD findings. Results indicated that a main effect of BDI-II group emerged ($F(2, 300) = 9.3, p < .001$), wherein participants with moderate and high BDI-II scores displayed lower relative left frontal activity than those with low scores (both $p < .001$ and $d = .50$ and $.48$, respectively). These findings are consistent with lifetime MDD and current MDD findings, which demonstrated no interactions between depression and sex for the CSD reference.

Discussion

The present study examined the relationship between asymmetries in frontal brain activity and depression, utilizing both categorical DSM diagnoses as well as measures of depressive severity. The key findings are that: 1) CSD-referenced frontal EEG asymmetry differentiates those individuals with lifetime MDD from those without, an effect not related to current depression, and consistent with the notion that frontal asymmetry can function as an endophenotype of risk for depression; 2) EEG asymmetry derived from other reference montages — those that have been used with some frequency in the literature to produce a pattern of results that is plagued by some inconsistencies — produced a pattern of findings that suggests some degree of relationship between frontal asymmetry and depressive severity in women, but not consistently so for men.

Lifetime MDD and EEG Asymmetry

Results examining lifetime MDD status are consistent with the proposition that current source density-referenced frontal EEG asymmetry is an endophenotype related to risk for depression in both women and men, unconfounded by current symptom severity. Participants with lifetime MDD (both current MDD and past MDD) displayed less relative left frontal activity than never-depressed participants, and measures of current depression

(BDI-II and HRSD scales) did not account for this effect. These findings indicate that current source density-referenced EEG asymmetry may be a liability marker, identifying a vulnerability to develop depression (e.g., Allen et al., 2004b, Gotlib et al., 1998; Henriques & Davidson, 1990, 1991) and related dysphoric conditions (Accortt & Allen, 2006; Coan & Allen, 2008), although prospective studies are needed to definitively test this proposition.

Current Depression Severity and Sex Differences in EEG Asymmetry

Unlike the current source density reference, frontal asymmetry was not related to lifetime MDD using the average, Cz, or linked mastoid references, which are the reference montages commonly used in the EEG asymmetry literature. Instead, the nature of the relationship between depression and EEG asymmetry was modified by current depression severity as measured by the BDI-II and sex. Analyses examining BDI-II depression groups indicated that for average and linked mastoid references, women with high levels of depressive symptoms exhibited relatively less left frontal brain activity at rest than women with low levels of depressive symptoms, with women with moderate levels of depression displaying means in between low and high groups. These findings for women replicate several studies with predominately or exclusively female samples that find current depression is linked to relatively less left than right frontal activity (see Coan and Allen, 2003, and Thibodeau et al., 2006).

In contrast to women, who demonstrated a relatively consistent pattern of EEG asymmetry results as a function of current depression severity, men exhibited a weak relationship and also inconsistent patterns of findings across these three reference montages. For example, men with moderate depressive symptoms, but not those with high levels of depressive symptoms, displayed relatively less left frontal activity than men with low levels of depression for the Cz reference; conversely, linked mastoid results demonstrated that men with high levels of depression exhibited relatively greater left frontal brain activity at rest than men with low levels of depression. Results for the average reference were non-significant. The linked-mastoid findings are consistent with one study that directly compared gender differences using an average reference (Miller et al., 2002), yet inconsistent with another study that examined a male sample using a linked-ears reference (Jacobs & Snyder, 1996). Although the latter study found that men with relative left frontal EEG asymmetry displayed lower BDI scores, significant results were confined to only one site (whereas results for the present study extend across a wide region of frontal cortex), with EEG asymmetry measured on only one occasion, and the range of BDI scores in that study appeared very limited, with few men endorsing scores in the high depressive range of the present study (BDI-II scores of 21 or higher).

In summary, women and men displayed opposing patterns of frontal EEG asymmetry as a function of current depression severity for the linked mastoid reference (and the pattern of means for the average reference was consistent with the linked-mastoid pattern). Whether these findings signify different mechanisms underlying depression in men and women remains an open question. Although it is possible that these asymmetry findings may reveal different causal pathways, they might also reflect different symptom constellations in depression in men and women (e.g., Kornstein et al., 2000; Silverstein, 2002). For example, pure MDD+ men suffer more sudden spells of anger and aggression than pure MDD+ women (Winkler, Pjrek, & Kasper, 2005), suggesting anger could be a moderator of EEG asymmetry that could explain sex differences in depression, since relatively greater left frontal activity characterizes individuals with trait anger at rest (e.g., Harmon-Jones & Allen, 1998). Additional evidence in support of this argument of anger as a moderator of frontal brain asymmetry are findings demonstrating that boys without oppositional defiant disorder display relatively less left frontal EEG activity, consistent with the EEG asymmetry pattern exhibited by MDD- men in the present study, whereas oppositional-defiant boys

(who had higher rates of depression than healthy boys) demonstrated no asymmetry (Baving, Laucht, & Schmidt, 2000). Therefore, limited research supports the assertion that anger and related aggressive behaviors may influence patterns of frontal brain activity differently in depressed and non-depressed men.

Choice of Reference and EEG Asymmetry

The present study indicates that a pattern of relatively less left frontal EEG activity as measured by the current source density reference reflects a trait-like marker of depression risk for men and women, whereas a similar pattern of EEG asymmetry measured by average and linked mastoid references appears to reflect a severity-related marker of current depression, at least in women. Since the current source density algorithm attenuates broad electrical sources, such as distal parietal and occipital sources where EEG alpha activity is typically quite large (Hagemann et al., 2001; Kayser & Tenke, 2006), it is more probable that the trait-like index of current source density-referenced EEG asymmetry reflects predominantly frontal sources. By contrast, the severity-related indices of EEG asymmetry derived from AVG, LM, and Cz reference montages will potentially reflect considerable nonfrontal activity, since these references reflect both proximal and distal sources, in part due to lower signal-to-noise ratios in the measurement of frontal EEG activity than those derived from the CSD reference (Hagemann et al., 2001). In order for frontal EEG asymmetry to be an endophenotype for depression risk, it must converge with other possible risk indicators for depression (e.g., genetic variations in serotonin genes involved in the transmission and maintenance of depression), and CSD-referenced EEG asymmetry appears most promising in this endeavor.

The Cz reference produced atypical results for men and null findings for women, yet this is not inconsistent with the fact that a majority of significant effects for the Cz reference in depression-related studies (e.g., 19 in the Thibodeau et al., 2006 meta-analysis) used infants or children as participants, not young adults. Only four studies with adults from the Thibodeau et al. (2006) meta-analysis demonstrated Cz-related differences in depressed groups (and two of the four consisted of small samples of middle-aged adult participants: Baehr, Rosenfeld, Baehr, & Earnest, 1998; Henriques & Davidson, 1991), whereas here we replicate larger studies that reported null results with the Cz reference in young adults (e.g., Bruder et al., 1997; Reid et al., 1998). EEG asymmetry researchers have shown that there is little convergence between Cz and other references (e.g., Hagemann et al., 2001; Reid et al., 1998) and have advocated the use of references other than Cz since variations in power at this active EEG site can distort the direction and strength of asymmetry recorded from lateral sites in either hemisphere (e.g., Davidson, 1998; Hagemann et al., 2001).

The findings using average-, linked mastoid-, and Cz-referenced asymmetry scores are only partially consistent with the extant literature. They are consistent in that: 1) To the extent that asymmetry relates to depression under these reference montages, it does so in women, and most of the extant depression and EEG asymmetry literature is based on female samples (Coan & Allen, 2004; Thibodeaux et al., 2006); and, 2) relatively less left frontal activity was observed among those with higher levels of depressive symptoms. The findings are inconsistent in that they fail to support the idea that frontal EEG asymmetry recorded under these reference montages is related to a lifetime history of MDD. For example, Gotlib et al. (1998) found that individuals with any history of depression, regardless of whether currently symptomatic, showed relatively less left frontal activity as assessed using a Cz reference montage, and Henriques and Davidson (1990) found that previously depressed but euthymic individuals showed relatively less left frontal activity than never depressed controls using the LM reference montage. One key difference between these studies and the present study might be that the present study obtained a sample that spanned a wide range of depressive severity, such that the participants with no history of major depression in the present study

did not have an absence of depressive symptoms. The lifetime MDD- participants in the present study had a mean BDI-II score of 6.2, whereas the never-depressed controls in the Henriques and Davidson (1990) study had comparatively little symptomatology (mean BDI of 1.4). Because lifetime MDD- participants in the current study had at least some level of symptomatology, the present study may have been at a disadvantage to find differences between lifetime MDD+ and lifetime MDD- if frontal EEG asymmetry under these montages tracks symptom severity to some extent.

Methodological Issues and Limitations

The design of the present study was advantageous for examining the relationship between lifetime MDD and EEG asymmetry in women and men due to 1) the recruitment of a large sample of medically healthy, interview-diagnosed, medication-free individuals with no comorbid anxiety disorders; 2) repeated sessions of EEG recording that provided a reliable estimate of trait asymmetry; and, 3) a systematic examination of reference montages, including the first to examine the current-source density montage in depression. The consistency of the findings extended across both medial and lateral regions of frontal cortex, but different references indexed different aspects of depression, with the CSD-referenced data serving as a viable endophenotype, whereas average- and linked mastoid-referenced data appear to show a female-specific relationship to depressive severity. The relationship to depressive severity was found across persons, not within persons over time, and there is no support from the literature that, within persons, changes in EEG asymmetry track changes in levels of symptoms over intervals of weeks or months (Allen, Urry, Hitt, & Coan, 2004; Debener et al., 2000). Future work can examine whether changes in severity over longer intervals tracks frontal asymmetry using average- and linked-mastoid references.

Limitations of the present study include a younger cohort, suggesting that these findings may not be assumed to apply for later-life onset depression. Our early-onset sample, however, might be expected to have a recurrent or chronic course of depression (cf. findings with chronic depression, Klein et al., 1999) and thus may generalize to individuals with severe depression later in life. In addition, these results may not be generalizable to depressed individuals with comorbid DSM-IV conditions such as anxiety and substance use disorders, which are common in MDD patients (e.g., Kessler et al., 2003). Furthermore, it is unclear whether these findings may extend to persons with MDD who are being treated with psychotropic medications, but to date no evidence suggests that antidepressants impact frontal EEG asymmetry. Finally, the present study offers a cross-sectional investigation to address a within-subjects question, namely whether an individual's pattern of regional frontal brain activity predicts subsequent development of MDD, so prospective, longitudinal data are needed to firmly establish that EEG asymmetry is a vulnerability marker for depression.

Synopsis

Findings from CSD-referenced data indicated that a pattern of less relative left frontal activity (inferred from relatively greater left frontal alpha power) was evident for individuals with lifetime MDD compared to never-depressed individuals, regardless of sex or current severity of depressive symptoms. These findings clearly suggest the promise of CSD-referenced frontal EEG asymmetry as an endophenotype of depression risk. EEG asymmetry for average and linked mastoid references, however, showed evidence of sex-specific contributions related to current depressive symptom severity, reducing the likelihood that EEG asymmetry under these references montages can serve profitably as an endophenotypic marker of depression. For these references a pattern of less relative left frontal activity characterized women with currently high levels of depressive symptoms as compared to women with low levels of current depression. For men, by contrast, in the largest study of

EEG asymmetry of depression to date, findings from the linked mastoid reference revealed that men with moderate or high levels of depressive symptoms showed relatively greater left frontal activity, a finding opposite to women, but nonetheless replicating one previous study with a sizable male sample (Miller et al., 2002). The implications of this finding for men must await further investigation, but it is clear that future studies of frontal brain asymmetry and risk for depression must take sex differences into consideration.

Because resting frontal EEG asymmetry has demonstrated inconsistencies in its relationship to depression and emotion in previous studies, because its underlying mechanisms are poorly if at all understood, and because it is characterized by many methodological quirks (including its predominant manifestation as difference score and reference-dependent effects), resting frontal EEG asymmetry has occupied a decidedly (and deservedly) tentative status as a potential endophenotypic marker of depression (Coan & Allen, 2008). Although the present study has not entirely resolved the many methodological and theoretical mysteries long associated with frontal EEG asymmetry as an index of psychopathology, it has brought the measure significantly up to date in a larger and more representative sample of individuals than has previously been reported. If associations between frontal EEG asymmetry and depression were illusory or otherwise absent, the current study was sufficiently powered to advocate just such a conclusion. The strength of the results reported here suggests, by contrast, that associations between frontal EEG asymmetry and affective psychopathology are real, important, and worthy of considerable scientific inquiry as the pursuit of endophenotypic markers of depression vulnerability continues.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported in part by grants from the National Institutes of Health (R01-MH066902) and the National Alliance for Research on Schizophrenia and Depression (NARSAD) to John Allen. The authors wish to thank Eliza Ferguson, Jamie Velo, Dara Halpern, Craig Santerre, Eynav Accortt, Amanda Brody, Jay Hegde and myriad research assistants for their help on this project

References

- Accortt E, Allen JJB. Frontal EEG asymmetry and premenstrual dysphoria. *Journal of Abnormal Psychology* 2006;115:179–184. [PubMed: 16492109]
- Allen JJB, Coan JA, Nazarian M. Issues and assumptions on the road from raw signals to metrics of frontal EEG asymmetry in emotion. *Biological Psychology* 2004a;67:183–218. [PubMed: 15130531]
- Allen JJB, Urry HL, Hitt SK, Coan JA. The stability of resting frontal electroencephalographic asymmetry in depression. *Psychophysiology* 2004b;41:269–280. [PubMed: 15032992]
- Anokhin AP, Heath AC, Myers E. Genetic and environmental influences on frontal EEG asymmetry: A twin study. *Biological Psychology* 2006;71:289–295. [PubMed: 16054745]
- Baehr E, Rosenfeld JP, Baehr R, Earnest C. Comparison of two EEG asymmetry indices in depressed patients vs. normal controls. *International Journal of Psychophysiology* 1998;31:89–92. [PubMed: 9934624]
- Bagiella E, Sloan RP, Heitjan DF. Mixed effects models in psychophysiology. *Psychophysiology* 2000;37:13–20. [PubMed: 10705763]
- Baving L, Laucht M, Schmidt MH. Oppositional children differ from healthy children in frontal brain activation. *Journal of Abnormal Child Psychology* 2000;28:267–275. [PubMed: 10885684]
- Beck, AT.; Steer, RA.; Brown, GK. *The Beck Depression Inventory–II*. Harcourt Assessment; San Antonio: 1996.

- Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Archives of General Psychiatry* 1961;4:561–571. [PubMed: 13688369]
- Bruder GE, Fong R, Tenke CE, Leite P, Towey JP, Stewart JE, et al. Regional brain asymmetries in major depression with or without an anxiety disorder: A quantitative electroencephalographic study. *Biological Psychiatry* 1997;41:939–948. [PubMed: 9110099]
- Chapman LJ, Chapman JP. The measurement of handedness. *Brain and Cognition* 1987;6:175–183. [PubMed: 3593557]
- Coan, JA.; Allen, JJB. The state and trait nature of frontal EEG asymmetry in emotion.. In: Hugdahl, K.; Davidson, R.J., editors. *The Asymmetrical Brain*. second ed. MIT Press; Cambridge, MA: 2003. p. 565-615.
- Coan JA, Allen JJB. Frontal EEG asymmetry as a moderator and mediator of emotion. *Biological Psychology* 2004;67:7–49. [PubMed: 15130524]
- Coan, JA.; Allen, JJB. Affective style and risk for psychopathology.. In: Beauchaine, TP.; Hinshaw, SP., editors. *Child Psychopathology*. Wiley Press; Hoboken: 2008.
- Coan JA, Allen JJB, McKnight PE. A capability model of individual differences in frontal EEG asymmetry. *Biological Psychology* 2006;72:198–207. [PubMed: 16316717]
- Davidson RJ. Anterior electrophysiological asymmetries, emotion, and depression: Conceptual and methodological conundrums. *Psychophysiology* 1998;35:607–614. [PubMed: 9715104]
- Davidson RJ, Marshall JR, Tomarken AJ, Henriques JB. While a phobic waits: Regional brain electrical and autonomic activity in social phobics during anticipation of public speaking. *Biological Psychiatry* 2000;47:85–95. [PubMed: 10664824]
- Dawson G, Frey K, Panagiotides H, Osterling J, Hessel D. Infants of depressed mothers exhibit atypical frontal brain activity: A replication and extension of previous findings. *Journal of Child Psychology and Psychiatry* 1997;38:179–186. [PubMed: 9232464]
- Debener S, Beauducel A, Nessler D, Brocke B, Heilemann H, Kayser J. Is resting anterior EEG alpha asymmetry a trait marker for depression? *Neuropsychobiology* 2000;41:31–37. [PubMed: 10629433]
- First, MG.; Spitzer, RL.; Gibbon, M.; Williams, JB. *Structured clinical interview for DSM-IV Axis I disorder—clinical version, administration booklet*. Biometrics Research Department; New York, NY: 1997.
- Forbes EE, Shaw DS, Fox NA, Cohn JF, Silk JS, Kovacs M. Maternal depression, child frontal asymmetry, and child affective behavior as factors in child behavior problems. *Journal of Child Psychology and Psychiatry* 2006;47:79–87. [PubMed: 16405644]
- Gotlib IH, Ranganath C, Rosenfeld JP. Frontal EEG alpha asymmetry, depression, and cognitive functioning. *Cognition & Emotion* 1998;12:449–478.
- Gottesman II, Gould TD. The endophenotypic concept in psychiatry: Etymology and strategic intentions. *American Journal of Psychiatry* 2003;160:636–645. [PubMed: 12668349]
- Greenberg PE, Kessler RC, Birnbaum HG, Leong SA, Lowe SW, Berglund PA, et al. The economic burden of depression in the United States: How did it change between 1990 and 2000? *Journal of Clinical Psychiatry* 2003;64:1465–1475. [PubMed: 14728109]
- Hagemann D. Individual differences in anterior EEG asymmetry: Methodological problems and solutions. *Biological Psychology* 2004;67:157–182. [PubMed: 15130530]
- Hagemann D, Naumann E, Thayer JF. The quest for the EEG reference revisited: A glance from brain asymmetry research. *Psychophysiology* 2001;38:847–857. [PubMed: 11577908]
- Hagemann D, Naumann E, Thayer JF, Bartussek D. Does resting electroencephalograph asymmetry reflect a trait?: An application of latent state-trait theory. *Journal of Personality and Social Psychology* 2002;82:619–641. [PubMed: 11999928]
- Hamilton M. A rating scale for depression. *Journal of Neurology, Neurosurgery, and Psychiatry* 1960;23:56–62.
- Harmon-Jones E, Allen JJB. Anger and frontal brain activity: EEG asymmetry consistent with approach motivation despite negative affective valence. *Journal of Personality and Social Psychology* 1998;74:1310–1316. [PubMed: 9599445]
- Heller W, Nitschke JB. The puzzle of regional brain activity in depression and anxiety: The importance of subtypes and comorbidity. *Cognition & Emotion* 1998;12:421–447.

- Henriques JB, Davidson RJ. Regional brain electrical asymmetries discriminate between previously depressed and healthy control subjects. *Journal of Abnormal Psychology* 1990;99:22–31. [PubMed: 2307762]
- Henriques JB, Davidson RJ. Left frontal hypoactivation in depression. *Journal of Abnormal Psychology* 1991;100:535–545. [PubMed: 1757667]
- Hollingshead, AB. Four factor index of social status. Department of Sociology, Yale University; New Haven, CT: 1975.
- Iacono WG. Identifying psychophysiological risk for psychopathology: examples from substance abuse and schizophrenia research. *Psychophysiology* 1998;35:621–637. [PubMed: 9844425]
- Jacobs GD, Snyder D. Frontal brain asymmetry predicts affective style in men. *Behavioral Neuroscience* 1996;110:3–6. [PubMed: 8652069]
- Judd LL, Akiskal HS, Zeller PJ, Paulus M, Leon AC, Maser JD. Psychosocial disability during the long-term course of unipolar depressive disorder. *Archives of General Psychiatry* 2000;57:375–380. [PubMed: 10768699]
- Kayser J, Tenke CE. Principal components analysis of Laplacian waveforms as a generic method for identifying ERP generator patterns: I. Evaluation with auditory oddball tasks. *Clinical Neurophysiology* 2006;117:348–368. [PubMed: 16356767]
- Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, et al. The epidemiology of major depressive disorder: Results from the National Comorbidity Survey Replication (NCS-R). *Journal of the American Medical Association* 2003;289:3095–3105. [PubMed: 12813115]
- Kessler RC, DuPont RL, Berglund P, Wittchen H. Impairment in pure and comorbid generalized anxiety disorder and major depression at 12 months in two national surveys. *American Journal of Psychiatry* 1999;156:1915–1923. [PubMed: 10588405]
- Klein DN, Schatzberg AF, McCullough JP, Dowling F, Goodman D, Howland RH, et al. Age of onset in chronic major depression: Relation to demographic and clinical variables, family history, and treatment response. *Journal of Affective Disorders* 1999;55:149–157. [PubMed: 10628884]
- Kornstein SG, Schatzberg AF, Thase ME, Yonkers KA, McCullough JP, Keitner GI, et al. Gender differences in chronic major and double depression. *Journal of Affective disorders* 2000;60:1–11. [PubMed: 10940442]
- Leventhal T, Brooks-Gunn J. The neighborhoods they live in: The effects of neighborhood residence on child and adolescent outcomes. *Psychological Bulletin* 2000;126:309–337. [PubMed: 10748645]
- Miller A, Fox NA, Cohn JF, Forbes EE, Sherrill JT, Kovacs M. Regional patterns of brain activity in adults with a history of childhood-onset depression: Gender differences and clinical variability. *American Journal of Psychiatry* 2002;159:934–940. [PubMed: 12042180]
- Perrin F, Pernier J, Bertrand O, Echallier JF. Spherical splines for scalp potential and current density mapping. *Electroencephalography and clinical Neurophysiology* 1989;72:184–187. [PubMed: 2464490]
- Perrin F, Pernier J, Bertrand O, Echallier JF. Corrigenda. *Electroencephalography and clinical Neurophysiology* 1990;76:565–566.
- Pizzagalli DA, Jahn AL, O'Shea JP. Toward an objective characterization of an anhedonic phenotype: A signal detection approach. *Biological Psychiatry* 2005;57:319–327. [PubMed: 15705346]
- Pizzagalli DA, Nitschke JB, Oakes TR, Hendrick AM, Horras KA, Larson CL, et al. Brain electrical tomography in depression: The importance of symptom severity, anxiety, and melancholic features. *Biological Psychiatry* 2002;52:73–85. [PubMed: 12113998]
- Reid SA, Duke LM, Allen JJB. Resting frontal electroencephalographic asymmetry in depression: Inconsistencies suggest the need to identify mediating factors. *Psychophysiology* 1998;35:389–404. [PubMed: 9643053]
- Silverstein B. Gender differences in the prevalence of somatic versus pure depression: A replication. *American Journal of Psychiatry* 2002;159:1051–1052. [PubMed: 12042198]
- Smit DJA, Posthuma D, Boomsma DI, de Geus EJC. The relation between frontal EEG asymmetry and the risk for anxiety and depression. *Biological Psychology* 2007;74:26–33. [PubMed: 16875773]

- Stewart JL, Towers DN, Coan JA, Allen JJB. The oft-neglected role of parietal EEG asymmetry and risk for major depressive disorder. submitted.
- Tenke CE, Kayser J. Reference-free quantification of EEG spectra: Combining current source density (CSD) and frequency principal components analysis (fPCA). *Clinical Neurophysiology* 2005;116:2826–2846. [PubMed: 16257577]
- Thibodeau R, Jorgensen RS, Kim S. Depression, anxiety, and resting frontal EEG asymmetry: A meta-analytic review. *Journal of Abnormal Psychology* 2006;115:715–729. [PubMed: 17100529]
- Tomarken AJ, Dichter GS, Garber J, Simien C. Resting frontal brain activity: Linkages to maternal depression and socio-economic status among adolescents. *Biological Psychology* 2004;67:77–102. [PubMed: 15130526]
- Tomarken AJ, Davidson RJ, Wheeler RE, Kinney L. Psychometric properties of resting anterior EEG asymmetry: Temporal stability and internal consistency. *Psychophysiology* 1992;29:576–592. [PubMed: 1410187]
- Towers DN, Allen JJB. A better estimate of the internal consistency reliability of frontal EEG asymmetry scores. *Psychophysiology* 2009;46:132–142. [PubMed: 19055503]
- Winkler D, Pjrek E, Kasper S. Anger attacks in depression – Evidence for a male depressive syndrome. *Psychotherapy and Psychosomatics* 2005;74:303–307. [PubMed: 16088268]

Note: BDI = Beck Depression Inventory. LOC = Loss of consciousness. MDD = major depressive disorder. PTSD = posttraumatic stress disorder. NOS = Not otherwise specified. OCD = obsessive compulsive disorder. GAD = generalized anxiety disorder. ADHD = attention deficit hyperactivity disorder.

Figure 1.
Flowchart of participant screening and enrollment.

Figure 2.

Panel A shows frontal alpha asymmetry scores (8-13 Hz at F2-F1, F4-F3, F6-F5, F8-F7) by lifetime MDD status for each reference montage (AVG = average; CSD = current source density; CZ = Cz; LM = linked mastoid), across all four frontal regions depicted on the head insert. Error bars reflect standard error. Panel B shows results of a follow-up assessment indicating the relationship of lifetime MDD status to CSD-referenced asymmetry is not solely accounted for by current MDD status. Y-axis is $\ln \mu V^2$ for AVG, Cz, and LM references, and $\ln \mu V^2 / \text{cm}^2$ for CSD referenced data.

Table 1DSM-IV Diagnoses Endorsed in Lifetime MDD+ Group ($N = 143$)

Diagnosis	Biological Sex	Frequency
Current MDD only	Men	5
	Women	9
Past MDD only	Men	20
	Women	55
Current MDD and Past MDD	Men	10
	Women	29
Current MDD and Current Dysthymia	Men	0
	Women	2
Past MDD and Current Dysthymia	Men	1
	Women	5
Current MDD, Past MDD, and Current Dysthymia	Men	3
	Women	4

Table 2

Group Demographics

MDD Status	Biological Sex	Caucasian %	BDI-II M (SE)	HRSD M (SE)	SES M (SE)
Lifetime MDD+	Men (N = 39)**	71.8	15.2 (1.3)	9.6 (0.9)	44.3 (2.2)
	Women (N = 104)*	77.9	17.6 (0.8)	11.7 (0.5)	41.9 (1.3)
Lifetime MDD-	Men (N = 56)*	66.1	5.7 (1.1)	3.9 (0.7)	42.6 (1.8)
	Women (N = 107)	71.0	6.2 (0.8)	4.0 (0.5)	44.3 (1.3)

Note: MDD= Major Depressive Disorder. BDI-II = Beck Depression Inventory II. HRSD = Hamilton Rating Scale for Depression. SES = socioeconomic status as determined by Hollingshead (1975).

* 1 participant did not report SES.

** 2 participants did not report SES.