Published in final edited form as:

Inorg Chem. 2009 September 7; 48(17): 8342–8356. doi:10.1021/ic900975y.

Molecular Oxygen and Sulfur Reactivity of a Cyclotrimeratrylene (CTV) Derived Trinuclear Copper(I) Complex

Debabrata Maiti[†], Julia S. Woertink[‡], Reza A. Ghiladi[†], Edward I. Solomon[‡], and Kenneth D. Karlin^{†,*}

[†]Department of Chemistry, The Johns Hopkins University, Baltimore, Maryland 21218

[‡]Department of Chemistry, Stanford University, Stanford, California 94305

Abstract

Our continuing efforts into developing copper coordination chemistry relevant to dioxygen-processing copper proteins has led us to design and synthesize a cyclotrimeratrylene (CTV)-based trinucleating ligand, CTV-TMPA, which employs tetradentate tris(2-pyridylmethyl)-amine chelates (TMPA) for their copper ion binding sites. Binding of three copper ions per CTV-TMPA unit was established by various chemical and spectroscopic methods such as UV-vis and resonance Raman (rR) spectroscopies. The following complexes were observed: A tricopper(I) complex [(CTV-TMPA)Cu₃]³⁺ (**1**), a CO adduct [(CTV-TMPA)Cu₃(CO)₃]³⁺ (**1-CO**; $\nu(\text{C}=\text{O}) = 2094 \text{ cm}^{-1}$), a triphenylphosphine adduct [(CTV-TMPA)Cu₃(PPh₃)₃]³⁺ (**1-PPh₃**), a tricopper(II) complex [(CTV-TMPA)Cu₃]³⁺ (**1-Ox**) and its tris-monochloride or tris-monobromide adducts. Also, introduction of dioxygen to the -80°C solutions of **1** leads to O₂-adducts, the first example of a synthetic copper complex which can stabilize a mononuclear Cu^{II}-superoxo and dinuclear peroxo species simultaneously within one complex {[Cu] = 1.53 mM in THF: (μ -1,2-peroxo complex, $\lambda_{\text{max}} = 543$ (ϵ 9650) nm): $\nu(\text{O}-\text{O}) = 825$ ($(\Delta^{18}\text{O}_2) = -47$) cm^{-1} ; $\nu(\text{Cu}-\text{O}) = 506$ ($(\Delta^{18}\text{O}_2) = -26$) cm^{-1} ; (superoxo complex, $\lambda_{\text{max}} = 427$ (ϵ 3150) nm): $\nu(\text{O}-\text{O}) = 1129$ ($(\Delta^{18}\text{O}_2) = -60$) cm^{-1} ; $\nu(\text{Cu}-\text{O}) = 463$ ($(\Delta^{18}\text{O}_2) = -27$) cm^{-1} }. Elemental sulfur reacts reversibly with **1** leading to a (proposed) hexanuclear species {[(CTV-TMPA)Cu₃]₂(μ -1,2-S₂²⁻)₃]⁶⁺ (**1-S**) { $\lambda_{\text{max}} = 544$ (ϵ 7270 nm)}, possessing one dicopper(II)-disulfide structural type: {THF solvent} $\nu(\text{S}-\text{S}) = 489$ ($(\Delta^{34}\text{S}) = -10$) cm^{-1} ; $\nu(\text{Cu}-\text{S}) = 506$ ($(\Delta^{34}\text{S}) = -5$) cm^{-1} }. Derivation of spectroscopic, structural and chemical conclusions were aided by the study of a close mononuclear analogue with one pyridyl group of the TMPA parent possessing a 6-CH₂OCH₃ substituent, this being part of the CTV-TMPA architecture.

Introduction

Biological systems containing multinuclear copper centers have motivated synthetic inorganic chemists to attempt to mimic bio-site structure, spectroscopy and active-site function.^{1–7} As this topic pertains to copper ion, binuclear copper centers occur in the O₂-carrier hemocyanins and the related tyrosinase monooxygenases and synthetic model systems for such proteins are now abundant.^{5,6,8,9} Copper cluster are well established for the “multicopper oxidases” (MCOs).^{10–15} These extensively studied proteins reveal synergism between two or more copper sites in the activation of dioxygen.^{10,11,16} The currently well-characterized members include ascorbate oxidase (AO) (Figure 1, top), laccase (Lc), ceruloplasmin (Cp), Fet3, bilirubin oxidase (BO) and CueO. In these proteins, the electron-transfer cycle is conveniently mediated by four copper sites which store

Correspondence: karlin@jhu.edu.

oxidizing power upon complete four-electron reduction of dioxygen to water prior to delivering oxidizing equivalents in a sequential manner to a substrate such as ascorbate or iron(II) ion. X-ray crystallographic determinations of the most prominent member of this series, AO, along with others, confirm that three of these four copper ions are intimately engaged to form a catalytic triad, one type 2 (mononuclear) and one type 3 (couple binuclear) copper center. The fourth copper ion is a distinct type-1 (or “blue”) copper center that oxidizes substrates which bind nearby and most likely contributes to the overall stoichiometry by long range (~12.5 Å in AO) but through-bond electron-transfer to the trinuclear cluster.

Multi-copper cluster also exist in the active site of nitrous oxide reductase (N₂OR) (Figure 1, bottom) which possesses four copper ions in close proximity to form a (histidine)₇Cu₄(μ₄-S) cluster (referred to as Cu₇).^{1,17,18} This mediates the final step in bacterial denitrification, N₂O + 2e⁻ + 2H⁺ → H₂O + N₂ and plays a critical environmental role by preventing release of greenhouse gas nitrous oxide. Based upon biochemical and spectroscopic studies, particulate methane monooxygenase (pMMO) (found in methanotrophic eubacteria) and ammonia monooxygenase (AMO) (ammonia-oxidizing eubacteria) are believed by some to contain copper cluster.^{19,20} These oxygenases catalyze a number of industrially important transformations, including the conversion of methane to methanol, ethane hydroxylation and the oxidation of other hydrocarbons, halogenated organics, and carbon monoxide.

A variety of trinuclear copper model compounds have been obtained by self-assembly reactions.^{21–34} In an alternative approach, conformationally constrained trinucleating ligands have been introduced to bind three metal ions in close proximity. Most of these are rather flexible and fail to impose a perfectly well defined arrangement of the metals. Some of the synthetic ligand systems employed from the groups of Karlin,^{35–38} Kim,³⁹ Itoh,^{40,41} Tolman,⁴² Casella,⁴³ Fenton^{3,44} and Chan⁴⁵ are depicted in Chart 1. Our earlier efforts^{35–38} into modeling the active sites of copper-cluster containing enzymes led us to synthesize trinucleating ligands including mesitylene-based organics (Chart 1, MesPy1, MesPy2)⁴⁶ which employ tridentate copper-chelates as their copper binding sites. Structural characterization of the resulting tricopper complexes revealed that the trimethyl-mesitylene motif may not provide the cluster-like geometry required for Cu₃/O₂ intramolecular reactivity, since the complexes have two copper centers located on one side of the mesityl plane, with the third on the other ‘opposite’ side. In a similar context, Kim and Itoh employed ‘1,3,5-triethyl-benzene ‘steering’ motifs showing that this forces all three copper centers to be on the same side of the mesityl plane.^{39–41}

In this report, we present our continuing efforts towards modeling the active sites of the multicopper enzymes by use of a ligand scaffold based upon the bowl-shaped cyclotrimeratriylene (CTV) molecule⁴⁷ possessing appended pyridyl alkylamine moieties.⁴⁶ The hope was that the more structured CTV architecture^{47–51} will help promote close di- or tri-copper interactions as occur in the multicopper enzyme active sites. Investigations of dioxygen reactivity with mono- and binuclear copper(I) complexes employing similar tetradentate pyridyl-alkylamine copper chelates have been previously carried out,^{5,6,9,52–54} thereby establishing patterns for copper-dioxygen adduct formation and representing a good starting point for obtaining insights with the CTV ligand and achieving our current objectives.

Experimental Section

Materials and Methods

Unless otherwise stated all solvents and chemicals used were of commercially available analytical grade. Methanol, dichloromethane, acetonitrile (CH₃CN), tetrahydrofuran (THF),

pentane, 2-methyltetrahydrofuran (MeTHF), acetone and diethyl ether were used after passing them through a 60 cm long column of activated alumina (Innovative Technologies, Inc.) under argon. Preparative thin-layer chromatography was performed on a Harrison Research Chromatotron Model 8924 equipped with a 4mm Adsorbosil-Plus P (silica gel, Alltech Associates) plate. Thin-layer chromatography was performed on 'Baker-Flex' aluminum oxide (IB-F) and silica gel (IB2-F) plates (J. T. Baker). Alumina (EM-Science, AX-0612, 80–200 mesh) and silica gel 60 (EM-Science, 7734, 70–230 mesh) were also purchased from commercial sources. All ligands were synthesized and characterized in the air unless otherwise stated. Preparation and handling of air-sensitive materials were carried out under an argon atmosphere using standard Schlenk techniques. Solvents and solutions were deoxygenated by either repeated freeze-pump-thaw cycles (5 ×), or by bubbling of argon (> 25 min.) directly through the solution. Solid samples were stored and transferred in an MBraun LabMaster 130 inert atmosphere (<1 ppm O₂, <1 ppm H₂O) glovebox under nitrogen atmosphere. NMR spectra were measured either on a Varian NMR instrument at 400 MHz or Bruker 400 MHz spectrometer. All spectra were recorded in 5-mm-o.d. NMR tubes, and chemical shifts were reported as δ values downfield from an internal standard of Me₄Si (¹H). EI and FAB mass spectra were obtained using a VG70S instrument. ESI mass spectra were acquired using a Finnigan LCQDeca ion-trap mass spectrometer equipped with an electrospray ionization source (Thermo Finnigan, San Jose, CA). Samples were dissolved in CH₃OH and introduced into the instrument at a rate of 10 μ L/min using a syringe pump via a silica capillary line. The heated capillary temperature was 250 °C and the spray voltage was 5kV. X-band electron paramagnetic resonance (EPR) spectra were recorded on a Bruker EMX CW-EPR spectrometer controlled with a Bruker ER 041 XG microwave bridge operating at X-band (~9.472 GHz). The low temperature experiments were carried out in tetrahydrofuran (THF) at 77K using a N₂(l) finger dewar. UV-vis spectra were recorded with either a Cary-50 Bio spectrophotometer equipped with a fiber optic coupler (Varian) and a fiber optic dip probe (Hellma: 661.302-QX-UV-2mm-for-low-temperature) or a Hewlett-Packard Model 8453A diode array spectrophotometer equipped with a two-window quartz H.S. Martin Dewar filled with cold MeOH (25 °C to –85 °C) which was maintained and controlled by a Neslab VLT-95 low temp circulator. Spectrophotometer cells used were made by Quark Glass with column and pressure/vacuum side stopcock and 2 mm path length. The copper(I) complex used for low temperature UV-vis and all reactivity studies reported below are the ClO₄[–] and B(C₆F₅)₄[–] salt complexes unless otherwise stated. *While we have experienced no problems in working with perchlorate compounds, they are potentially explosive and care must be taken not to work with large quantities.* Dioxygen was dried by passing through a short column of supported P₄O₁₀ (Aquasorb, Mallinkrodt) and was bubbled into reaction solutions via an 18-gauge, 24-inch-long stainless steel syringe needle. IR spectra were collected using a Mattson Instruments Galaxy series FT-IR (model 4030) that was controlled by the PC program WinFIRST. Elemental analyses were performed by Desert Analytics (Tucson, AZ).

4-(2-propenyloxy)-3-methoxybenzyl alcohol

To a flame-dried 3-neck 2000 mL round-bottom flask equipped with a stir bar and reflux condenser were added in the following order: anhydrous K₂CO₃ (225.0 g, 1.628 mol), 4-hydroxy-3-methoxybenzyl alcohol (200.0 g, 1.297 mol), 1000 mL acetone, and allyl bromide (130 mL, 1.502 mol). The resulting yellow solution was refluxed under argon for 18 hours. After cooling to room temperature, the solvent was reduced *in vacuo* to give a yellow solid, after which 1000 mL dichloromethane was added. The solution was extracted with water (2 × 750 mL), brine (1 × 750 mL), dried over anhydrous Na₂SO₄, filtered, and the resulting solution was concentrated *in vacuo* to give a light yellow solid. Recrystallization of the crude material by dissolving in 600 mL boiling 2:1 diisopropyl/diethyl ethers and subsequent cooling to 0 °C yielded 4-(2-propenyloxy)-3-methoxybenzyl

alcohol (152.0 g, 783 mmol) as a white solid in 60.3% yield. ^1H NMR (CDCl_3): δ 3.89 (s, 3H, OCH_3), 4.62 (m, br, OCH_2 & OH), 5.28 (d, 1H, $J = 10.5$ Hz, allylic H), 5.40 (d, 1H, $J = 17.3$ Hz, allylic H), 6.09 (m, 1H, allylic H), 6.85 – 6.94 (m, 3H, Ar H).

($\pm$)2,7,12-trimethoxy-3,8,13-tris(2-propenyloxy)-10,15-dihydro-5H-tribenzo [a,d,g]cyclononene

To a 3-neck 2000 mL round bottom flask equipped with a stir bar and containing 1250 mL methanol solvent was added 4-(2-propenyloxy)-3-methoxybenzyl alcohol (152.0 g, 783 mmol), yielding a light yellow solution. Upon cooling to 0 °C, 70% perchloric acid (600 mL) was added dropwise over a period of one hour. The resulting red solution was allowed to warm to room temperature, after which a white precipitate formed. After stirring for an additional 18 hours, dichloromethane (1500 mL) was added to quench the reaction, and the dark red organic layer was collected and washed with water (4×1000 mL), brine (2×750 mL), dried over Na_2SO_4 , and filtered. The red-brown solution was then concentrated in *vacuo* to give a sticky red-brown solid, to which 800 mL diethyl ether was added and the mixture stirred. Filtration of the cloudy red-brown solution gave ($\pm$) 2,7,12-trimethoxy-3,8,13-tris(2-propenyloxy)-10,15-dihydro-5H-tribenzo[a,d,g]cyclononene (57.3 g, 108 mmol) as a white solid in 41.5% yield. ^1H NMR (CDCl_3): δ 3.52 (d, 3H, $J = 13.8$ Hz, H_{eq}), 3.83 (s, 9H, OCH_3), 4.58 (s, br, 6H, OCH_2), 4.74 (d, 3H, $J = 13.8$ Hz, H_{ax}), 5.25 (d, 3H, $J = 10.5$ Hz, allylic H), 5.37 (d, 3H, $J = 17.3$ Hz, allylic H), 6.06 (m, 3H, allylic H), 6.79 (s, 3H, H_1), 6.85 (s, 3H, H_2).

($\pm$)2,7,12-Trimethoxy-3,8,13-trihydroxy-10,15-dihydro-5H-tribenzo [a,d,g] cyclononene (Cyclotriguaiacylene, CTG)

To a 3-neck 2000 mL round bottom flask equipped with a stir bar and a reflux condenser was added in the following order: ($\pm$) 2,7,12-trimethoxy-3,8,13-tris(2-propenyloxy)-10,15-dihydro-5H-tribenzo[a,d,g]cyclo-nonene (57.3 g, 108 mmol), 600 mL acetonitrile, 180 mL water, triethylammonium formate (60.0 g, 410 mmol), triphenylphosphine (PPh_3 , 2.03 g, 7.740 mmol), and a catalytic amount of palladium(II) acetate (250 mg, 1.110 mmol). The cloudy yellow mixture was refluxed for 4 hours, after which the acetonitrile was removed via rotary evaporation, and 1500 mL of ethyl acetate was added. The organic layer was filtered, washed with water (2×1000 mL), brine (1×1000 mL), dried over Na_2SO_4 , and again filtered. Concentration resulted in a pale green-yellow solid to which some diethyl ether was added and stirred. Isolation of the CTG (21.0 g, 51.4 mmol) by filtration gave an off-white solid in 47.6% yield. ^1H NMR (CDCl_3): δ 3.48 (d, 3H, $J = 13.7$ Hz, H_{eq}), 3.84 (s, 9H, OCH_3), 4.69 (d, 3H, $J = 13.7$ Hz, H_{ax}), 6.81 (s, 3H, H_1), 6.86 (s, 3H, H_2). EI^+ MS: ($m + \text{H}^+$): m/z 408.

($\pm$) 2,7,12-Trimethoxy-3,8,13-tris((2-(6-chloromethyl) pyridyl)methoxy)-10,15-dihydro-5H-tribenzo[a,d,g]cyclononene (CTV-Halide)

To a flame dried 2-neck 500 mL round bottom flask equipped with a stir bar and rubber septum was added in the following order under argon: K_2CO_3 (5.0 g, 36.1 mmol), 200 mg KI, 500 mg Bu_4NI , CTG (1.5 g, 3.68 mmol), 2,6-bis(chloromethyl)pyridine (6.0 g, 34.1 mmol), and 200 mL high-purity acetone. The cloudy yellow reaction mixture was stirred at room temperature under argon for 5 hours, after which an additional amount of 2,6-bis(chloromethyl)pyridine (6.0 g, 34.1 mmol) dissolved in 75 mL acetone was added under argon flow. After an additional 65 hours of stirring at ambient temperature, the reaction mixture was filtered, and the filtrate was concentrated in *vacuo*. The resulting yellow solid was dissolved in 500 mL dichloromethane, and the clear yellow solution was washed with water (2×500 mL), brine (1×500 mL), dried over Na_2SO_4 , filtered, and reduced via rotary evaporation to give a yellow solid. The crude material was applied to a silica column (80–200 mesh, 4.0 cm. o.d. $\times$ 18.0 cm.) where excess 2,6-bis(chloromethyl)pyridine was eluted

with dichloromethane. Acetone was then allowed to pass down the column, thereby eluting the partially purified CTV-halide product. After concentration, the off-white solid material was further purified using a Chromatotron (eluent: 1.5% EtOH in CH₂Cl₂). Removal of solvent yielded CTV-halide (1.88 g, 2.27 mmol) as a white solid in 62% yield. TLC (silica, 1.5% CH₃OH in CH₂Cl₂) R_f: 0.30. ¹H NMR (CDCl₃): δ 3.43 (d, 3H, J = 13.8 Hz, H_{eq}), 3.74 (s, 9H, OCH₃), 4.67 (d, 3H, J = 13.8 Hz, H_{ax}), 4.68 (s, 6H, CH₂Cl), 5.24 (s, 6H, OCH₂), 6.68 (s, 3H, H₁), 6.81 (s, 3H, H₂), 7.38 (d, 3H, J = 7.80 Hz, Ar H), 7.46 (d, 3H, J = 7.80 Hz, Ar H), 7.71 (t, 3H, J = 7.80 Hz, Ar H). FAB MS (m + H⁺): m/z 826.

(±) 2,7,12-Trimethoxy-3,8,13-tris(6-(tris((2-pyridyl)methyl)amine)-methoxy)-10,15-dihydro-5H-tribenzo[a,d,g]cyclononene (CTV-TMPA)

To a flame dried 2 neck 250 mL round bottom flask equipped with a stir bar was added in the following order under argon: K₂CO₃ (5.0 g, 36.1 mmol), tetrabutylammonium iodide (2.5 g, 6.78 mmol), bis(2-picolyl)amine (2.5 g, 12.6 mmol), 125 mL distilled acetonitrile, diisopropylethylamine (2.5 mL, 14.4 mmol), and CTV-halide (1.2 g, 1.45 mmol). The reaction mixture was heated to 35°C and allowed to stir at this temperature under inert atmosphere overnight. Then, the solvent was removed via rotary evaporation and the yellow-brown oil was dissolved in 500 mL dichloromethane, washed with water (2 × 500 mL), brine (1 × 500 mL), dried over Na₂SO₄ and filtered. To this clear yellow solution was added phthalic anhydride (2.24 g, 15.1 mmol) and the reaction mixture was allowed to stir under argon for two additional hours. The solution was then washed with 1 N NaOH (3 × 500 mL), brine (1 × 500 mL), dried over Na₂SO₄, filtered, and the solvent removed *in vacuo* to give dark yellow oil. Upon dissolving the oil in 10 mL ethyl acetate, a white crystalline precipitate was formed. This precipitate was removed via filtration, and the resulting clear dark yellow solution was applied to an alumina column (unactivated, 80–200 mesh, 4.0 cm. o.d. × 18 cm.) and eluted with 3% CH₃OH in EtOAc. Removal of solvent gave CTV-TMPA (900 mg, 0.68 mmol) as an off-white solid in 47% yield. TLC (alumina, 3% CH₃OH in EtOAc) R_f: 0.30 – 0.35. ¹H NMR (CDCl₃): δ 3.39 (d, 3H, J = 13.7 Hz, H_{eq}), 3.63 (s, 9H, OCH₃), 3.89 (s, br, 18H, NCH₂), 4.63 (d, 3H, J = 13.7 Hz, H_{ax}), 5.20 (s, 6H, OCH₂), 6.63 (s, 3H, H₁), 6.80 (s, 3H, H₂), 7.13–7.68 (m, 27H, Ar H), 8.53 (d, 6H, J = 4.2 Hz, Ar H). FAB MS (m + H⁺): m/z 1316. Anal. Calcd. For C₈₁H₇₈N₁₂O₆: C, 73.95; H, 5.98; N, 12.78. Found: C, 73.60; H, 5.56; N, 11.68. Assignments of H_{ax} and H_{eq} peaks were carried out following literature reports.^{55,56}

[(CTV-TMPA)Cu^I]₃(ClO₄)₃

In a 25 mL Schlenk flask equipped with a stir bar was added under argon CTV-TMPA (300 mg, 0.23 mmol) and [Cu(CH₃CN)₄](ClO₄) (220 mg, 0.69 mmol). Addition of 10 mL deoxygenated acetonitrile resulted in a yellow solution, which was allowed to stir under inert atmosphere for 30 minutes. Deoxygenated diethylether was added until the solution began to turn cloudy, at which point the yellow solution was passed through a coarse porosity Schlenk filter frit into a 100 mL air-free flask. Further addition of deoxygenated diethyl ether (75 mL), followed by vigorous stirring, led to the formation of a yellow precipitate. The clear mother liquor was decanted under a flow of argon and the precipitate was washed with 80 mL deoxygenated diethyl ether. After filtration under argon, the yellow solid was placed under vacuum for two hours which resulted in a free-flowing solid (295 mg, 72% yield). ¹H NMR (CD₃CN): δ 3.61 (d, 3H, J = 13.5 Hz, H_{eq}), 3.72 (s, 9H, OCH₃), 3.96 (s, br, 18H, NCH₂), 4.82 (d, 3H, J = 13.4 Hz, H_{ax}), 5.30 (dd, 6H, J = 12.6 & 31.8 Hz, OCH₂), 7.00 (s, 3H, H₁), 7.15 (s, 3H, H₂), 7.29 – 7.77 (m, 27H, Ar H), 8.51 (s, 6H, Ar H).

[(CTV-TMPA)Cu^I]₃(B(C₆F₅)₄)₃ (1)

CTV-TMPA (0.250 g, 0.190 mmol) and [Cu^I(CH₃CN)₄](B(C₆F₅)₄)⁵⁷ (0.172 g, 0.190 mmol) were placed in a 25 mL Schlenk flask under argon. THF (2 mL) were added under argon to

the mixture of solids to form a light yellow solution. The resulting solution was stirred under argon for 30 min. Pentane (14 mL) was then added to the solution to precipitate a light yellow solid. The supernatant was decanted, and the solid was recrystallized three times from THF/pentane under argon and dried under vacuum (1 h), giving 0.458 g of light yellow powder (68% yield). Anal. Calcd. For $C_{153}H_{78}B_3Cu_3F_{60}N_{12}O_6$: C, 51.86; H, 2.22; N, 4.74. Found: C, 51.58; H, 2.92; N, 4.10.

Electrochemistry

Cyclic voltammetry was carried out with a Bioanalytical Systems BAS-100B electrochemistry analyzer. The sample cell used was a standard three-electrode system with platinum wire auxiliary as the counter electrode. A glassy carbon electrode (GCE, BAS MF 2012) was used as the working electrode. The reference electrode was Ag/Ag⁺. Dimethylformamide (DMF) as solvent was deoxygenated by argon bubbling. The measurements were performed at room temperature in DMF containing 0.1 M tetrabutylammonium hexafluorophosphate (TBAHP) and 1 – 0.1 mM copper complex.

Reaction of [(CTV-TMPA)Cu^I]₃(B(C₆F₅)₄)₃ (**1**) with [Fe^{III}Cp₂]⁺

Complex [(CTV-TMPA) Cu^I]₃(B(C₆F₅)₄)₃ (**1**) (0.020 g, 0.006 mmol) was dissolved in 3.0 mL THF and taken in a UV-vis cuvette inside the inert atmosphere box. On the benchtop, an initial spectrum was recorded. In a separate 5 mL Schlenk flask in the glovebox, [Fe^{III}Cp₂]PF₆ (Aldrich; 0.050g, 0.151 mmol) was dissolved in 0.5 mL degassed THF. A total of 56 μL (0.017 mmol) of the THF solution of [Fe^{III}Cp₂]PF₆ was added in the course of 5 min (~9 μL each time) via microliter syringe to the copper(I) solution **1**. The solution was purged with Ar after each addition of ferrocenium ion and a UV-vis spectrum recorded. An EPR of this solution was recorded. EPR (9.471 GHz, MeTHF, 77K): $g_{||} = 2.246$, $g_{\perp} = 2.046$, $A_{||} = 152$ G, $A_{\perp} = 29.3$ G. The EPR of the previously reported crystallographically characterized [($(L^{CH_2OMe})Cu^{II}(Cl)_2$)(B(C₆F₅)₄)₂]⁹³ (0.180 g, 0.080 mmol) complex in 3.1 mL THF was compared with the above EPR of the fully oxidized **1** under similar experimental conditions. In a separate UV-vis cuvette, similar experiments were carried out and the addition of 1 more equiv of [Fe^{III}Cp₂]⁺ solution did not cause further UV-vis or EPR spectral changes.

[(CTV-TMPA)Cu^{II}(Cl)]₃(B(C₆F₅)₄)₃ (**1-Cl**)

[(CTV-TMPA)Cu^I]₃(B(C₆F₅)₄)₃ (**1**) (0.080 g, 0.023 mmol) was placed in a 25 mL Schlenk flask under argon. Six drops of degassed CHCl₃ were added to THF (2.0 mL, degassed) and the solvent mixture was added to the copper(I)-complex containing flask. Under argon, the mixture was stirred for 3 hr while the solution turned to green. Pentane (15 mL, degassed) was then added to precipitate a solid. The copper complex obtained was further recrystallized three times from THF/pentane and dried under vacuum (2 h), giving 0.053 g of light yellow powder (64% yield). EPR (9.471 GHz, THF, 77 K): $g_{||} = 2.23$, $g_{\perp} = 2.05$, $A_{||} = 143$ G, $A_{\perp} = 33.5$ G. Anal. Calcd. For [(CTV-TMPA)Cu^{II}(Cl)]₃(B(C₆F₅)₄)₃ (**1-Cl**), $C_{153}H_{78}B_3Cl_3Cu_3F_{60}N_{12}O_6$: C, 50.35; H, 2.15; N, 4.61. Found: C, 50.16; H, 2.12; N, 4.32. A cyclic voltammetry study and ESI-MS analysis (with the perchlorate salt) were also carried out.

Generation of copper-carbonyl complex from the reaction of CO and **1**: quantitative measurement of the release of CO upon addition of PPh₃

Bubbling CO through a THF solution of [(CTV-TMPA)Cu^I]₃(B(C₆F₅)₄)₃ (**1**) generates the corresponding carbonyl adduct [(CTV-TMPA) Cu^I₃(CO)₃](B(C₆F₅)₄)₃ (**1-CO**). IR spectroscopy: $\nu_{CO} = 2094$ cm⁻¹. ¹H NMR (DMSO): δ 3.56 (d, 3H, H_{eq}), 3.75 (s, 9H, OCH_3), 4.2 (s, br, 18H, NCH_2), 4.86 (d, 3H, H_{ax}), 5.37 (6H, OCH_2), 7.14 (s, 3H, H_1), 7.31

(s, 3H, $\underline{\text{H2}}$), 7.36 – 7.89 (m, 27H, Ar $\underline{\text{H}}$), 8.55 (s, 6H, Ar $\underline{\text{H}}$). Anal. Calcd. For [(CTV-TMPA)Cu^{II}₃(CO)₃](B(C₆F₅)₄)₃ (**1-CO**), C₁₅₆H₇₈B₃Cu₃F₆₀N₁₂O₉: C, 51.65; H, 2.17; N, 4.63. Found: C, 51.44; H, 2.54; N, 4.13.

To confirm that all three copper(I) sites bind CO, PPh₃ was added to **1-CO** and the CO release was quantitated. In a drybox, [(CTV-TMPA)Cu^I₃](B(C₆F₅)₄)₃ (0.0025 g, 0.0007 mmol) was dissolved in 10 mL THF within a UV-vis cuvette. Out on benchtop, the cuvette was cooled to –90 °C in the cooling Dewar flask of the UV-vis instrument. Carbon monoxide was bubbled through the solution for ~5 s and excess CO was removed completely via 5 vacuum/argon cycles. Also, inside a drybox within a separate 5 mL Schlenk flask, PPh₃ (0.0055 g, 0.020 mmol) was dissolved in 1 mL degassed THF. A total of a 100 μ L (0.0020 mmol) of PPh₃ solution was added via microliter syringe to the UV-vis cuvette containing the solution of **1-CO**. A 100 μ L solution of (F₈TPP)Fe^{II}(THF)₂ {F₈TPP = tetrakis(2,6-difluorophenyl)porphyrinate(2-)} (0.0018 g, 0.021 mmol) was added via a microliter syringe to the resulting solution. UV-vis spectrum of this final solution was recorded and was compared with the UV-vis spectrum generated from a separate solution of [(F₈TPP)Fe^{II}(THF)₂] (0.0018 g, 0.021 mmol) in 10.2 mL THF which had been bubbled with CO. A ~2.9 equiv CO removal upon addition of 3 equiv PPh₃ to one equiv **1-CO** was observed.

Reaction of [(CTV-TMPA)Cu^I₃](B(C₆F₅)₄)₃ (**1**) with O₂

A 3.0 mL THF solution of **1** (0.015 g, 0.0042 mmol) was taken in a UV-vis cuvette assembly under argon; this was cooled to –80 °C and an initial UV-Vis spectrum was recorded. Dioxygen was bubbled for 30 s through the solution using a long syringe needle and the new spectrum recorded upon removal of excess O₂. The deep purple product is formulated as **1-O₂** (see Discussion).

Thermal Transformation Product of **2-O₂**

[(CTV-TMPA)Cu^I₃](B(C₆F₅)₄)₃ (**1**) (0.009 g, 0.0025 mmol) was dissolved in 4.0 mL THF and transferred to a UV-vis cuvette assembly inside the MBraun glovebox. Outside on the benchtop, the cuvette was cooled to –80 °C and an initial spectrum was recorded. Dioxygen was bubbled through the solution with a long syringe needle. This results in an immediate color change from colorless to deep purple (excess dioxygen removed). The UV-vis spectra were recorded at –80 °C with respect to time.

In a separate 25 mL Schlenk flask, [(CTV-TMPA)Cu^I₃](B(C₆F₅)₄)₃ (**1**) (0.044 g, 0.012 mmol) was dissolved in 2 mL THF and O₂ was bubbled through the solution upon cooling to –80 °C. This was allowed to warm to room temperature, and then pentane (20 mL) was then added to precipitate an intensely purple solid. The copper complex obtained was further recrystallized four times from THF/pentane and dried under vacuum (4 h), giving 0.033 g of a dark purple powder (76 % yield). Anal. Calcd. For [(CTV-TMPA)Cu^{II}₃(OH)₃](B(C₆F₅)₄)₃ (**1-Dec**), C₁₅₃H₈₁B₃Cu₃F₆₀N₁₂O₉: C, 51.13; H, 2.27; N, 4.68. Found: C, 50.95; H, 2.37; N, 5.15. Note, however that and IR spectrum (Nujol mull) did not reveal an expected O-H stretch in the 3400–3700 cm^{–1} region. An EPR spectrum showed a single broad isotropic signal, see Supporting Information. We did not attempt to obtain mass spectrometric data.

Synthesis of [(CTV-TMPA)Cu^I₃(PPh₃)₃](B(C₆F₅)₄)₃ (**1-PPh₃**)

A 1.5 mL THF solution of [(CTV-TMPA)Cu^I₃](B(C₆F₅)₄)₃ (**1**) (0.067 g, 0.019 mmol) was taken in a 25 mL Schlenk flask and PPh₃ (0.015 g, 0.058 mmol) was added (100 μ L THF) anaerobically via a microliter syringe. The resulting solution was stirred under argon for 20 min. Pentane (15 mL) was then added to the solution to precipitate a white solid. The supernatant was decanted, and the solid was recrystallized four times from THF/pentane

under argon and dried under vacuum (3 h), giving 0.068 g of white powder (84 % yield). Anal. Calcd. For $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3(\text{PPh}_3)_3](\text{B}(\text{C}_6\text{F}_5)_4)_3$ (**1-PPh₃**), $\text{C}_{207}\text{H}_{123}\text{B}_3\text{Cu}_3\text{F}_{60}\text{N}_{12}\text{O}_6\text{P}_3$: C, 57.42; H, 2.86; N, 3.88. Found: C, 57.61; H, 3.01; N, 3.51. ^1H NMR (DMSO): 3.49 (d, 3H, H_{eq}), 3.56–4.33 (m, 27H, OCH_3 and NCH_2), 4.76 (d, 3H, H_{ax}), 5.04 (6H, OCH_2), 7.01 (s, 3H, H_1), 7.10 (s, 3H, H_2), 7.19 – 7.91 (m, 42H, Ar H), 8.42 (s, 6H, Ar H).

^{31}P -NMR study of $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3(\text{PPh}_3)_3](\text{B}(\text{C}_6\text{F}_5)_4)_3$ (**1-PPh₃**)

Inside the glovebox, $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3(\text{PPh}_3)_3](\text{B}(\text{C}_6\text{F}_5)_4)_3$ (**1-PPh₃**) (0.014 g, 0.0032 mmol) and $\text{P}(\text{Mes})_3$ (0.0034 g, 0.0098 mmol) was taken into a NMR sample tube and was dissolved in degassed d_8 -THF. Out on the benchtop, with a rubber septum on top to maintain anaerobic conditions, a ^{31}P -NMR was recorded. ^{31}P -NMR (d_8 -THF): δ 3.79 (3P, **1-PPh₃**), –35.92 (3.093 P, PMes_3).

Generation of disulfido-dicopper (II) complex, $[(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3]_2(\mu\text{-1,2-S}_2^{2-})_3]^{6+}$ (**1-S**)

$[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3](\text{B}(\text{C}_6\text{F}_5)_4)_3$ (**1**) (0.0192g, 0.0054 mmol) was dissolved in 3.8 mL THF under argon in a UV-vis cuvette with a glass stopper. Solid elemental sulfur (0.0017 g) was dissolved in 0.5 mL THF upon stirring in a 5 mL Schlenk flask under argon. This sulfur containing solution was cooled to -80°C using an acetone/dry-ice bath. The glass stoppers from the cuvette and the flask were interchanged for rubber septa. From the stock solution, 50 μL of the sulfur containing solution (0.0053 mmol) was added to the solution in the cuvette. Purging argon for 5 s caused thorough mixing of sulfur solution with the copper(I) solution. The resulting solution was kept at -80°C for 5 min and a UV-vis spectrum was recorded. Three further additions of the sulfur solution (50 μL each time) accompanied by Ar bubbling and recording of UV-vis spectra after 10 minutes were carried out. Further addition of sulfur solution did not cause any observable UV-vis change. A nice purple colored solution ($\lambda_{\text{max}} = 544\text{ nm}$) fully formed, designated as complex **1-S** (See Results and Discussion). Species **1-S** is stable at -80°C ; upon warming to room temperature it loses its UV-vis intensity, whereas based on the UV-vis changes recooling to -80°C (under Ar) results in complete reformation of **1-S**. A blackish-purple colored copper complex was generated upon warming the -80°C **1-S** solution to RT. This solution was filtered with Whatman filter paper (Glass Microfibre Filters, GF/C, Circles, 24 mm $\varnothing$, Catalog No. 1822024) plugged in pipette under argon and the filtration was repeated five times with a new filter paper each time to confirm absolute removal of unreacted (or released upon warming) elemental sulfur from the reaction solution. A pentane solution (20 mL) was added to precipitate the copper product which was further recrystallized four times from THF/pentane under argon and then dried under vacuum (6 h). Anal. Calcd. For $[(\text{CTV-TMPA})\text{Cu}_3\text{S}_3](\text{B}(\text{C}_6\text{F}_5)_4)_3$: $\text{C}_{153}\text{H}_{78}\text{B}_3\text{Cu}_3\text{F}_{60}\text{N}_{12}\text{O}_6\text{S}_3$; C, 50.49; H, 2.16; N, 4.62; S, 2.64. Found: C, 49.26; H, 2.29; N, 4.11; S, 3.3. The high S and low C & N content, despite many recrystallizations (with filtration), may indicate the presence of some polysulfane complex impurities.

Resonance Raman (rR) spectroscopy

Resonance Raman spectroscopic measurements were undertaken using a Princeton Instruments ST-135 back-illuminated CCD detector on a Spex 1877 CP triple monochromator with 1200, 1800, and 2400 grooves/mm holographic spectrograph gratings. The excitation was provided by Coherent 190C-K Kr+ and Innova Sabre 25/7 Ar+ CW lasers. The spectral resolution was $< 2\text{ cm}^{-1}$. Samples were prepared in standard low-grade NMR tubes. Sample concentrations were approximately 1.0 mM and 1.53 mM in copper for dioxygen reactivity. Sample concentrations were approximately 2.45 mM in copper for sulfur reactivity. All the samples were run at 77 K in a liquid N_2 finger Dewar (Wilmad).

The precursor copper(I) sample solutions were prepared in the glovebox in NMR tubes under N₂ and were cooled to −80 °C on the benchtop, after which O₂ bubbling was carried out via syringe. The solutions were frozen in liquid N₂ and sent to Stanford University for rR spectroscopic interrogation.

Results and Discussion

Synthesis of Cyclotrivenatrylene (CTV) Based Trinucleating Ligand, CTV-TMPA

As mentioned, we designed this tripodal ligand system based on the building block cyclotrivenatrylene (2,3,7,8,12,13-hexamethoxy-10,15-dihydro-5H-tribenzo[a,d,g]cyclononene) to promote close metal-metal interactions. This molecule has a rigid, bowl-like shape and possesses three-fold symmetry. Functionalized CTV ligands have for example been previously used to provide sub-site differentiation for [4Fe-4S] protein model compounds,⁵⁸ and this structural motif seems well-suited for metal-cluster modeling.⁵⁰⁻⁵⁸

The functionalization of the cyclotrivenatrylene framework provides a chiral ligand system CTV-TMPA; the conformational inversion of their nine-membered ring is very slow at room temperature.⁴⁷ The CTV-TMPA ligand employs the TMPA {= tris(2-pyridylmethyl)amine} tetradentate copper chelate whose corresponding mononuclear Cu^I/O₂ chemistry is well established with respect to not only the types of dioxygen adducts possible, but also their kinetics and thermodynamics of formation.^{9,52-54} In order to append a copper binding site to the CTV periphery, it was determined that the best course of action was to proceed via functionalization of the cyclotrigenatrylene (CTG) precursor, which possesses a phenolic group ideal for nucleophilic substitution reactions (Scheme 1). We have previously demonstrated the ability to modify tetraarylporphyrins which contain meso-substituted 2-hydroxyphenyl arms with the copper-chelate TMPA {TMPA = tris(2-pyridylmethyl)amine},^{61,62} and here we employed the same methodology and reaction conditions. The addition of 2,6-bis(chloromethyl)pyridine to CTG results in the formation of CTV-halide, whose alkylhalide functional group allows for the attachment of a variety of primary and secondary amines. We then chose bis(2-picolyl)amine for the synthesis of a trinucleating ligand, CTV-TMPA (Scheme 1).

Synthesis and characterization of a tricopper(I) complex

The light yellow tricopper(I) complex of CTV-TMPA, [(CTV-TMPA)Cu^I]₃(B(C₆F₅)₄)₃ (**1**) (Scheme 2), was generated by the addition of three equivs of [Cu^I(CH₃CN)₄](B(C₆F₅)₄) to one equiv CTV-TMPA ligand in deoxygenated THF solvent under argon followed by precipitation with pentane. Isolation of a pure solid was achieved by recrystallization from THF/pentane. The copper(I) complex was also synthesized using the perchlorate (ClO₄[−]) counteranion. The copper complexes [(CTV-TMPA)Cu^I]₃³⁺ (**1**) were characterized by elemental analysis and ¹H-NMR spectroscopy, see Experimental Section.

Mononuclear analog for copper ion in CTV-TMPA

CTV-TMPA possesses three TMPA units in which one of the three pyridine ring of TMPA is 6-CH₂-O-R substituted and the three units are connected to a bowl-shaped cyclotrivenatrylene moiety. We, previously, reported⁵² the synthesis and characterization of the mono 6-CH₂-O-CH₃ substituted TMPA ligand, L^{CH₂OMe} (diagram) which is indeed a close mimic of any given mononuclear ligand portion of CTV-TMPA. X-ray structures of the copper complexes [(L^{CH₂OMe})Cu^I]⁺ (**2**), [(L^{CH₂OMe})Cu^{II}(Cl)]₂²⁺ and [(L^{CH₂OMe})Cu^{II}(Br)]₂²⁺ were described.⁵² Several attempts to crystallize copper-complexes with CTV-TMPA were unsuccessful, thus we sought to produce evidence for tricopper binding and chemistry by chemical approaches.

Electrochemistry

The half-wave potentials for the copper(I) complexes, [(CTV-TMPA) Cu^I₃]³⁺ (**1**) and [(L^{CH₂OMe})Cu^I]⁺ (**2**) were measured under argon in dimethylformamide (DMF) by cyclic voltammetry (CV). In DMF, **1** and **2** displayed a single quasi-reversible redox behavior with peak-current ratios of i_{pa}/i_{pc} = 0.84 & 0.87, respectively (Table 1). Peak separations were less than 150 mV at a scan rate of 100 mV/s. The ferrocene-ferrocenium couple under the same conditions showed ΔE_p = 85 mV and $E_{1/2}$ = + 55 mV vs Ag/AgNO₃. A typical CV scan for [(CTV-TMPA) Cu^I₃]³⁺ (**1**) is depicted in Figure 2.

The electrochemical behavior of [(CTV-TMPA)Cu^I₃]³⁺ (**1**) can be explained by having the presence of three non-interacting copper ion centers where the redox process for each occurs at about the same potential. The appearance of only one CV wave has been previously reported for a number of multi-copper systems.^{63–66} Generally, one wave is seen if ΔE_p between two separate processes (i.e. Cu^{II}-Cu^{II} + e[−] → Cu^{II}-Cu^I + e[−] → Cu^I-Cu^I) is < ~ 100–200 mV. We have not attempted a more detailed analysis which might be used to estimate $E_{1/2}$ for the individual redox reactions,^{63,67} however this has been carried out in a number of cases.⁶⁵ In one example, for a tri-platinum complex bound directly to the CTV catechol framework, Bohle and Stasko⁶⁸ observed three separate reversible redox couples, assigned as individual Pt-catecholate-to-Pt-seminquinoid species.

[(CTV-TMPA)Cu^I₃]³⁺ (**1**) has very similar $E_{1/2}$ value (Table 1) to that of the close mononuclear mimic [(L^{CH₂OMe})Cu^I]⁺ (**2**). These $E_{1/2}$ values are ~110 mV more positive compared to the ‘parent’ [(TMPA)Cu^I(CH₃CN)]⁺ (**3**) complex which lacks a single 6-pyridyl substituent. Factors that can influence ligand-copper complex $E_{1/2}$ values include (i) nature of chelating ligands, (ii) donor atom types and (iii) the geometry around tetra or pentacoordinate complexes.^{5,7,69–71} A steric effect arises due to the 6-CH₂-O-R substituent (accompanied by a change of geometry) is the likely cause of the positive shift in $E_{1/2}$ seen in [(CTV-TMPA)Cu^I₃]³⁺ (**1**) and [(L^{CH₂OMe})Cu^I]⁺ (**2**) relative to parent [(TMPA)Cu^I(CH₃CN)]⁺ (**3**) complex. Such suppositions come about from the study of similarly constructed ligand-copper systems.^{72–74}

The ease of electrochemical oxidations of the three copper centers in $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**) correlates in a general way with the $[(\text{L}^{\text{CH}_2\text{OMe}})\text{Cu}^{\text{I}}]^+$ (**2**) reactivity with O_2 , as described below. $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**) does react with O_2 at low temperature forming copper-dioxygen adducts. However, their tendency to give 1:1 and 2:1 adducts depends on geometric effects within the CTV architecture (*vide infra*).

Addition of 3 $[\text{Fe}^{\text{III}}\text{Cp}_2]^+$ to $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**)

The tricopper(I) complex **1** can be oxidized to all copper(II) via addition of $[\text{Fe}^{\text{III}}\text{Cp}_2]^+$ as oxidant (Scheme 3). UV-vis absorption monitoring of solutions of $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**) after addition of $[\text{Fe}^{\text{III}}\text{Cp}_2]^+$ shows that three equiv are needed to complete the oxidation of **1**; after that, no further spectral changes are observed when further (i.e., excess) $[\text{Fe}^{\text{III}}\text{Cp}_2]^+$ is added (Figure 4). EPR spectra of the resulting fully oxidized copper species, $[(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3]^{3+}$ (**1-Ox**) (Figure 5), closely correlated in intensity (i.e., semi-quantitative spin integration) with EPR spectra of 1.5 equiv *dimeric* copper(II) complex $[(\text{L}^{\text{CH}_2\text{OMe}})\text{Cu}^{\text{II}}(\text{Cl})]_2^{2+}$ (i.e. 3 equiv Cu(II)), which in solution breaks up into monomeric species. Thus, we can conclude that all three copper(II) centers are located in a similar tetra-coordinate environment originating from the three equiv copper centers in this CTV-TMPA ligand system. Species **1-Ox** is insensitive to the presence of O_2 .

Generation of $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3(\text{PPh}_3)_3]^{3+}$ (**1-PPh₃**)

Triphenylphosphine is well known as a very good ligand for Cu(I) complexes, and it is sometimes used in the establishing the nature of Cu_nO_2 reactivity or other metal-dioxygen complexes.^{9,63,75} The affinity of PPh_3 for copper(I) originates from the soft nature of both species; once a copper(I)- PPh_3 adduct is formed, it is usually unreactive toward O_2 .^{9,63,75}

The triphenylphosphine adduct of **1**, $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3(\text{PPh}_3)_3]^{3+}$ (**1-PPh₃**) may be generated by addition of three equiv PPh_3 to the copper(I) solution of THF and can be isolated by addition of pentane. This product, **1-PPh₃** is stable and insensitive to O_2 . We have taken advantage of ^{31}P -NMR spectroscopy (with PMes_3 internal standard added) to confirm the number of PPh_3 equivalents binding to **1** (Scheme 4). A 1:3 mixture of **1-PPh₃** and $\text{P}(\text{Mes})_3$ shows that the NMR integration ratios are 1: 1.031 for the two peaks observed at 3.79 (**1-PPh₃**) and -35.92 (PMes_3) ppm; thus indicating three PPh_3 binding sites in $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**) (Figure S1),⁷⁶ as expected.

Generation of $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3(\text{CO})_3]^{3+}$ (**1-CO**) and CO Release by Addition of PPh_3

The carbon monoxide adduct of **1**, $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3(\text{CO})_3]^{3+}$ (**1-CO**) can be generated by bubbling CO through the copper(I) solution (Scheme 5) and the product obtained was characterized by elemental analysis, ^1H -NMR and IR spectroscopies. Addition of PPh_3 and quantitative detection of the carbon monoxide released allowed us to confirm that **1-CO** possesses three CO molecules, one per copper. Thus, the amount of CO released from the THF solution of $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3(\text{CO})_3]^{3+}$ (**1-CO**) upon PPh_3 addition was determined following the addition of 3 equiv of the iron-porphyrinate complex $[(\text{F}_8\text{TPP})\text{Fe}^{\text{II}}(\text{THF})_2]$ $\{\text{F}_8\text{TPP} = \text{tetrakis}(2,6\text{-difluorophenyl})\text{porphyrinate}(2-)\}$ $\{\text{UV-vis (THF; } \lambda_{\text{max}}, \text{ nm): 421 nm, 542 nm}\}$.⁷⁷ The resultant carbonmonoxy-THF complex $[(\text{F}_8\text{TPP})\text{Fe}^{\text{II}}(\text{CO})(\text{THF})]$ $\{\text{UV-vis (THF; } \lambda_{\text{max}}, \text{ nm): 411 nm, 525 nm}\}$ (also compared with same concentration of $[(\text{F}_8\text{TPP})\text{Fe}^{\text{II}}(\text{THF})_2]$ bubbled with CO) indicated the release of ~2.9 equiv of CO for one equiv $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3(\text{CO})_3]^{3+}$ (**1-CO**) (Figure 6). Thus, **1-CO** possesses three CO-bound cuprous centers. Consistent with the results described above (i.e., for electrochemistry and the properties of copper in **1-PPh₃** and **1-Ox**) the CO bound centers in **1-CO** are comparable since a single IR stretch ($\nu_{\text{C=O}} = 2094 \text{ cm}^{-1}$) is observed.⁷⁶ The ν_{CO} value is typical for that observed for other TMPA analogues ($\nu_{\text{CO}} = 2091\text{--}2094 \text{ cm}^{-1}$) and the closest mimic $\text{L}^{\text{CH}_2\text{OMe}}$ ($\nu_{\text{CO}} = 2094 \text{ cm}^{-1}$).^{52,78} This and previous studies indicate an

overall tetra-coordinated $\text{Cu}^{\text{I}}\text{N}_3(\text{CO})$ structure (at least as a solid) with one pyridyl group (likely the 6-substituted pyridyl arm) dangling. When all pyridyl groups bond, giving a $\text{Cu}^{\text{I}}\text{N}_4(\text{CO})$ coordination, the ν_{CO} drops to the 2075 cm^{-1} range due to increased Cu^{I} -to-CO back donation.^{78,79}

Generation of $[(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3(\text{Cl})_3](\text{B}(\text{C}_6\text{F}_5)_4)_3$ (**1-Cl**) via a dechlorination reaction

As we have seen many times previously and is known in other cases, TMPA-based copper(I) complexes are reactive towards organohalides, resulting in the production of halide-copper(II) products.^{52,72} A recent report from our laboratories suggests such reactions may occur via organocopper intermediates.⁸⁰ Here, we utilized this organohalide reaction chemistry (eq. 1) to generate a chloride complex $[(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3(\text{Cl})_3](\text{B}(\text{C}_6\text{F}_5)_4)_3$ (**1-Cl**). Exposure of **1** to CHCl_3 (in THF) at room temperature led to an immediate reaction (according to eq. 1) and **1-Cl** was isolated in high yield.

EPR spectroscopy (Figure S3), elemental analysis, cyclic voltammetry (Figure S4) and ESI-MS (Figure S10 and S11) studies of **1-Cl** were carried out.⁷⁶ An ESI-MS spectrum of $[(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3(\text{Cl})_3](\text{ClO}_4)_3$ (in CH_3CN at RT) showed strong peaks in the range $m/z = 1808\text{--}1814$ for $(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3(\text{Cl})_3(\text{ClO}_4)_2^+$ and the observed peak pattern matched well with that predicted.⁷⁶ The bromide analog $[(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3(\text{Br})_3]^3+$ was also synthesized following a similar method involving dehalogenation of CHBr_3 with $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^3+$; the EPR (Figure S8) and ESI-MS (Figure S12 and S13) spectra was recorded. ESI-MS spectra of $[(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3(\text{Br})_3](\text{ClO}_4)_3$ (in CH_3CN at RT) showed strong peaks at $m/z = 1940\text{--}1948$ for $(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3(\text{Br})_3(\text{ClO}_4)_2^+$ and the observed peak pattern matched well with that theoretically predicted.⁷⁶ The synthesis of $[(\text{L}^{\text{CH}_2\text{OMe}})\text{Cu}^{\text{II}}(\text{Cl})]\text{B}(\text{C}_6\text{F}_5)_4$ was previously reported. As related to present study, EPR and cyclic voltammetry studies of $[(\text{L}^{\text{CH}_2\text{OMe}})\text{Cu}^{\text{II}}(\text{Cl})]\text{B}(\text{C}_6\text{F}_5)_4$ were carried out and the peak pattern and positions were found to be very similar to that for $[(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3(\text{Cl})_3]^3+$ (**1-Cl**).

Reaction of O_2 with $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^3+$ (**1**)

The copper chemistry of the mononuclear mimic ($\text{L}^{\text{CH}_2\text{OMe}}$) of the copper sites in CTV-TMPA has been recently reported. Copper(I)/ O_2 reactivity of $\text{L}^{\text{CH}_2\text{OMe}}$ leads to formation of a dinuclear purple species (similar to parent TMPA) formulated as $[(\text{L}^{\text{CH}_2\text{OMe}})\text{Cu}^{\text{II}}]_2(\mu\text{-}1,2\text{-O}_2^{2-})]^{2+}$ (**2-O₂**; see diagram below) based on its UV-vis $\{\lambda_{\text{max}} = 540\text{ nm} (\epsilon, 9550\text{ M}^{-1}\text{cm}^{-1}) \text{ and } 610\text{ nm, sh, } (\epsilon, 6500\text{ M}^{-1}\text{cm}^{-1})\}$ and resonance Raman (rR) ($\nu_{(\text{O-O})} = 848 (\Delta[^{18}\text{O}_2] = -47)\text{ cm}^{-1}$, $\nu_{(\text{Cu-O})} = 550 (\Delta[^{18}\text{O}_2] = -26\text{ cm}^{-1}$ in THF) spectroscopic properties.⁸¹ We have also recently characterized the first formed mononuclear $\text{Cu}^{\text{I}}/\text{O}_2$ initial adduct with a TMPA derivative, $\text{NMe}_2\text{-TMPA}$ {tris(2-(4-dimethylaminopyridyl)methyl)amine}, leading to a brilliant green colored copper-superoxo species $[(\text{NMe}_2\text{-TMPA})\text{Cu}^{\text{II}}(\text{O}_2^-)]^+$ (see diagram below): $\lambda_{\text{max}} = 418 (\epsilon = 4300\text{ M}^{-1}\text{cm}^{-1})$, $615 (\epsilon = 1100\text{ M}^{-1}\text{cm}^{-1})$, and $767 (\epsilon = 840\text{ M}^{-1}\text{cm}^{-1})$ and rR spectroscopy ($\nu_{(\text{O-O})} = 1121 (\Delta[^{18}\text{O}_2] = -63)\text{ cm}^{-1}$, $\nu_{(\text{Cu-O})} = 472 (\Delta[^{18}\text{O}_2] = -20\text{ cm}^{-1}$ in THF).⁸²

Following oxygenation of $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**) in THF at -80°C , there was a rapid change from colorless to an intense purple colored solution. The UV-vis spectrum of this product **1-O₂** (Figure 7) includes intense feature at 543 and 427 nm. The former absorption, along with lower broad energy closely resembles the distinctive pattern known for the 'parent' complexes $[(\text{TMPA})\text{Cu}^{\text{II}}]_2(\mu\text{-1,2-O}_2^{2-})^{2+}$ (**3-O₂**) and $[(\text{L}^{\text{CH}_2\text{OMe}})\text{Cu}^{\text{II}}]_2(\mu\text{-1,2-O}_2^{2-})^{2+}$ (**2-O₂**) as well as other structurally similar *trans*-peroxo binuclear Cu_2O_2 -adducts.^{5,81} This is confirmed by resonance Raman (rR) spectroscopy, *vide infra*.

Resonance Raman Spectroscopy of Solutions from $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**)/O₂

Resonance Raman spectra of **1-O₂** were collected with $\lambda_{\text{excit}} = 530\text{ nm}$ with $^{16}\text{O}_2$ and $^{18}\text{O}_2$ isotopic substitution. For **1-O₂** sample concentrations of $\sim 1.0\text{ mM}$ in THF, two isotope-dependent stretches were observed at 842 and 825 cm^{-1} , which shift to 795 and 778 cm^{-1} upon $^{18}\text{O}_2$ substitution and are assigned as peroxo O-O stretches (Figure 8). In addition, there are two isotope-sensitive stretches observed at 506 and 548 cm^{-1} , which shift to 480 and 522 cm^{-1} and are assigned as symmetric Cu-O stretches (Figure 9). An $^{18}\text{O}_2$ -insensitive stretch is observed at 438 cm^{-1} and is assigned as a Cu-N_{amine} stretch. Cu-nitrogen ligand vibrations could be resolved in some cases (*vide infra*), but here for complex **1-O₂**, an overlapping solvent band from THF (258 cm^{-1}), precludes its observation.

The frequencies and 18-O shifts observed for **1-O₂** correlate well to those of an $\mu\text{-1,2-end-on}$ peroxo species, Table 2.⁵ For $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**)/O₂ derived solutions, both UV-vis absorption and rR data in the peroxo region (with $\lambda_{\text{excit}} = 530\text{ nm}$) are very similar to those observed for the parent $[(\text{L}^{\text{CH}_2\text{OMe}})\text{Cu}^{\text{II}}(\mu\text{-1,2-O}_2^{2-})]_2^{2+}$ (**2-O₂**) complex. As was previously observed for the parent $\text{L}^{\text{CH}_2\text{OMe}}$ complex, shifts in the rR and absorption peak positions relative to $[(\text{TMPA})\text{Cu}^{\text{II}}(\mu\text{-1,2-O}_2^{2-})]_2^{2+}$ (**3-O₂**) arise from the presence of 6-substituted pyridyl arms of the TMPA.⁸¹

The UV-vis spectra of solutions generated from O₂ reaction with $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**) also show a prominent absorption feature at 427 nm (Figure 7). Resonance Raman spectra were collected with $\lambda_{\text{excit}} = 413\text{ nm}$ and an O-O stretch at 1129 cm^{-1} , shifting to 1069 cm^{-1} upon $^{18}\text{O}_2$ substitution, was observed (Figure 10). This stretching frequency and isotopic shift, along with the position and intensity of the 427 nm absorption, are consistent with those of a mononuclear superoxo-copper(II) complex.^{5,82-83} A Cu-O stretch was also observed for the complex with 413 nm excitation at 463 cm^{-1} , shifting to 436 cm^{-1} upon $^{18}\text{O}_2$ substitution (Figure 11).

Interestingly, rR spectra collected with $\lambda_{\text{excit}} = 530\text{ nm}$ on samples at higher concentration (1.53 mM of **1**) show only one O-O stretch at 825 cm^{-1} ($\Delta(^{18}\text{O}_2) = -47$) (Figure 12) and a Cu-O stretching at 506 cm^{-1} ($\Delta(^{18}\text{O}_2) = -26$) (Figure 13).⁸⁴ Thus, at this higher concentration, only one peroxo species is observed. This result confirms that for the 1.0 mM sample concentration, the 506 cm^{-1} symmetric Cu-O and the 825 cm^{-1} O-O stretches derive from one peroxo complex, while the 548 cm^{-1} symmetric Cu-O and 842 cm^{-1} O-O stretches derive from a second, distinct peroxo complex.

The UV-vis and resonance Raman spectroscopic data indicate that both the superoxo and peroxo species form from $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**)/O₂ chemistry. In THF at 1.0 mM concentration, one type of superoxo (O_2^-) and two different types of peroxo ($\mu\text{-1,2-O}_2^{2-}$) species are present (Figure 8 and 9). Thus, a total of three different species are formed from **1**/O₂. Interestingly, oxygenation with a more concentrated copper solution (1.53 mM of **1**) in THF results in only one of the peroxo species and the superoxo species, as evidenced by rR spectroscopy (*vide supra*). The two different $\mu\text{-1,2-peroxos}$ observed at a lower concentration are likely an intra- and intermolecular species, while at higher concentration the observed peroxo is likely an *intermolecular* species. This type of concentration

dependent behavior has been previously observed.⁸⁵ Thus, to our knowledge, oxygenation of tricopper(I) complex **1** leads to the first example of synthetic copper complex which can simultaneously stabilize a mononuclear superoxo and dinuclear peroxo species (*vide infra*).

Decomposition of $[(\text{CTV-TMPA})\text{Cu}_3^{\text{I}}]^{3+}(\text{1})/\text{O}_2$ derived adducts

Previous systematic studies with tetradentate ligand-copper(I)/ O_2 systems show both superoxo and peroxo species are favored enthalpically but strongly disfavored entropically, accounting for their instability at higher temperatures.⁵³ As shown (*vide supra*), oxygenation of **1** results in a mixture of binuclear peroxo (one or two species) and mononuclear superoxo complexes. While UV-vis and rR spectra could be obtained, in fact these species are moderately unstable and complete decomposition occurs within ~2 hr even at -80°C (Figure 14). We also note that both superoxo and peroxo species appear to decompose at the same rate and a purple ($\lambda_{\text{max}} = 685\text{ nm}$) colored solid could be obtained by workup at room temperature. Although it appeared to a homogeneous material, the exact nature of this species could not be determined (but see the Experimental Section).

$[(\text{CTV-TMPA})\text{Cu}_3^{\text{I}}]^{3+}(\text{1})$ Reaction with Sulfur

Recently we described the reversible reaction of $[(\text{L}^{\text{CH}_2\text{OMe}})\text{Cu}^{\text{I}}]^+$ (**2**) with elemental sulfur, the result being the formation of a single complex, the end-on bound disulfide-dicopper(II) complex $[(\text{L}^{\text{CH}_2\text{OMe}})\text{Cu}^{\text{II}}]_2(\mu\text{-}1,2\text{-S}_2^{2-})^{2+}$ (**2-S**) ($\lambda_{\text{max}} = 540\text{ nm}$, $\nu_{(\text{S-S})} = 492\text{ cm}^{-1}$; $\nu_{(\text{Cu-S})_{\text{sym}}} = 309\text{ cm}^{-1}$).⁵² This contrasts somewhat with the behavior of the parent complex where $[(\text{TMPA})\text{Cu}^{\text{I}}(\text{CH}_3\text{CN})]^+(\text{3})/\text{S}_8$ chemistry produces an equilibrium mixture of at least three complexes. Yet, an X-ray structure and rR spectroscopy elucidated corresponded to $[(\text{TMPA})\text{Cu}^{\text{II}}]_2(\mu\text{-}1,2\text{-S}_2^{2-})^{2+}$ (**3-S**).^{89,90} As noted, the only difference between TMPA and $\text{L}^{\text{CH}_2\text{OMe}}$ is the presence of the pyridyl $6\text{-CH}_2\text{-O-R}$ motif in the latter, and of course this is present for all three copper ion binding sites within CTV-TMPA. Thus we expected that $[(\text{CTV-TMPA})\text{Cu}_3^{\text{I}}]^{3+}(\text{1})/\text{S}_8$ chemistry might also generate characterizable end-on bound disulfide-dicopper(II) complexes similar to what occurs for **2-S**.

Addition of elemental sulfur (as S_8)⁹¹ to **1** at -80°C under argon causes an immediate change in color of the solution from yellow to purple; such solutions are very stable, but decreased absorptions occur rapidly even upon slight warming. As described above, $[(\text{CTV-TMPA})\text{Cu}_3^{\text{I}}]^{3+}(\text{1})$ reacts rapidly and irreversibly with O_2 at room temperature, however it is required that lower temperatures be used in order to characterize and study Cu_3/O_2 adducts.⁵ Titrations of S_8 with $[(\text{CTV-TMPA})\text{Cu}_3^{\text{I}}]^{3+}(\text{1})$ show that (based on UV-vis criteria, and resonance Raman spectroscopy, see below) only one species forms in solution, $[(\text{CTV-TMPA})\text{Cu}_3^{\text{II}}]_2(\mu\text{-}1,2\text{-S}_2^{2-})_3^{6+}$ (**1-S**), $\lambda_{\text{max}} = 544\text{ nm}$ (Figure 15). The proposed structure for **1-S** is shown in the diagram here and discussed further below.

Temperature dependant sulfur binding in [(CTV-TMPA)Cu^I₃]³⁺ (1)

We find that by varying the temperature of solutions of [(CTV-TMPA)Cu^I₃]³⁺(1)/S₈, we can nicely vary the extent of sulfur binding. Heating/cooling cycles on solutions of [(CTV-TMPA)Cu^I₃]³⁺(1)/S₈ could be easily carried out (Figure 16). That elemental sulfur is released from a warmed (to RT) –80 °C solution is evident from the fact that no sulfur from an outside source is needed to fully reform the purple solution upon re-cooling. The cycle can be repeated multiple times without addition of more S₈ or an absorbance intensity change. We note that the dioxygen reactivity of [(CTV-TMPA)Cu^I₃]³⁺ (1) is not similarly reversible. We also note that a similar temperature variation of [{(L^{CH₂OMe})Cu^{II}}₂(μ-1,2-S₂²⁻)]²⁺ (2-S) resulted in complete loss of sulfur at room temperature.⁵² However, in the present case, 1-S did retain some sulfur coordination even at room temperature. Thus, attempts to isolate solid 1-S (with decreased intensity, hence a mixture of 1 and 1-S) were carried out successfully and (re)dissolving the solid material showed a similar intensity to that of a warmed-up solution resulting from 1/sulfur reactivity.

[(CTV-TMPA)Cu^I₃]³⁺(1)/S₈ Product Resonance Raman Spectroscopy

Resonance Raman spectra of [(CTV-TMPA)Cu^I₃]³⁺ (1)/S₈ with ³²S and ³⁴S isotopic substitution were collected with λ_{ex} = 530 nm at 77 K. The peaks at 494 cm⁻¹ and 307 cm⁻¹ shift to 482 cm⁻¹ and 302 cm⁻¹, respectively, with ³⁴S isotopic substitution (Figure 17). Based on spectral similarities with [{(L^{CH₂OMe})Cu^{II}}₂(μ-1,2-S₂²⁻)]²⁺ (2-S),⁹³ we can assign the complex as having a single type of end-on μ-1,2 disulfido-dicopper(II) complex, [(CTV-TMPA)Cu^{II}₃]₂(μ-1,2-S₂²⁻)₃⁶⁺ (1-S). The 307 cm⁻¹ peak can be assigned as a ν(Cu-S)_{sym} vibration. The peaks at 494 cm⁻¹ and 476 cm⁻¹ in the ³²S Raman spectrum are the result of a Fermi resonance of the ν(S-S) peak with a local non-enhanced mode of the same symmetry and correspond to a pre-interaction ν(S-S) of 489 cm⁻¹.⁵²

The high ν(S-S) value of 489 cm⁻¹ for [(CTV-TMPA)Cu^{II}₃]₂(μ-1,2-S₂²⁻)₃⁶⁺ (1-S³²S) which was also observed previously for [{(TMPA)Cu^{II}}₂(μ-1,2-S₂²⁻)]²⁺ (3-S) and [{(L^{CH₂OMe})Cu^{II}}₂(μ-1,2-S₂²⁻)]²⁺ (2-S), is indicative of the strong interaction between the S₂²⁻ and Cu^{II} ion centers, which removes electron density from the S-S π*_σ orbital and leads to a strengthened S-S bond. The 259 cm⁻¹ is assigned as a ν(Cu-N)_{py} stretch⁹² and the two additional Raman peaks, observed at 415 and 452 cm⁻¹, are the result of a Fermi splitting of ν(Cu-N)_{amine} and correspond to a pre-interaction ν(Cu-N)_{amine} of 449 cm⁻¹.

Complex Formulation; Proposed Structure

We suggest that the product of the [(CTV-TMPA) Cu^I₃]³⁺(1)/S₈ reaction is a hexanuclear complex [(CTV-TMPA)Cu^{II}₃]₂(μ-1,2-S₂²⁻)₃⁶⁺ (1-S) (see diagram above) (λ_{max} = 544 nm, Figure 15). This proposal derives from the very close similarity in characteristic UV-vis and rR spectroscopic features of 1-S as compared to the well characterized and comparable complexes employing the L^{CH₂OMe} and TMPA ligands. Complex 1-S possesses only the [(ligand)Cu^{II}]₂(μ-1,2-S₂²⁻)²⁺ moiety, a binuclear structure. Thus, for our CTV-TMPA tri-copper framework, the simplest formulation and complex likely to fulfill this requirement of possessing equivalent binuclear units would be the hexanuclear complex depicted in the diagram above. Determination of the exact structure would have to await an X-ray structure determination.

Summary/Conclusions

Our continuing efforts into modeling the active sites of copper-cluster containing enzymes have led us to synthesize CTV-based trinucleating ligand, CTV-TMPA, which employs tetradentate chelates as their copper binding sites. A series of chemical studies with the copper(I)-complex of CTV-TMPA, [(CTV-TMPA)Cu^I₃]³⁺, were carried out, including

studied on cyclic voltammetry, CO-binding and release, PPh₃-binding, complete oxidation of copper centers to a tricopper(II) complex and oxidative dehalogenation chemistry. The Cu^I/O₂ chemistry leads to the first example of a synthetic copper complex which can stabilize a mononuclear superoxo and dinuclear peroxo species simultaneously within one system. The resonance Raman studies of CTV-TMPA-Cu₃^I/O₂ adduct(s) at lower concentration (1.0 mM) demonstrate two types of end-on peroxo and one type of superoxo species, whereas at a bit higher concentration (1.53 mM), formation of only one type of peroxo along with a superoxo species are observed. The reaction of elemental sulfur with [(CTV-TMPA) Cu₃^I]³⁺ (**1**) leads to formation of a single dicopper-disulfide moiety with characteristic UV-vis and rR spectroscopic signatures, proposed to lead to an overall hexanuclear structure.

Our newly designed CTV based trinuclear ligand and tricopper(I) chemistry at this point appears not to lead to O₂-chemistry which might bear on, for example, the cooperative tricopper/O₂ reactivity observed in MCOs. Still, we have learned and we think advanced our knowledge and thinking about copper ion cluster design and chemistry.⁹³ The seemingly daunting task to design tri- or tetranuclear copper complexes which might mediate efficient dioxygen four-electron reduction (of interest in MCO (bio)chemistry or in fuel cell catalysts design) remains of great significance and interest.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Solomon E, Sarangi R, Woertink J, Augustine A, Yoon J, Ghosh S. *Acc. Chem. Res.* 2007;40:581–591. [PubMed: 17472331]
2. Holm, RH.; Solomon, EI., editors. *Chem. Rev.* Vol. 104. 2004. Biomimetic Inorganic Chemistry.
3. Fenton DE, Okawa H. *J. Chem. Soc. Dalton Trans* 1993:1349–1357.
4. Karlin KD. *Science* 1993;261:701–708. [PubMed: 7688141]
5. Mirica LM, Otsenwaelder X, Stack TDP. *Chem. Rev* 2004;104:1013–1045. [PubMed: 14871148]
6. Lewis EA, Tolman WB. *Chem. Rev* 2004;104:1047–1076. [PubMed: 14871149]
7. Lee, Y.; Karlin, KD. Highlights of Copper Protein Active-Site Structure/Reactivity and Synthetic Model Studies. In: Metzler-Nolte, N.; Kraatz, H-B., editors. *Concepts and Models in Bioinorganic Chemistry*. New York: Wiley-VCH; 2006. p. 363-395.
8. Op't Holt BT, Vance MA, Mirica LM, Heppner DE, Stack TDP, Solomon EI. *J. Am. Chem. Soc* 2009;131:6421–6438. [PubMed: 19368383]
9. Hatcher LQ, Karlin KD. *J. Biol. Inorg. Chem* 2004;9:669–683. [PubMed: 15311336]
10. Yoon J, Fujii S, Solomon EI. *Proc. Nat. Acad. Sci* 2009;106:6585–6590. [PubMed: 19346471]
11. Solomon EI, Augustine AJ, Yoon J. *Dalton Trans* 2008:3921–3932. [PubMed: 18648693]
12. Solomon EI, Chen P, Metz M, Lee S-K, Palmer AE. *Angew. Chem. Int. Ed* 2001;40:4570–4590.
13. Messerschmidt A. *Adv. Inorg. Chem* 1993;40:121–185.
14. Lee, D-H.; Lucchese, B.; Karlin, KD. Multimetal Oxidases. In: Que, L., Jr; Tolman, WB., editors. *Bio-coordination chemistry*. Vol. 8. Oxford: Elsevier Ltd; 2004. p. 437-457.
15. Bento I, Carrondo M, Lindley P. *J. Biol. Inorg. Chem* 2006;11:539–547. [PubMed: 16791638]
16. Kataoka K, Sugiyama R, Hirota S, Inoue M, Urata K, Minagawa Y, Seo D, Sakurai T. *J. Biol. Chem* 2009;284:14405–14413. [PubMed: 19297322]
17. Haltia T, Brown K, Tegoni M, Cambillau C, Saraste M, Mattila K, Djinnovic-Carugo K. *Biochem. J* 2003;369:77–88. [PubMed: 12356332]

18. Ghosh S, Gorelsky SI, DeBeerGeorge S, Chan JM, Cabrito I, Dooley DM, Moura JIG, Moura I, Solomon EI. *J. Am. Chem. Soc* 2007;129:3955–3965. [PubMed: 17352474]
19. Chan SI, Yu SS-F. *Acc. Chem. Res* 2008;41:969–979. [PubMed: 18605740]
20. Himes RA, Karlin KD. *Curr. Opin. Chem. Biol* 2009;13:119–131. [PubMed: 19286415]
21. Szeverényi Z, Knopp U, Zuberbühler AD. *Helv. Chim. Acta* 1982;65:2529–2539.
22. Hubberstey P, Russell CE. *J. Chem. Soc. Chem. Commun* 1995:959–960.
23. Chaudhuri P, Karpenstein I, Winter M, Butzlaff C, Bill E, Trautwein AX, Flörke U, Haupt H-J. *J. Chem. Soc. Chem. Commun* 1992:321–322.
24. Root DE, Henson MJ, Machonkin T, Mukherjee P, Stack TDP, Solomon EI. *J. Am. Chem. Soc* 1998;120:4982–4990.
25. Bonnefous C, Bellec N, Thummel RP. *Chem. Commun* 1999:1243–1244.
26. Kodera M, Tachi Y, Kita T, Kobushi H, Sumi Y, Kano K, Shiro M, Koikawa M, Tokii T, Ohba M, Okawa H. *Inorg. Chem* 2000;39:226–234. [PubMed: 11272529]
27. Liu XM, de Miranda MP, McInnes EJJ, Kilner CA, Halcrow MA. *Dalton Trans* 2004:59–64. [PubMed: 15356742]
28. Mirica LM, Stack TDP. *Inorg. Chem* 2005;44:2131–2133. [PubMed: 15792444]
29. Chen J, Wang X, Shao Y, Zhu J, Zhu Y, Li Y, Xu Q, Guo Z. *Inorg. Chem* 2007;46:3306–3312. [PubMed: 17355128]
30. DiNicola C, Karabach YY, Kirillov AM, Monari M, Pandolfo L, Pettinari C, Pombeiro AJL. *Inorg. Chem* 2007;46:221–230. [PubMed: 17198431]
31. Manzur J, Mora H, Vega A, Spodine E, Venegas-Yazigi D, Garland MT, ElFallah MS, Escuer A. *Inorg. Chem* 2007;46:6924–6932. [PubMed: 17655221]
32. Song YF, vanAlbada GA, Tang J, Mutikainen I, Turpeinen U, Massera C, Roubeau O, Costa JS, Gamez P, Reedijk J. *Inorg. Chem* 2007;46:4944–4950. [PubMed: 17497851]
33. Groysman S, Holm RH. *Inorg. Chem* 2009;48:621–627. [PubMed: 19138143]
34. Di Nicola C, Garau F, Karabach Yauhen Y, M. D. R. S. Martins Luísa, Monari M, Pandolfo L, Pettinari C, Pombeiro AJL. *Eur. J. Inorg. Chem* 2009;2009:666–676.
35. Karlin KD, Gan Q-F, Farooq A, Liu S, Zubietta J. *Inorg. Chim. Acta* 1989;165:37–39.
36. Karlin KD, Gan Q-F, Farooq A, Liu S, Zubietta J. *Inorg. Chem* 1990;29:2549–2551.
37. Karlin KD, Gan Q-f, Tyeklár Z. *Chem. Commun* 1999:2295–2296.
38. Frey ST, Sun HHJ, Murthy NN, Karlin KD. *Inorg. Chim. Acta* 1996;242:329–338.
39. Walsdorff C, Park S, Kim J, Heo J, Park K-M, Oh J, Kim K. *J. Chem. Soc. Dalton Trans* 1999:923–929.
40. Ohi H, Tachi Y, Itoh S. *Inorg. Chem* 2006;45:10825–10835. [PubMed: 17173442]
41. Ohi H, Tachi Y, Itoh S. *Inorg. Chem* 2004;43:4561–4563. [PubMed: 15257581]
42. Brown EC, Johnson B, Palavicini S, Kucera BE, Casella L, Tolman WB. *Dalton Trans* 2007:3035–3042. [PubMed: 17622421]
43. Battaini G, Granata A, Monzani E, Gullotti M, Casella L. *Adv. Inorg. Chem* 2006:185–233.
44. Adams H, Bailey NA, Dwyer JS, Fenton DE, Hellier PC, Hempstead PD, Latour JM. *J. Chem. Soc. Dalton Trans* 1993:1207–1216.
45. Chen PPY, Yang RBG, Lee JCM, Chan SI. *Proc. Nat. Acad. Sci* 2007;104:14570–14575. [PubMed: 17804786]
46. Ghiladi, RA. Thesis. Johns Hopkins University; 2003.
47. Collet A. *Tetrahedron* 1987;43:5725–5759.
48. Zhang S, Echegoyen L. *J. Am. Chem. Soc* 2005;127:2006–2011. [PubMed: 15701037]
49. Ameijde, Jv; Liskamp, RMJ. *Org. Biomolec. Chem* 2003;1:2661–2669.
50. Bohle DS, Stasko DJ. *Inorg. Chem* 2000;39:5768–5770. [PubMed: 11151378]
51. vanStrijdonck GPF, vanHaare JAEH, Honen PJM, vandenSchoor RCGM, Feiters MC, vanderLinden JGM, Steggerda JJ, Nolte RJM. *J. Chem. Soc. Dalton Trans* 1997:449–461.
52. Maiti D, Woertink JS, Vance MA, Milligan AE, Narducci Sarjeant AA, Solomon EI, Karlin KD. *J. Am. Chem. Soc* 2007;129:8882–8892. [PubMed: 17592845]

53. Zhang CX, Kaderli S, Costas M, Kim E-i, Neuhold Y-M, Karlin KD, Zuberbühler AD. *Inorg. Chem* 2003;42:1807–1824. [PubMed: 12639113]
54. Karlin KD, Kaderli S, Zuberbühler AD. *Acc. Chem. Res* 1997;30:139–147.
55. Collet A, Gabard J. *J. Org. Chem* 1980;45:5400–5401.
56. Canceill J, Collet A, Gottarelli G. *J. Am. Chem. Soc* 1984;106:5997–6003.
57. Liang H-C, Kim E, Incarvito CD, Rheingold AL, Karlin KD. *Inorg. Chem* 2002;41:2209–2212. [PubMed: 11952376]
58. van Strijdonck GPF, van Haare JAEH, Honen PJM, van den Schoor RCGM, Feiters MC, van der Linden JGM, Steggerda JJ, Nolte RJM. *J. Chem. Soc. Dalton Trans* 1997:449–461.
59. Vériot G, Dutasta J-P, Matouzenko G, Collet A. *Tetrahedron* 1995;51:389–400.
60. Matouzenko G, Veriot G, Dutasta JP, Collet A, Jordanov J, Varret F, Perrin M, Lecocq S. *New Journal of Chemistry* 1995;19:881–885.
61. Ghiladi RA, Ju TD, Lee D-H, Moënné-Loccoz P, Kaderli S, Neuhold Y-M, Zuberbühler AD, Woods AS, Cotter RJ, Karlin KD. *J. Am. Chem. Soc* 1999;121:9885–9886.
62. Ghiladi RA, Karlin KD. *Inorg. Chem* 2002;41:2400–2407. [PubMed: 11978105]
63. Lee D-H, Wei N, Murthy NN, Tyeklár Z, Karlin KD, Kaderli S, Jung B, Zuberbühler AD. *J. Am. Chem. Soc* 1995;117:12498–12513.
64. Karlin KD, Tyeklár Z, Farooq A, Haka MS, Ghosh P, Cruse RW, Gultneh Y, Hayes JC, Toscano PJ, Zubieta J. *Inorg. Chem* 1992;31:1436–1451.
65. Oshio H. *J. Chem. Soc. Dalton Trans* 1990:2985–2989.
66. Mazurek W, Bond AM, O'Connor MJ, Wedd AG. *Inorg. Chem* 1986;25:906–915.
67. Richardson DE, Taube H. *Inorg. Chem* 1981;20:1278.
68. Bohle DS, Stasko D. *Chem. Comm* 1998:567–568.
69. Rorabacher DB. *Chem. Rev* 2004;104:651–697. [PubMed: 14871138]
70. Hathaway, BJ. Copper. In: Wilkinson, G., editor. *Comprehensive Coordination Chemistry*. Vol. Vol. 5. New York: Pergamon; 1987. p. 533-774.
71. Hatcher LQ, Karlin KD. *Adv. Inorg. Chem* 2006;58:131–184.
72. Lucchese B, Humphreys KJ, Lee D-H, Incarvito CD, Sommer RD, Rheingold AL, Karlin KD. *Inorg. Chem* 2004;43:5987–5998. [PubMed: 15360248]
73. Nagao H, Komeda N, Mukaida M, Suzuki M, Tanaka K. *Inorg. Chem* 1996;35:6809–6815. [PubMed: 11666847]
74. Wei N, Murthy NN, Karlin KD. *Inorg. Chem* 1994;33:6093–6100.
75. Paul PP, Tyeklár Z, Jacobson RR, Karlin KD. *J. Am. Chem. Soc* 1991;113:5322–5332.
76. See Supporting Information.
77. Thompson DW, Kretzer RM, Lebeau EL, Scaltrito DV, Ghiladi RA, Lam K-C, Rheingold AL, Karlin KD, Meyer G. *J. Inorg. Chem* 2003;42:5211–5218.
78. Fry HC, Lucas HR, Narducci Sarjeant AA, Karlin KD, Meyer GJ. *Inorg. Chem* 2008;47:241–256. [PubMed: 18052158]
79. Kretzer RM, Ghiladi RA, Lebeau EL, Liang H-C, Karlin KD. *Inorg. Chem* 2003;42:3016–3025. [PubMed: 12716196]
80. Maiti D, Narducci Sarjeant AA, Itoh S, Karlin KD. *J. Am. Chem. Soc* 2008;130:5644–5645. [PubMed: 18393421]
81. Maiti D, Woertink JS, Narducci Sarjeant AA, Solomon EI, Karlin KD. *Inorg. Chem* 2008;47:3787–3800. [PubMed: 18396862]
82. Maiti D, Fry HC, Woertink JS, Vance MA, Solomon EI, Karlin KD. *J. Am. Chem. Soc* 2007;129:264–265. [PubMed: 17212392]
83. Itoh S. *Curr. Opin. Chem. Biol* 2006;10:115–122. [PubMed: 16504568]
84. The unequal intensities of the two Cu-O stretches shown in Figure 13 is the result of two factors. First, this intensity difference (also reflected in the intensity differences in the $\nu(\text{O-O})$'s in Figure 12) reflects minor differences in the concentration of $^{16}\text{O}_2$ vs. $^{18}\text{O}_2$ **1-O₂** samples. The concentration of **1-O₂** is slightly lower for the $^{18}\text{O}_2$ -generated sample presented than for the $^{16}\text{O}_2$ -generated sample. The second factor, which introduces error in intensity ratios (particularly for

spectra with lower signal-to-noise ratio as for Figures 12 and 13) is the method used to scale the spectra. Each spectrum presented is scaled to a nearby THF solvent mode, which is used as an internal standard. The spectrum shown in Figure 13 was scaled to the weak 586 cm^{-1} THF vibration while the O-O stretches shown in Figure 12 were scaled to the intense 880 cm^{-1} THF vibration. Thus error is introduced into the peak intensities as a result of the scaling procedure used in processing the rR data. The error in the calculated intensity of the weak 586 cm^{-1} THF vibration results in error in the final intensity of the scaled Cu-O stretches in Figure 13.

85. Pidcock E, Obias HV, Abe M, Liang H-C, Karlin KD, Solomon EI. *J. Am. Chem. Soc* 1999;121:1299–1308.
86. Baldwin MJ, Ross PK, Pate JE, Tyeklár Z, Karlin KD, Solomon EI. *J. Am. Chem. Soc* 1991;113:8671–8679.
87. Jacobson RR, Tyeklár Z, Karlin KD, Liu S, Zubieta J. *J. Am. Chem. Soc* 1988;110:3690–3692.
88. Schatz M, Raab V, Foxon SP, Brehm G, Schneider S, Reiher M, Holthausen MC, Sundermeyer J, Schindler S. *Angew. Chem. Int. Ed* 2004;43:4360–4363.
89. Chen P, Fujisawa K, Helton ME, Karlin KD, Solomon EI. *J. Am Chem. Soc* 2003;125:6394–6408. [PubMed: 12785779]
90. Helton ME, Chen P, Paul PP, Tyeklár Z, Sommer RD, Zhakarov LN, Rheingold AL, Solomon EI, Karlin KD. *J. Am. Chem. Soc* 2003;125:1160–1161. [PubMed: 12553805]
91. Elemental sulfur can exist in solution in a number of forms other than S_8 , see page 499 in Cotton, FA.; Wilkinson, G. *Advanced Inorganic Chemistry*. Sixth ed.. New York: John Wiley; Here, we will consistently describe the chemistry and reaction chemistry in terms of use of S_8 .
92. The high concentration of **1-S** used for the rR data presented in Figure 17 (2.45 mM) resulted in an intense, well-resolved $\nu(\text{Cu-N})\text{py}$ peak which could be reproducibly observed over the solvent vibrations
93. Perhaps the $6\text{-CH}_2\text{-O}$ linker between the TMPA and CTV units is too flexible, and a more rigid connection via synthetically constraining of the three chelates moieties would facilitate tricopper/small molecule interactions

Figure 1.

Top: Active site structure of ascorbate oxidase showing the type 2/type 3 catalytic triad, and the type 1 'blue' copper electron-transfer center. Coordinates (1AOZ) were taken from the Protein Data Bank (Brookhaven) and displayed using the program Rasmol. **Bottom:** Active site structure of Cu₂ sites in N₂OR. Coordinates (1QN1) were taken from the Protein Data Bank (Brookhaven) and displayed using the program Rasmol.

Figure 2.
Cyclic voltammogram of $[(\text{CTV-TMPA})\text{CuI}_3]^{3+}$ (**1**) in DMF at room temperature.

Figure 4.

UV-vis spectra at $-80\text{ }^{\circ}\text{C}$ in THF, illustrating formation of the $[(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3](\text{B}(\text{C}_6\text{F}_5)_4)_3$ upon addition of $[\text{Fe}^{\text{III}}\text{Cp}_2]^+$ (up to 3 equiv, 0.5 equiv each time) to tricopper(I) complex **1**. The downward arrow indicates the changes occurring during the transformation of copper(I) to copper(II). The upward arrow in the inset indicates increase in $d-d$ absorption during addition of $[\text{Fe}^{\text{III}}\text{Cp}_2]^+$, thus formation of copper(II), ($\lambda_{\text{max}} = 590\text{ nm}$). No further spectral changes are observed after addition of 3 equiv $[\text{Fe}^{\text{III}}\text{Cp}_2]^+$.

Figure 5. EPR spectrum following reaction of 3 $[\text{Fe}^{\text{III}}\text{Cp}_2]^+$ with $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**) at 77 K in 2-methyltetrahydrofuran (MeTHF) solvent {EPR parameters, $g_{\parallel} = 2.25$, $g_{\perp} = 2.05$, $A_{\parallel} = 152$ G, $A_{\perp} = 30$ G.}.

Figure 6.

UV-vis absorption spectra of $[(F_8TPP)Fe^{II}(THF)_2]$ in THF (**red**) at $-90\text{ }^{\circ}C$ and the spectra generated after addition of CO to generate $[(F_8TPP)Fe^{II}(CO)(THF)]$ (**brown**). The **green** spectrum is generated from a separate (same final copper concentration as that for **red** spectrum) addition of $[(F_8TPP)Fe^{II}(THF)_2]$ to the solution of $3PPh_3$ plus $[(CTV-TMPA)CuI_3(CO)_3]^{3+}$ (**1-CO**). A ~ 2.9 equiv $[(F_8TPP)Fe^{II}(CO)(THF)]$ formation is indicated (after baseline correction).

Figure 7. UV-vis spectra at $-80\text{ }^{\circ}\text{C}$ in THF, illustrating formation of **1-O₂** (red) from [(CTV-TMPA)CuI₃]³⁺ (**1**) (green) plus dioxygen. See text for assignments of the 427 and 543 nm absorptions.

Figure 8. rR spectra ($\nu_{\text{O-O}}$ region) of $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**) (**black**), $1\text{-}^{16}\text{O}_2$ (**red**), and $1\text{-}^{18}\text{O}_2$ (**blue**) ($\lambda_{\text{excit}} = 530 \text{ nm}$ at 77 K) in THF (1.0 mM).

Figure 9. rR spectra ($\nu_{\text{Cu-O}}$ region) of $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}(\mathbf{1})$ (black), $\mathbf{1}\text{-}^{16}\text{O}_2$ (red), and $\mathbf{1}\text{-}^{18}\text{O}_2$ (blue) ($\lambda_{\text{excit}} = 530 \text{ nm}$ at 77 K) in THF (1.0 mM $\mathbf{1}$). See text for band assignments.

Figure 10.

rR spectra (ν_{O-O} region) of $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}(\mathbf{1})$ (black), $1\text{-}^{16}\text{O}_2$ (red), and $1\text{-}^{18}\text{O}_2$ (blue) ($\lambda_{\text{excit}} = 413 \text{ nm}$ at 77 K) in THF (1.0 mM $\mathbf{1}$).

Figure 11.
rR spectra ($\nu_{\text{Cu-O}}$ region) of [(CTV-TMPA)Cu^I₃]³⁺(**1**) (**black**), **1**-¹⁶O₂ (**red**), and **1**-¹⁸O₂ (**blue**) ($\lambda_{\text{excit}} = 413 \text{ nm}$ at 77 K) in THF (1.0 mM **1**)

Figure 12.

rR spectra (ν_{O-O} region) of $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**) (**black**), $1\text{-}^{16}\text{O}_2$ (**red**), and $1\text{-}^{18}\text{O}_2$ (**blue**) (at 77 K, $\lambda_{\text{excit}} = 530$ nm) in THF (1.53 mM **1**).

Figure 13. rR spectra ($\nu_{\text{Cu-O}}$ region) of [(CTV-TMPA)Cu^I₃]³⁺ (**1**) (**black**), **1-¹⁶O₂** (**red**), and **1-¹⁸O₂** (**blue**) (at 77 K, $\lambda_{\text{excit}} = 530$ nm) in THF (1.53 mM **1**).

Figure 14.

UV-vis spectra at $-80\text{ }^{\circ}\text{C}$ in THF, illustrating the decomposition (i.e., thermal transformation) of purple Cu-O₂ species (**1-O₂**). [(CTV-TMPA)Cu^I₃]³⁺ (**1**) (**green** spectrum) following bubbling with O₂ generates **1-O₂** (**red** spectrum) which then undergoes subsequent transformations (**brown**, **orange** and **black** spectra) leading finally to a final **1-Dec** species (**blue** spectrum). Also, see Experimental Section.

Figure 15.

UV-vis spectra at $-80\text{ }^{\circ}\text{C}$ in tetrahydrofuran (THF) solvent, illustrating that only the end-on disulfide complex $[\{(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3\}_2(\mu\text{-}1,2\text{-S}_2^{2-})_3]^{6+}$ (**1-S**) is generated from $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}(\text{1})/4\text{S}$. The yellow/orange spectrum is that for $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}(\text{1})$; addition of increasing amounts of sulfur results leads to the spectra shown in **black** (1S), **brownish-purple** (2S), **blue** (3S) and **green** (4S). No spectral changes were observed upon further addition of sulfur to the reaction solution.

Figure 16.

UV-vis spectra at $-80\text{ }^{\circ}\text{C}$ in THF, illustrating formation of the end-on disulfide complex $[\{(\text{CTV-TMPA})\text{Cu}^{\text{II}}_3\}_2(\mu\text{-}1,2\text{-S}_2^{2-})_3]^{6+}$ (**1-S**) from $[(\text{CTV-TMPA})\text{Cu}^{\text{I}}_3]^{3+}$ (**1**)/ S_8 reactivity (**red** spectrum). The **blue** spectrum shown resulted upon warming a solution of **1-S** (**red** spectrum). Re-cooling to $-80\text{ }^{\circ}\text{C}$ again leads to the reformation of the **red** spectrum. This cycle can be repeated several times (see text).

Figure 17.

Solvent subtracted resonance Raman spectra of **1-S** with naturally abundant sulfur (red) and with isotopically enriched ^{34}S (blue) in THF ($\lambda_{\text{excit}} = 530 \text{ nm}$, 77 K). See text for assignments and discussion.

Scheme 1.

The successful generation of CTV-TMPA was confirmed by ^1H -NMR spectroscopic analysis, FAB and ESI-MS spectrometry along with elemental analysis.

Scheme 2.

Scheme 3.

Scheme 4.

Scheme 5.

Chart 1.

Table 1

Cyclic voltammetry data for copper(I) complexes in DMF.

Complexes	$E_{1/2}$ (mV) vs Fc/Fc^+	ΔE_p (mV)	i_{pa}/i_{pc}
$[(TMPA)Cu^I(CH_3CN)]^+ (3)$	-610	80	0.80
$[(L^{CH_2OMe})Cu^I]^+ (2)$	-500	115	0.87
$[(CTV-TMPA)Cu^I_3]^{3+} (1)$	-490	145	0.84

Table 2
Spectroscopic features for copper-dioxygen adducts related to those observed in 1/O₂ solutions.

μ -1,2-Peroxodicopper(II)-Complexes	Solvent	UV-vis/nm (ϵ -M ⁻¹ cm ⁻¹)	ν , cm ⁻¹ , ¹⁶ O- ¹⁸ O (¹⁸ O- ¹⁸ O)	ν , cm ⁻¹ , Cu- ¹⁶ O (Cu- ¹⁸ O)	ref
[(CTV-TMPA)Cu ^{II} _m (μ -1,2-O ₂ ²⁻)] ⁿ⁺ (1-O₂) (1.0 mM)	THF	543 (9650), 605 (8100)	842 (795) 825 (778)	506 (480) 548 (522)	^a
[(CTV-TMPA)Cu ^{II} _m (μ -1,2-O ₂ ²⁻)] ⁿ⁺ (1-O₂) (1.53 mM)	THF		825 (778)	506 (480)	^a
[[(TMPA)Cu ^{II}] ₂ (μ -1,2-O ₂ ²⁻)] ²⁺ (3-O₂) (parent complex)	EtCN	525 (11300), 615 (5800)	832 (788)	561 (535)	86
	Et ₂ O	520 (19200)	827 (783)	561 (535)	87
[[(L ^{CH₂OMe})Cu ^{II}] ₂ (μ -1,2-O ₂ ²⁻)] ²⁺ (2-O₂)	THF	540 (9550), 610 (6500)	848(801)	550(524)	81
End-on bound Cu^{II}(O₂⁻) complex	Solvent	UV-vis/nm (ϵ -M ⁻¹ cm ⁻¹)	ν , cm ⁻¹ ¹⁶ O- ¹⁶ O (¹⁸ O- ¹⁸ O)	ν , cm ⁻¹ Cu- ¹⁶ O (Cu- ¹⁸ O)	ref
[(CTV-TMPA)Cu ^{II} _x (O ₂ ⁻)] ^{y+} (1-O₂) (1.0 mM)	THF	427 (3150)	1129 (1069)	1129 (1069)	^a
[(Me ₂ N-TMPA)Cu ^{II} (O ₂ ⁻)] ⁺	THF	418 (4300)	1121 (1058)	1121 (1058)	82
[(TMG ₃ tren)Cu ^{II} (O ₂ ⁻)] ⁺	Acetone	447 (3400)	1117 (1059)	^b	88

^aThis work.

^bNot reported.

Table 3

Summary of spectroscopic features for related copper-sulfur adducts.

μ -1,2-Disulfide-dicopper(II) - Complexes	Solvent	UV-vis/nm (ϵ -M ⁻¹ cm ⁻¹)	ν , cm ⁻¹ , ³² S, ³² S (³⁴ S, ³⁴ S)	ν , cm ⁻¹ , Cu- ³² S (Cu- ³⁴ S)	ref
[{(CTV-TMPA)Cu ^{II}] ₃ (μ -1,2-S ₂ ²⁻) ₃] ⁶⁺ (1-S)	THF	544 (7270)	489(479)	307(302)	^a
[{(TMPA)Cu ^{II}] ₂ (μ -1,2-S ₂ ²⁻) ²⁺ (3-S)	CH ₃ CN	568 ^b	499(490)	316(312)	90
[{(L ^{CH₂OMe})Cu ^{II}] ₂ (μ -1,2-S ₂ ²⁻) ²⁺ (2-S)	THF	540(4600) ^c	492(482)	309(304)	52

^aThis work.

^bTHF,

^c2-methyltetrahydrofuran.