Published in final edited form as:

J Surg Res. 2010 December ; 164(2): 214–215. doi:10.1016/j.jss.2010.02.015.

MESENCHYMAL STEM CELLS—A NEW APPROACH TO INTESTINAL ISCHEMIA/REPERFUSION INJURY?

Aaron M. Abarbanell, M.D.¹

¹ Department of Surgery, Indiana University School of Medicine, Indianapolis, Indiana

Keywords

stem cells; intestinal ischemia/reperfusion; paracrine signaling; inflammation

Mesenchymal stem cells (MSCs) possess many properties that may make them suitable as transplanted therapeutic agents for ischemia/reperfusion injury (I/R). These properties include but are not limited to paracrine signaling [1], immunomodulation of the recipient environment [2,3], and possibly differentiation into the injured tissue [4]. Several studies have investigated the effects of MSCs with respect to I/R of the heart [5], brain [6] and kidney [7]. However, the role of MSCs has not been determined in the setting of intestinal I/R. In this issue, Jiang and colleagues studied this question in a rat model of I/R [8]. Using an established model, the superior mesenteric artery (SMA) was occluded for 45 minutes, and at the start of reperfusion MSCs were injected into the submucosa of the small bowel intestine. Treatment of the small bowel with MSC decreased the severity of intestinal barrier injury, bacterial translocation and serum lactate levels in this study.

Given the preponderance of evidence supporting the potential therapeutic roles for MSCs with respect to other organ systems, the authors are to be commended for pursuing this line of investigation with respect to the small bowel. In light of the observed beneficial effects of MSC treatment on the bowel, there are many questions to be answered. For instance what is the mechanism by which MSCs exert their salutary effects, and how can we clinically deliver these MSCs? What would be the effects of a pre-ischemic strategy compared to a post-ischemic strategy? Finally, what is the appropriate number of cells to deliver?

Certainly some answers may be inferred from previous studies that indicate that MSCs produce potent trophic factors that are associated with increased angiogenesis and decreased apoptosis [6]. However, the current study did not examine cytokine levels in the bowel after I/R. It is likely given the improved gut integrity in the treatment group as well as the histologic data that apoptosis is decreased in this study. Nonetheless, this supposition and other possible associations need to be more rigorously tested in future experiments.

Bowel I/R is encountered in multiple clinical scenarios such as necrotizing enterocolitis, trauma, cardiac surgery, sepsis, abdominal aortic surgery, strangulated hernias and bowel transplantation. To date, numerous pathways have been linked to the injury related to bowel

Correspondence address: Aaron M. Abarbanell, M.D., 2017 Van Nuys Medical Science Building, 635 Barnhill Drive, Indianapolis, Indiana 46202, aabarban@iupui.edu, Phone: 317-278-8625, Fax: 317-274-6896.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

I/R [9]. Thus it is likely that a therapeutic strategy for bowel I/R will be multi-factorial. As the understanding of MSC biology continues to evolve, the role of these cells may indeed be promising for combating the devastating effects of intestinal I/R injury.

References

1. Caplan AI, Dennis JE. Mesenchymal stem cells as trophic mediators. *J Cell Biochem* 2006;98:1076. [PubMed: 16619257]
2. Atoui R, Asenjo JF, Duong M, et al. Marrow stromal cells as universal donor cells for myocardial regenerative therapy: their unique immune tolerance. *Ann Thorac Surg* 2008;85:571. [PubMed: 18222266]
3. Atoui R, Shum-Tim D, Chiu RC. Myocardial regenerative therapy: immunologic basis for the potential "universal donor cells". *Ann Thorac Surg* 2008;86:327. [PubMed: 18573459]
4. Psaltis PJ, Zannettino A, Worthley SG, et al. Mesenchymal Stromal Cells - Potential for Cardiovascular Repair. *Stem Cells*. 2008
5. Wang M, Tsai BM, Crisostomo PR, et al. Pretreatment with adult progenitor cells improves recovery and decreases native myocardial proinflammatory signaling after ischemia. *Shock* 2006;25:454. [PubMed: 16680009]
6. Dharmasaroja P. Bone marrow-derived mesenchymal stem cells for the treatment of ischemic stroke. *J Clin Neurosci* 2009;16:12. [PubMed: 19017556]
7. Togel F, Hu Z, Weiss K, et al. Administered mesenchymal stem cells protect against ischemic acute renal failure through differentiation-independent mechanisms. *Am J Physiol Renal Physiol* 2005;289:F31. [PubMed: 15713913]
8. Jiang H, Qu L, Li Y, et al. Bone Marrow Mesenchymal Stem Cells Reduce Intestinal Ischemia/Reperfusion Injuries in Rats. *J Surg Res*. 2009
9. Mallick IH, Yang W, Winslet MC, et al. Ischemia-reperfusion injury of the intestine and protective strategies against injury. *Dig Dis Sci* 2004;49:1359. [PubMed: 15481305]