Published in final edited form as:

Nat Protoc. 2010 ; 5(2): 294–302. doi:10.1038/nprot.2009.223.

Preparation of the Caspase-3/-7 substrate Ac-DEVD-pNA via Solution-Phase Peptide Synthesis

Quinn P. Peterson¹, David R. Goode², Diana C. West², Rachel C. Botham², and Paul J. Hergenrother^{1,2,*}

¹Department of Biochemistry, University of Illinois Urbana-Champaign, 600 S. Matthews, Urbana, IL 61801

²Department of Chemistry, University of Illinois Urbana-Champaign, 600 S. Matthews, Urbana, IL 61801

Abstract

The caspase enzymes are integral to cellular inflammation and apoptotic cascades, and are commonly studied by cell biologists, medicinal chemists, and chemical biologists. In particular, the assessment of caspase enzymatic activity is a standard method to evaluate cell death pathways and novel apoptosis-modulating agents. Caspase enzymatic activity can be conveniently monitored with peptidic chromogenic or fluorogenic substrates, with certain peptide sequences imparting selectivity for certain caspases. The synthesis of these peptide substrates is typically performed via solid-phase synthesis, a method that is not ideal for production of the gram quantities needed for high-throughput screening. Described herein is a facile method for the synthesis of the Ac-DEVD-pNA caspase-3 substrate using solution-phase peptide synthesis. This protocol, involving iterative PyBOP-mediated couplings and Fmoc deprotections, is rapid, operationally simple, and can be used to generate over one gram of product.

INTRODUCTION

Caspases are a group of cysteine proteases that play critical roles in the inflammatory response (caspases-1, -4, -5) and apoptotic cascade (caspases-2, -3, -6, -7, -8, -9, -10, -12).¹ Caspases reside in the cell as low activity zymogens called procaspases.² During apoptosis, a caspase activation cascade begins with a signal that activates an “initiator” procaspase to the active caspase enzyme, with procaspase-8 and -10 being activated through the dimerization of cellular death receptors,³ and procaspase-9 being activated through cytochrome c release from the mitochondria and formation of the apoptosome.^{4,5} Once activated, caspase-8 and caspase-9 catalyze the hydrolysis of procaspase-3 and procaspase-7, forming the “executioner” caspases-3 and -7.^{6–8} Caspase-3 (and to a lesser extent caspase-7) is responsible for the proteolytic cleavage of over 100 different protein substrates,^{9,10} resulting in the systematic dismantling of the cell and apoptosis (Figure 1).

Given their central role in apoptotic cell death, aberrant caspase activity can have disastrous consequences for an organism. For example, in many degenerative diseases an increase in

*To whom correspondence should be addressed: hergenro@uiuc.edu.

AUTHOR CONTRIBUTION STATEMENT

All authors contributed extensively to the work presented in this paper.

Competing Financial Interest

The authors declare that they have no competing financial interests.

caspase activity is observed, and this is believed to lead to the premature cell death. Evidence for elevated caspase activity has been seen in Alzheimer's disease,^{11–13} Parkinson's disease,^{12,14} Huntington's disease,¹⁵ ALS¹⁶ and several others. In these cases, caspase inhibitors have been long sought as potential cytoprotective agents, and indeed the first caspase inhibitors are now in clinical trials for the treatment of liver disease, myocardial infarction, and rheumatoid arthritis.¹⁷ Conversely, resistance to apoptosis is a hallmark of cancer,¹⁸ and cancer cells can evade apoptosis through a modulation in caspase activity. Caspase-8,^{19–24} and -10/25²⁶ are mutated in some cancers, and elevated levels of endogenous caspase inhibitor proteins (e.g., XIAP, survivin) are commonly observed in a variety of cancers.^{27–29} Interestingly, the levels of procaspase proteins are elevated in many tumor types,^{30–35} and this has led to the hypothesis that drugs that directly activate procaspase enzymes could selectively induce apoptotic death in cancer cells.^{35–37}

Shortly after the discovery of the caspases a series of elegant studies defined the preferred cleavage site for each caspase isozyme.^{38,39} As indicated by their name (cysteine *aspartic* acid *protease*), caspases recognize short peptidic sequences and universally cleave on the C-terminal side of aspartic acid residues. The substrate recognition sequence for each caspase has been largely elucidated; for the most part, these enzymes recognize tetrapeptide sequences. The discovery of the preferred recognition sequence for each caspase led directly to the development of peptidic inhibitors and substrates for each isozyme. In the development of inhibitors,^{39,40} an electrophilic “warhead” (typically an aldehyde or α -fluoro ketone) is appended to the carboxy terminus of the scissile aspartic acid. Likewise, substrates can be created by appending the appropriate chromophore or fluorophore to this same position. The use of the *para*-nitro aniline (pNA) chromophore is especially popular, and as such the pNA tetrapeptide substrates for caspases-1, -3, -6, -8, and -9 are available from many commercial vendors.

Caspase-3, with its multiple cellular substrates, is one of the most studied and important caspases. Procaspase-3 is activated to caspase-3 through the action of both caspase-8 and caspase-9, meaning that procaspase-3 is a lynchpin upon which both the death receptor and mitochondrial apoptotic pathways converge (Figure 1).³ As such, there is much interest in and usage of the preferred peptidic substrates for caspase-3, Ac-DEVD-pNA (colorimetric) and Ac-DEVD-AMC (fluorescent), in cell biology, chemical biology, and medicinal chemistry. For example, Ac-DEVD-pNA and Ac-DEVD-AMC are widely used to monitor caspase-3/-7 activity in cell culture, have been used for the *in vitro* identification of caspase-3/-7 inhibitors,^{40–50} and in the identification and assessment of small molecule activators of procaspase-3/-7.^{35–37} Caspase-3 or caspase-7 will catalyze the hydrolysis of Ac-DEVD-pNA, releasing the pNA chromophore which has a λ_{max} of 405 nm (Figure 2).

While the average cost of Ac-DEVD-pNA from commercial vendors (5 mg for ~\$60) is reasonable when small quantities of this reagent are needed, it is prohibitively expensive to purchase the gram quantities required to screen a small molecule library for caspase inhibition/activation. For example, to screen a 200,000 compound library for caspase-3 inhibition under standard 384-well plate assay conditions would require ~1.25 g of Ac-DEVD-pNA. Described herein is a protocol to synthesize 1 gram of this Ac-DEVD-pNA substrate. Through this method, Ac-DEVD-pNA can be produced for approximately 1/50th of the commercial cost using rapid, continuous solution-phase peptide synthesis.^{51,52} The solution-phase peptide synthesis method provides several advantages over solid-phase peptide synthesis for the generation of large quantities of short peptides. In solid-phase peptide synthesis, a large excess (as high as 5–40 equivalents) of amino acids are used to drive the coupling reactions to completion. In the solution-phase synthesis described herein, the amino acids are used in only slight excess (1.1 equivalents). In addition, peptides can easily be produced on the gram scale using solution-phase peptide synthesis. Ac-DEVD-

pNA synthesized through this method has been used to generate substrate for the assessment of caspase inhibitors,⁴⁰ for the high-throughput screening for procaspase-3 activators,³⁵ and for mechanistic studies on procaspase-3 activating compounds,^{36,37} and cytoprotective compounds.⁵³

Experimental Design

As shown in Figure 3, the Ac-DEVD-pNA peptide is synthesized from C- to N-terminus. The synthetic route begins by coupling *para*-nitroaniline to Fmoc-Asp(Ot-Bu)-OH using POCl₃.⁵⁴ After workup, the resulting Fmoc-Asp(Ot-Bu)-pNA amide (**1** in Figure 3) is purified by silica gel and isolated in 77% yield. This product is then carried through iterative Fmoc deprotection, amino acid coupling, and workup steps without any intermediate purification: protected valine is first coupled to provide **2**, then Fmoc deprotection and PyBOP coupling with protected glutamic acid provides **3**, and finally Fmoc deprotection and coupling with acylated aspartic acid provides the fully protected peptide **4**. This protected peptide is difficult to isolate via extraction and column chromatography. Attempted extraction of the resulting product into ethyl acetate, chloroform, methylene chloride and hexane results in a persistent emulsion and insoluble precipitate. Efforts to purify the crude material using column chromatography resulted in an insoluble material that precipitates on the column. However, this peptide can be isolated by trituration in diethyl ether. Protected peptide **4** is then globally deprotected using trifluoroacetic acid (TFA) to provide the final product. The product is isolated by trituration in diethyl ether to give an overall yield of 40% for the 3 couplings and the global deprotection.

This procedure has been performed on a 1 to 10 mM scale with minimal differences in protocol. When working on smaller scales, reaction times may be decreased to 20 minutes per coupling/deprotection step. Conversely, larger scales may require longer reaction times and should be monitored to verify completion. Other substrates or inhibitors can likely be synthesized using different chromophores or fluorophores on the carboxy terminus. Individuals wishing to perform such modified protocols should ensure that there is a free amine on the chromophore or fluorophore of choice, and the initial coupling reaction to the protected aspartic acid residue may require optimization. After coupling to aspartic acid, an analogous protocol would be followed for the remaining coupling/deprotection steps.

Standard solid-phase peptide syntheses involve coupling to a resin, followed by C- to N-terminal peptide construction in a similar iterative sequence involving deprotection, purification, and coupling. Purification of products is achieved by filtration, where impurities are washed through a filter which retains the solid support and the growing peptide chain. Because the peptide is forming on a solid support and not in solution, coupling reactions are not as efficient and require large excess of each amino acid to overcome this limitation. Furthermore, due to low loading efficiency on the solid support, a large amount of resin is often required produce gram quantities of the final peptide. The solution-phase synthesis described herein is advantageous when large quantities of a relatively short peptide must be synthesized. In this example, gram quantities of the Ac-DEVD-pNA caspase-3/-7 substrate can be synthesized, in good overall yield, using nearly stoichiometric quantities of reagents. This route also allows for the addition of the chromophore prior to couplings, enabling the purification of this product to be carried out early in the synthesis. In contrast, in solid-phase synthesis the protected peptide is typically cleaved from the resin before being coupled to the chromophore.

Although the synthesis reported herein is specific for the Ac-DEVD-pNA substrate, analogous strategies could likely be used to produce other caspase substrates, inhibitors, or reporters including, Ac-DEVD-AFC, Ac-DEVD-AMC, FAM-DEVD-fmk and others. In addition, the method outlined in Figure 3 represents an improvement over our previously

reported synthesis,⁴⁰ as the earlier protocol requires an extraction (after final coupling) that results in an emulsion that is difficult to resolve. The method reported herein takes advantage of the insolubility of the product in common solvents in utilization of precipitation-based purifications. We anticipate this method and those related to it will be useful for the preparation of large quantities of caspase substrates required for high-throughput screening.

MATERIALS

REAGENTS

Brine Solution, saturated NaCl solution (Fisher Scientific cat. no. S271)

Sodium Bicarbonate, saturated solution (Fisher Scientific cat. no. S233)

Sodium Sulfate, Anhydrous (Sigma-Aldrich cat. no. 239313)

Acetone (Sigma-Aldrich cat. no. 179124)

Methylene Chloride (Sigma-Aldrich cat. no. D65100) **CAUTION**

Methanol (Fisher Scientific cat. no. A412) **CAUTION**

Hexanes (Fisher Scientific cat. no. H303) **CAUTION** Highly flammable, avoid long-term exposure and environmental toxin.

Ethyl Acetate (Sigma-Aldrich cat. no. 319902)

Diethyl Ether (Fisher Scientific cat. no. E138)

N, N-diisopropylethylamine, DIEA, Hünig's Base (Sigma-Aldrich cat. no. D12, 580-6) **CAUTION**

Tris(2-aminoethyl)amine, TAEA (Sigma-Aldrich cat. no. 225630) **CAUTION**

Trifluoroacetic Acid (Sigma-Aldrich) **CAUTION**

Phosphate Buffer

p-nitroaniline (Aldrich) **CAUTION**

Ac-Asp(O*t*Bu)-OH (ChemImpex cat. no. 04343)

Fmoc-Asp(O*t*Bu)-OH (Novabiochem cat. no. 04-12-1013)

Fmoc-Glu(O*t*Bu)-OH (Novabiochem cat. no. 04-12-1020)

Fmoc-Val-OH (Novabiochem cat. no. 04-12-1039)

PyBOP (Novabiochem cat. no. 01-62-0016) **CAUTION**

Pyridine (EMD cat. no. PX2020-6) **CAUTION**

Ninhydrin TLC stain

DPIP TLC Stain

NaCl (Fisher Scientific cat. no. S271)

POCl₃ (Sigma-Aldrich cat. no. 201170) **CAUTION**

EQUIPMENT

Büchner funnel (CoorsTek cat. no. 60243)

One-necked 500 mL round-bottom flasks (Chemglass)

500 mL Erlenmeyer flasks (Pyrex)
250 mL Erlenmeyer flask (Pyrex)
Filter Paper (Whatman)
Glass Funnel (Pyrex)
150 mL fritted filter funnel with coarse frit (Chemglass)
Glass Stoppers (Pyrex)
50 mL Graduated Cylinder (Kimax)
High vacuum pump (Welch)
Laboratory clamps (Fisher Scientific)
Laboratory jacks (Fisher Scientific)
Magnetic stir/hot plate (Corning)
Rotary evaporator (Büchi)
1000 mL Separatory funnel (Pyrex or Kimax)
500 mL Separatory funnel (Pyrex or Kimax)
Small stir bars (Fisher Scientific)
4 vane stir bar (Fisher Scientific)
Rubber septa (Chemglass)
100 × 190 cm recrystallization dish (Kimax)
Thermometer (-20 to 20 degrees C)
50 mL disposable syringe (Henke Sass Wolf)
1 mL disposable syringe (Henke Sass Wolf)
5 mL disposable syringe (Henke Sass Wolf)
100 mL beaker (Pyrex)
Silica Gel (Silicycle)
TLC plates (Silicycle)

REAGENT SETUP

Phosphate buffer is 90 g of $\text{NaHPO}_4 \cdot \text{H}_2\text{O}$ (EMD cat. no. SX0710) and 37.5 g NaH_2PO_4 (Sigma-Aldrich cat. no. S9763) in 500 mL H_2O pH 5.5. Hünig's base was dried via distillation over CaH_2 immediately before use. Into a distillation flask was placed an appropriate volume of Hünig's base, CaH_2 was added, the flask was heated, and the distillate was collected. Alternatively, a sure-seal bottle of Hünig's base may be purchased and used.

EQUIPMENT SETUP

Thoroughly clean and dry overnight (12 h) all glassware in an oven (100 °C) prior to use.

PROCEDURE

Recrystallization of *p*-nitroaniline

1. Add 10 grams of *p*-nitroaniline (pNA) to a 500 mL Erlenmeyer flask. Add 450 mL of H₂O. On a stirring hot plate with a stir bar, boil to completely dissolve all solid material. If needed, add additional H₂O until all solid has been dissolved.
CAUTION: *p*-Nitroaniline is toxic. Use proper protective equipment when handling this compound.
2. Set up a hot filter by placing a funnel over an Erlenmeyer flask with ~10 mL of boiling H₂O. Place a fluted piece of filter paper into the funnel and wet with a small volume of H₂O (~5 mL). Allow the funnel and filter paper to heat in the steam from the Erlenmeyer flask.
3. Filter the hot pNA solution through the hot filter to remove any insoluble particulates.
4. Boil the filtrate until crystals begin to form. Immediately remove from heat and allow to cool completely in an undisturbed location. Long yellow needles will form.
5. Filter the crystals through a piece of filter paper to obtain pure crystals. Wash the crystals with two washes (20 mL) of ice cold H₂O. Allow crystals to air dry on the filter paper.

PAUSE POINT: The recrystallized *p*-nitroaniline can be stored at room temperature for at least 1 year.

Coupling of *p*-nitroaniline to Fmoc-Asp(O*t*-Bu)-OH

- 6 To a clean, dry, 500 mL round bottom flask add 4.115 g (10 mmoles) of Fmoc-Asp(O*t*-Bu)OH, 1.381g (10 mmoles) of pNA, and 30 mL of pyridine. Add a stir bar.
CAUTION: Pyridine is toxic. Avoid breathing the fumes. Work with pyridine in a chemical fume hood and cover reaction vessels with septum when moving between the hood and the rotary evaporator to minimize the spread of vapors.
- 7 Make a salt/ice bath by placing a large 4 vane stir bar in the bottom of a 100 × 190 cm recrystallization dish. Fill the dish with ice and pour 200 grams of NaCl over the ice. Place the dish on a stir plate in a chemical fume hood and stir until the bath reaches a temperature of –10 to –15°C as monitored with a thermometer.
- 8 Place the round bottom flask containing the Fmoc-Asp(O*t*-Bu)OH, pNA, and pyridine into the salt/ice bath with the neck clamped to a support in the hood and the flask submerged 1.5 inches in the ice/water/salt mixture.
- 9 Slowly add 1.02 mL (11 mmoles) of POCl₃ via a syringe and stir for 30 minutes while maintaining the salt/ice bath in the –10 to –15°C range.
CAUTION: POCl₃ is corrosive. Avoid contact with skin.
- 10 Quench the reaction by adding 100 mL of ice water. Fill a 100 mL beaker with ice. Add water to 100 mL. Slowly pour ice and water into the reaction vessel and allow to stir for 1 minute. The quenching reaction is exothermic and results in the release of HCl.

- 11 Pour the contents of the reaction vessel into a 500 mL separatory funnel. Add 150 mL of ethyl acetate (EtOAc). Extract the reaction mixture. The product will be extracted into the organic (upper) phase.
- CAUTION:** Pressure can build up in the separatory funnel during extraction. Vent the separatory funnel several times during shaking by turning the separatory funnel upside down and opening the stopcock to release built up air pressure.
- 12 Drain the aqueous (lower) phase into a 250 mL Erlenmeyer flask. Pour the organic phase into a separate 500 mL Erlenmeyer flask.
- 13 Pour the aqueous phase back into the separatory funnel and extract two more times as above with 150 mL of EtOAc combining the organic phases.
- 14 Pour the 3 collected organic phases (~450mL) into a 1 L separatory funnel. Add 150 mL of a saturated bicarbonate solution and shake vigorously, venting the funnel frequently. Allow the phases to separate (it may take up to 2 hours for this separation) and remove the aqueous phase.
- 15 Add 150 mL of brine to the separatory funnel and shake. Allow phases to separate and remove the aqueous phase.
- 16 Pour the organic phase into a 1 L Erlenmeyer flask and add 20–30 grams of solid anhydrous sodium sulfate to remove any residual water. The sodium sulfate should be free flowing in the bottom of the flask.
- 17 Pour the dried organic phase through a fluted filter paper in a funnel and collect the filtrate in a clean dry round bottom flask.
- 18 Concentrate the solution to <100 mL via rotary evaporation.
- 19 Attach a large 150 mL fritted filter funnel with a coarse frit to a clamp in a fume hood. Fill the filter 2" high with silica gel. Pour the concentrated product over the 2" silica pad and elute the product with a 3:1 mixture of hexane/EtOAc under mild vacuum. The product should come off in the first 300 mL and the impurities such as pyridine or the unreacted starting materials should stay on the column. The elution of product may be monitored via TLC. $R_f = 0.71$.
- 20 To remove residual pyridine, the reaction mixture should be co-evaporated in a series of solvents. Concentrate to near dryness using a rotary evaporator. Redissolve the product in 50 mL of toluene to create an azeotrope with the pyridine. Concentrate this solution to near dryness. Add 50 mL of EtOAc. Concentrate to dryness. Add 50 mL of MeOH. Concentrate to dryness. The product will be a yellow residue until the final evaporation at which point it will be a yellow to white powder. The theoretical yield for this step should be 5.316 grams of pure product.

PAUSE POINT: This is a good stopping point. The purified product can be stored in solid form, at room temperature, protected from light, for up to 6 months with no observable degradation. The following steps through the workup of Ac-DEVD-pNA should be completed during the course of one day.

Deprotection of Fmoc-Asp(Ot-Bu)-pNA

- 21 Dissolve 2.658 g (5 mmol) of the Fmoc-Asp(Ot-Bu)-pNA in 30 mL of methylene chloride in a 500 mL round bottom flask. Add a stir bar and clamp to a support in a chemical fume hood directly over a stir plate. Add 30 mL (200 mmol) of TAEA slowly to the reaction vessel. Allow to stir for 30 minutes.

- 22 Monitor the reaction by TLC to verify conversion. The TLC should be run in 10% MeOH in EtOAc. The TLC can be stained with Ninhydrin to visualize the free amines that are exposed during deprotection. Dip the TLC plate into the Ninhydrin stain and heat with a heat gun until color is exposed. $R_f = 0.43$.

Workup

- 23 After conversion has been verified, add 60 mL of EtOAc to the reaction and then wash the reaction mixture with 50 mL of brine in a separatory funnel.
- 24 Extract the reaction mixture three times with 50 mL of phosphate buffer. Discard the aqueous fractions and dry the organic layer over anhydrous sodium sulfate using 20–30 g as in step 16.
- 25 Concentrate to dryness with a rotary evaporator. The product should be a fluffy yellow to white powder. The theoretical yield should be 1.547 g.

PyBOP Coupling of Fmoc-Val-OH to Asp-pNA

- 26 Redissolve the product carried over from the previous step in 30 mL of methylene chloride in a 500 mL round bottom flask with stirring. Add 1.867 g of Fmoc-Val-OH (5.5 mmoles) to the reaction vessel.

CRITICAL STEP: Step 27 and 28 must be done as close in time as possible to prevent epimerization. Weigh out the PyBOP for step 28 before adding DIEA in step 27.

- 27 Add 1.3 mL of DIEA (7.5 mmoles) to the reaction vessel slowly via syringe.
- 28 Add 2.861 g (5.5 mmoles) of PyBOP and stir reaction for 30 minutes. The reaction should turn from cloudy to a clear yellow over the course of the 30 minute reaction.
- 29 Verify coupling by TLC. The TLC should be run in 10% MeOH in EtOAc. The TLC may be stained with Ninhydrin to verify the absence of the free amine indicating stoichiometric conversion. $R_f = 0.51$

Deprotection of Fmoc-Val-Asp-pNA

- 30 After verification of coupling, add 40 equivalents of TAEA (200 mmoles, 30 mL) via syringe to the reaction mixture to deprotect the N-terminal amino acid. Allow the reaction to stir for 30 minutes at room temperature. The reaction should turn cloudy over the course of the 30 minute reaction.
- 31 Verify deprotection by TLC. The deprotection can be detected by the presence of the free amine using the Ninhydrin stain. $R_f = 0.33$

Workup

- 32 After verifying the deprotection has come to completion, add 30 mL of EtOAc.
- 33 Transfer the reaction to separatory funnel and wash with 50 mL of brine. Allow phases to separate.
- 34 Extract the reaction three times with 50 mL of phosphate buffer. Discard the aqueous phases and dry the organic phase over anhydrous sodium sulfate using 20–30 g as in step 16 and 24.
- 35 Concentrate to dryness with a rotary evaporator. The product should be a yellow residue.

Coupling of Fmoc-Glu(Ot-Bu)-OH to Val-Asp-pNA

- 36 Add 30 mL of methylene chloride to the residue obtained above and stir to dissolve.
- 37 Add 2.341 g (5.5 mmol) of Fmoc-Glu(Ot-Bu)-OH to the reaction vessel.
CRITICAL STEP: Step 38 and 39 must be done as close in time as possible to prevent epimerization. Weigh out the PyBOP for step 39 before adding DIEA in step 38.
- 38 Add 1.3 mL of DIEA (7.5 mmol) to the reaction vessel via syringe.
- 39 Add 2.862 g of PyBOP (5.5 mmol) to the reaction vessel and stir for 30 minutes.
- 40 Verify coupling by TLC in 10% MeOH in EtOAc. Stain with ninhydrin to verify the absence of any free amine. $R_f = 0.45$

Deprotection of Fmoc-Glu-Val-Asp-pNA

- 41 After verification of coupling, add 40 equivalents of TAEA (200 mmol, 30 mL) via syringe to the reaction mixture to deprotect the N-terminal amino acid. Allow the reaction to stir for 30 minutes at room temperature.
- 42 Verify deprotection by TLC. The deprotection can be detected by the presence of the free amine using the Ninhydrin stain. $R_f = 0.42$

Workup

- 43 After verifying the deprotection has come to completion, add 30 mL of EtOAc.
- 44 Transfer the reaction to separatory funnel and wash with 50 mL of brine.
- 45 Extract the reaction three times with 50 mL of phosphate buffer. Discard the aqueous phases and dry the organic phase over anhydrous sodium sulfate using 20–30 g as in steps 16, 24 and 34.
- 46 Concentrate to dryness with a rotary evaporator. The product should be yellow residue.

Coupling of Ac-Asp-OH to Glu-Val-Asp-pNA

- 47 Add 30 mL of methylene chloride to the residue obtained above and stir to dissolve.
- 48 Add 1.272 g (5.5 mmol) of Ac-Asp(Ot-Bu)-OH to the reaction vessel.
CRITICAL STEP: Step 49 and 50 must be done as close in time as possible to prevent epimerization. Weigh out the PyBOP for step 50 before adding DIEA in step 49.
- 49 Add 1.3 mL of DIEA (7.5 mmol) to the reaction vessel via syringe.
- 50 Add 2.862 g (5.5 mmol) of PyBOP to the reaction vessel. Allow to stir at RT for 30 min. Take a TLC to verify coupling in 10% methanol in EtOAc. Stain the TLC plate with DPIP. $R_f = 0.65$

Workup of protected Ac-DEVD-pNA

- 51 Concentrate the reaction mixture to dryness using a rotary evaporator.

- 52 Add 60 mL of diethyl ether and triturate the solid product, washing the solid by swirling the flask. If necessary, pipette the ether over the solid several times to ensure good mixing.
- 53 Filter and collect the precipitate. The impurities such as HOBT and the phosphoamide byproducts of the PYBOP coupling will be soluble in the diethyl ether. Wash the precipitate with 15 mL of diethyl ether. Allow the precipitate to dry and transfer to a round bottom flask.
- 54 Allow to dry under vacuum overnight. The product should be a white/off-white powder.

PAUSE POINT: At this point the solid may be stored at RT for a week or at -20°C for at least 6 months.

Global Deprotection

- 55 Add 30 mL of TFA via syringe to the solid product and stir for 30 minutes to globally deprotect the fully protected product.
- 56 Remove the TFA by rotary evaporation.
- 57 Triturate the solid with 30 mL of diethyl ether, washing the solid by swirling the flask. If necessary, pipette the ether over the solid several times to ensure good mixing. Allow the precipitate to settle to the bottom and decant the diethyl ether. Repeat two times.
- 58 Allow the final product to dry under vacuum overnight. The product is a light tan solid.

CRITICAL STEP: The product can be assessed by ^1H NMR, mass spectrometry, and HPLC to verify proper synthesis and complete deprotection. The product appears as a single peak on an analytical reverse-phase HPLC (see trace in Supporting Information). The product should be stored at -20°C and is stable for at least one year.

TIMING

- Steps 1–5, recrystallization of pNA, 4 h
- Step 6–20, coupling *p*-nitroaniline to Fmoc-Asp(*O**t*-Bu)-OH, 6 h
- Step 21 and 22, deprotection of Fmoc-Asp(*O**t*-Bu)-pNA, 0.5 h
- Step 23–25, workup of Asp(*O**t*-Bu)-pNA, 1 h
- Step 26–29, PyBOP coupling of Fmoc-Val-OH to Asp(*O**t*-Bu)-pNA, 0.5 h
- Step 30 and 31, Deprotection of Fmoc-Val-Asp(*O**t*-Bu)-pNA, 0.5 h
- Step 32–35, Workup of Val-Asp(*O**t*-Bu)-pNA, 1 h
- Step 36–40, Coupling of Fmoc-Glu(*O**t*-Bu)-OH to Val-Asp(*O**t*-Bu)-pNA, 0.5 h
- Step 41 and 42, Deprotection of Fmoc-Glu(*O**t*-Bu)-Val-Asp(*O**t*-Bu)-pNA, 0.5 h
- Step 43–46, Workup of Glu(*O**t*-Bu)-Val-Asp(*O**t*-Bu)-pNA, 1 h
- Step 47–50, Coupling of Ac-Asp(*O**t*-Bu)-OH to Glu(*O**t*-Bu)-Val-Asp(*O**t*-Bu)-pNA, 0.5 h
- Step 51–54, Workup of protected Ac-DEVD-pNA, 15 h

Step 55–58, Global Deprotection of Ac-DEVD-pNA, 20 h

TROUBLESHOOTING

If the fully protected product is not pure after washing with diethyl ether, the product may be further washed with diethyl ether to remove further impurities. Alternatively, the product may be purified via reverse phase HPLC.

ANTICIPATED RESULTS

As described, the synthesis will produce 700–1000 mg of Ac-DEVD-pNA. The procedure can be scaled if desired.

Analytical data

Ac-DEVD-pNA:

m.p.: 179–184°C

HPLC (C18, See Supporting Information for trace)

¹H-NMR (500 MHz, CD₃OD) (see supporting Information)

HRMS(ESI) (*m/z*) calculated for C₂₆H₃₄N₆O₁₃, 639.22; found, 639.2254

Acknowledgments

We are grateful to the National Institutes of Health (R01-CA120439) for the support of this work. Q.P.P. was partially supported by a Chemistry–Biology Interface Training Grant from the National Institutes of Health (Ruth L. Kirschstein National Research Service Award 1 T32 GM070421 from the National Institute of General Medical Sciences) and by a predoctoral fellowship from the ACS Division of Medicinal Chemistry. D.C.W. was partially supported by Ruth L. Kirschstein National Research Service Award 3F31CA130138-01S1.

REFERENCES

1. Shi Y. Mol Cell 2002;9:459–470. [PubMed: 11931755]
2. Bose K, Pop C, Feeney B, Clark AC. Biochemistry 2003;42:12298–12310. [PubMed: 14567691]
3. Hengartner MO. Nature 2000;407:770–776. [PubMed: 11048727]
4. Acehan D, Jiang X, Morgan DG, Heuser JE, Wang X, Akey CW. Mol. Cell 2002;9:423–432. [PubMed: 11864614]
5. Chao Y, Shiozaki EN, Srinivasula SM, Rigotti DJ, Fairman R, Shi Y. PLoS Biol 2005;3:e183. [PubMed: 15941357]
6. Yin Q, Park HH, Chung JY, Lin SC, Lo YC, da Graca LS, Jiang X, Wu H. Mol. Cell 2006;22:259–268. [PubMed: 16630893]
7. Walsh JG, Cullen SP, Sheridan C, Luthi AU, Gerner C, Martin SJ. Proc. Natl. Acad. Sci. U S A 2008;105:12815–12819. [PubMed: 18723680]
8. Riedl SJ, Fuentes-Prior P, Renatus M, Kairies N, Krapp S, Huber R, Salvesen GS, Bode W. Proc Natl Acad Sci U S A 2001;98:14790–14795. [PubMed: 11752425]
9. Luthi AU, Martin SJ. Cell Death Differ 2007;14:641–650. [PubMed: 17273173]
10. Timmer JC, Salvesen GS. Cell Death Differ 2007;14:66–72. [PubMed: 17082814]
11. Engidawork E, Gulesserian T, Yoo BC, Cairns N, Lubec G. Biochem Biophys Res Commun 2001;281:84–93. [PubMed: 11178964]
12. Blandini F, Sinforiani E, Pacchetti C, Samuele A, Bazzini E, Zangaglia R, Nappi G, Martignoni E. Neurology 2006;66:529–534. [PubMed: 16505307]
13. Roth KA. J Neuropathol Exp Neurol 2001;60:829–838. [PubMed: 11556539]
14. Nagatsu T, Sawada M. J Neural Transm Suppl 2007;113–120. [PubMed: 17982884]
15. Rubinsztein DC, Carmichael J. Expert Rev Mol Med 2003;5:1–21. [PubMed: 14585171]

16. Martin LJ. *J Neuropathol Exp Neurol* 1999;58:459–471. [PubMed: 10331434]
17. Green DR, Kroemer G. *J Clin Invest* 2005;115:2610–2617. [PubMed: 16200193]
18. Hanahan D, Weinberg RA. *Cell* 2000;100:57–70. [PubMed: 10647931]
19. Soung YH, Lee JW, Kim SY, Jang J, Park YG, Park WS, Nam SW, Lee JY, Yoo NJ, Lee SH. *Cancer Res* 2005;65:815–821. [PubMed: 15705878]
20. Soung YH, Lee JW, Kim SY, Sung YJ, Park WS, Nam SW, Kim SH, Lee JY, Yoo NJ, Lee SH. *Oncogene* 2005;24:141–147. [PubMed: 15531912]
21. Kim HS, Lee JW, Soung YH, Park WS, Kim SY, Lee JH, Park JY, Cho YG, Kim CJ, Jeong SW, Nam SW, Kim SH, Lee JY, Yoo NJ, Lee SH. *Gastroenterology* 2003;125:708–715. [PubMed: 12949717]
22. Mandruzzato S, Brasseur F, Andry G, Boon T, van der Bruggen P. *J Exp Med* 1997;186:785–793. [PubMed: 9271594]
23. Teitz T, Wei T, Valentine MB, Vanin EF, Grenet J, Valentine VA, Behm FG, Look AT, Lahti JM, Kidd VJ. *Nat Med* 2000;6:529–535. [PubMed: 10802708]
24. Teitz T, Lahti JM, Kidd VJ. *J Mol Med* 2001;79:428–436. [PubMed: 11511973]
25. Shin MS, Kim HS, Kang CS, Park WS, Kim SY, Lee SN, Lee JH, Park JY, Jang JJ, Kim CW, Kim SH, Lee JY, Yoo NJ, Lee SH. *Blood* 2002;99:4094–4099. [PubMed: 12010812]
26. Park WS, Lee JH, Shin MS, Park JY, Kim HS, Lee JH, Kim YS, Lee SN, Xiao W, Park CH, Lee SH, Yoo NJ, Lee JY. *Oncogene* 2002;21:2919–2925. [PubMed: 11973654]
27. Kania J, Konturek SJ, Marlicz K, Hahn EG, Konturek PC. *Dig Dis Sci* 2003;48:266–271. [PubMed: 12643601]
28. Zou H, Yang R, Hao J, Wang J, Sun C, Fesik SW, Wu JC, Tomaselli KJ, Armstrong RC. *J Biol Chem* 2003;278:8091–8098. [PubMed: 12506111]
29. Shiozaki EN, Chai J, Rigotti DJ, Riedl SJ, Li P, Srinivasula SM, Alnemri ES, Fairman R, Shi Y. *Mol Cell* 2003;11:519–527. [PubMed: 12620238]
30. Roy S, Bayly CI, Gareau Y, Houtzager VM, Kargman S, Keen SLC, Rowland K, Seiden IM, Thornberry NA, Nicholoso DW. *Proc. Natl. Acad. Sci* 2001;98:6132–6137. [PubMed: 11353841]
31. Grigoriev MY, Pozharissky KM, Hanson KP, Imyanitov EN, Zhivotovsky B. *Cell Cycle* 2002;1:337–342. [PubMed: 12461296]
32. Krepela E, Prochazka J, Liul X, Fiala P, Kinkor Z. *Biol Chem* 2004;385:153–168. [PubMed: 15101558]
33. Fink D, Schlagbauer-Wadl H, Selzer E, Lucas T, Wolff K, Pehamberger H, Eichler HG, Jansen B. *Melanoma Res* 2001;11:385–393. [PubMed: 11479427]
34. Estrov Z, Thall PF, Talpaz M, Estey EH, Kantarjian HM, Andreeff M, Harris D, Van Q, Walterscheid M, Kornblau SM. *Blood* 1998;92:3090–3097. [PubMed: 9787143]
35. Putt KS, Chen GW, Pearson JM, Sandhorst JS, Hoagland MS, Kwon J-T, Hwang S-K, Jin H, Churchwell MI, Cho M-H, Doerge DR, Helferich WG, Hergenrother PJ. *Nature Chem Biol* 2006;2:543–550. [PubMed: 16936720]
36. Peterson QP, Goode DR, West DC, Ramsey KN, Lee J, Hergenrother PJ. *J Mol Biol* 2009;388:144–158. [PubMed: 19281821]
37. Peterson QP, Hsu DC, Goode DR, Novotny CJ, Totten RK, Hergenrother PJ. *J Med Chem.* 2009 10.1021/jm900722z.
38. Thornberry N, Rano TA, Peterson EP, Rasper DM, Timkey T, Garcia-Calvo M, Houtzager VM, Nordstrom PA, Roy S, Vaillancourt JP, Chapman KT, Nicholoso DW. *J Biol Chem* 1997;27:17907–17911. [PubMed: 9218414]
39. Nicholson DW, Ali A, Thornberry NA, Vaillancourt JP, Ding CK, Gallant M, Gareau Y, Griffin PR, Labelle M, Lazebnik YA, Munday NA, Raju SM, Smulson ME, Yamin T-T, Yu VL, Miller DK. *Nature* 1995;376:37–43. [PubMed: 7596430]
40. Goode DR, Sharma AK, Hergenrother PJ. *Org Lett* 2005;7:3529–3532. [PubMed: 16048334]
41. Smolewski P, Grabarek J, Phelps DJ, Darzynkiewicz Z. *Int. J. Oncol* 2001;19:657–663. [PubMed: 11562738]
42. Hardy JA, Wells JA. *J Biol Chem.* 2009

43. Chu W, Rothfuss J, Chu Y, Zhou D, Mach RH. *J Med Chem* 2009;52:2188–2191. [PubMed: 19326941]
44. Chu W, Rothfuss J, d'Avignon A, Zeng C, Zhou D, Hotchkiss RS, Mach RH. *J Med Chem* 2007;50:3751–3755. [PubMed: 17585855]
45. Chu W, Zhang J, Zeng C, Rothfuss J, Tu Z, Chu Y, Reichert DE, Welch MJ, Mach RH. *J Med Chem* 2005;48:7637–7647. [PubMed: 16302804]
46. Guo Z, Xian M, Zhang W, McGill A, Wang PG. *Bioorg Med Chem* 2001;9:99–106. [PubMed: 11197351]
47. Chen Y-H, Zhang Y-H, Zhang H-J, Liu D-Z, Gu M, Li J-Y, Wu F, Zhu X-Z, Li J, Nan F-J. *J Med Chem* 2006;49:1613–1623. [PubMed: 16509578]
48. Lee D, Long SA, Murray JH, Adams JL, Nuttall ME, Nadeau DP, Kikly K, Winkler JD, Sung C-M, Ryan MD, Levy MA, Keller PM, DeWolf WE. *J Med Chem* 2001;44:2015–2026. [PubMed: 11384246]
49. Wu JC, Fritz LC. *Methods* 1999;17:320–328. [PubMed: 10196103]
50. Chapman JG, Magee WP, Stukenbrok HA, Beckius GE, Milici AJ, Tracey WR. *Eur J Pharmacol* 2002;456:59–68. [PubMed: 12450570]
51. Hoeg-Jensen T, Jakobsen MH, Holm A. *Tetrahedron Lett* 1991;32:6387–6390.
52. Carpino LA, Ghassemi S, Ionescu D, Ismail M, Sadat-Aalae D, Truran GA, Mansour EME, Siwruk GA, Eynon JS, Morgan B. *Org Process Res Dev* 2002;7:28–37.
53. Thompson CM, Quinn CA, Hergenrother PJ. *J Med Chem* 2009;52:117–125. [PubMed: 19072121]
54. Rijkers DTS, Adams HPHM, Hemker HC, Tesser GI. *Tetrahedron* 1995;51:11235–11250.

Figure 1.

Pathways of caspase activation. Release of pro-apoptotic proteins (such as cytochrome c) from the mitochondria promotes the formation of the apoptosome, a multimeric complex including Apaf-1, cytochrome c, and procaspase-9. In the apoptosome, procaspase-9 becomes activated to caspase-9, an enzyme that can then proteolytically cleave procaspases-3 and -7, resulting in their activation. In an analogous fashion, procaspases-8 and -10 are activated to caspases-8 and -10 after death ligands bind to surface death receptors, initiating the dimerization and recruitment of procaspase-8 and -10. Once activated, caspases-8 and -10 can also proteolytically cleave procaspases-3 and -7 to form the active caspases-3 and -7.

Figure 2.
The hydrolysis of Ac-DEVD-pNA to Ac-DEVD and pNA, as catalyzed by caspase-3 or caspase-7.

Figure 3.
Solution-phase synthesis of Ac-DEVD-pNA.