Published in final edited form as:

Methods Mol Biol. 2009 ; 492: 321–351. doi:10.1007/978-1-59745-493-3_20.

Analysis of Protein Glycosylation and Phosphorylation Using Liquid Phase Separation, Protein Microarray Technology, and Mass Spectrometry

Jia Zhao, Tasneem H. Patwa, Manoj Pal, Weilian Qiu, and David M. Lubman

Summary

Protein glycosylation and phosphorylation are very common posttranslational modifications. The alteration of these modifications in cancer cells is closely related to the onset and progression of cancer and other disease states. In this protocol, strategies for monitoring the changes in protein glycosylation and phosphorylation in serum or tissue cells on a global scale and specifically characterizing these alterations are included. The technique is based on lectin affinity enrichment for glycoproteins, all liquid-phase two-dimensional fractionation, protein microarray, and mass spectrometry technology. Proteins are separated based on pI in the first dimension using chromatofocusing (CF) or liquid isoelectric focusing (IEF) followed by the second-dimension separation using nonporous silica RP-HPLC. Five lectins with different binding specificities to glycan structures are used for screening glycosylation patterns in human serum through a biotin–streptavidin system. Fluorescent phosphodyes and phosphospecific antibodies are employed to detect specific phosphorylated proteins in cell lines or human tissues. The purified proteins of interest are identified by peptide sequencing. Their modifications including glycosylation and phosphorylation could be further characterized by mass-spectrometry-based approaches. These strategies can be used in biological samples for large-scale glycoproteome/phosphoproteome screening as well as for individual protein modification analysis.

Keywords

Glycosylation; Phosphorylation; Posttranslational modification; Protein microarrays; Liquid phase separation; Lectin; Mass spectrometry

20.1. Introduction

Protein glycosylation and phosphorylation are very common post-translational modifications (PTM). The alteration in protein glycosylation on the cell surface and in body fluids has been shown to correlate with the progression of cancer and other disease states (1–4). These glycosylation changes in tumor-secreted proteins reflect fundamental changes in the enzymes involved in the glycosylation pathway. The ability to efficiently profile and characterize the variation in glycoproteins could lead to the identification of disease-associated glycan alterations and new diagnostic markers. Phosphorylation and dephosphorylation of proteins regulate a variety of cellular signaling pathways typically associated with major changes in cell function (5–7). It is, therefore, essential to detect the alteration in phosphorylation patterns on a global scale and to be able to identify the critical proteins involved in cell-cycle regulation where such proteins could be important in cancer onset and progression. Glycosylation and phosphorylation have been extensively studied by

several technologies such as mass spectrometry and microarrays. Protein microarrays are useful as a high throughput screening method for whole-cell lysates, fractionated proteomes, tissues, and antigen–antibody reactions, whereas mass spectrometry offers the ability for in-depth analysis of the target alterations (8,9).

The strategies included in this protocol employ an all-liquid phase enrichment and fractionation methodology coupled to protein microarray technology and mass spectrometry. The procedures are briefly outlined in Scheme 20.1. For serum glycoprotein profiling, glycoproteins are enriched by lectin affinity chromatography. After depositing on the microarray surface, the liquid-phase separated glycoproteins are probed with different lectins with a wide range of binding specificities to different glycan moieties. The glycoprotein–lectin interaction is assessed using a biotin–streptavidin system. Human serum can be screened for different glycan structures using this strategy thereby allowing one to do comparative studies that monitor individual glycosylation changes within a glycoproteome representing different biological states (10). The glycans of target glycoproteins are released by PNGase F and further mass-mapped or fragmented for structure analysis by mass spectrometry. For phosphoprotein analysis, an all-liquid two-dimensional separation method has been explored to map the phosphoprotein expression of a cell lysate to study changes in phosphorylation patterns. Proteins are separated based on pI in the first dimension using chromatofocusing (CF) or liquid isoelectric focusing (IEF) followed by a second-dimension separation based on hydrophobicity using nonporous silica RP-HPLC (11–14). Fluorescent phosphodies and phosphospecific antibodies are used to detect specific phosphorylated proteins (15). The specific phosphorylation site of the target-phosphorylated protein is detected by enzymatic dephosphorylation or MS2 neutral loss experiment. This method provides a convenient means of profiling of the phosphoproteome in human cancer cells and is useful to identify potential biomarkers in cancer.

20.2. Materials

20.2.1. Sample Preparation

1. Urea, Thiourea, *n*-octyl β -D-glucopyranoside (OG) (Sigma).
2. Tris[2-carboxyethyl] phosphine (TCEP) (Pierce).
3. Glycerol.
4. Protease inhibitor tablet (Roche, Indianapolis, IN).
5. Tissue homogenizer.
6. Centrifuge (refrigerated centrifuge capable of $30,000 \times g$).
7. Desalting column: PD-10 (Sephadex G-25; Amersham Pharmacia Biotech).
8. Lysis buffer: 7.5 M urea, 2.5 M thiourea, 4% *n*-octyl-D-glucopyranoside (OG), 10 mM Tris[2-carboxyethyl]phosphine (TCEP), 10% (v/v) glycerol, 50 mM Tris and 40 μ L protease inhibitor solution (10 mg tablet in 1 mL PBS buffer).

20.2.2. Top-Twelve Protein Depletion (For Serum Sample)

1. ProtomeLab IgY-12 proteome partitioning kit (Beckman Coulter, Fullerton, CA).
2. Spin filters, 0.45 μ m, (Millipore, Billerica, MA).
3. 10 $\times$ Dilution buffer: 100 mM Tris-HCl, pH 7.4, 1.5 M NaCl.
4. 10 $\times$ Stripping buffer: 1 M glycine-HCl, pH 2.5.
5. 10 $\times$ Neutralization buffer: 1 M Tris-HCl, pH 8.0.

6. 15-mL, 5-kDa MWCO Amicon filters (Millipore).

20.2.3. Lectin Affinity Glycoprotein Extraction

1. Agarose bound lectin, Wheat Germ Agglutinin, (WGA) and Agarose bound Concanavalin A (Con A) (Vector Laboratories, Burlingame, CA).
2. Empty spin columns (column capacity: 0.9 mL) (Pierce, Rockford, IL).
3. Lectin affinity binding buffer: 0.15 M NaCl, 20 mM Tris-HCl, 1 mM MnCl₂, 1 mM CaCl₂, pH 7.4.
4. Elution buffer: 0.3 M *N*-acetyl-glucosamine and 0.3 M Methyl- α -d-mannopyranoside in 20 mM Tris-HCl and 0.5 M NaCl, pH 7.0.

20.2.4. Separation of Proteins Based on pI

20.2.4.1. Micro-Rotofor Isoelectric Focusing

1. Cathode electrolyte: 0.1 M NaOH.
2. Anode electrolyte: 0.1 M H₃PO₄.
3. Rotofor running buffer: 3% CHAPS, 6 M urea, 2 M thiourea, 10 mM DTT, 2% ampholyte 3–10 (for pH 3–10), 2% ampholyte composed of 0.2% ampholyte 3–10, 0.9% ampholyte 3–5, and 0.9% ampholyte 4–6 (for pH 3–6).
4. Micro-Rotofor system (Bio-Rad).
5. Power supply with constant power capability.
6. 3-mL syringe and 10-mL syringes.
7. Sealing tape.
8. Forceps.
9. Cold trap.
10. Vacuum pump.

20.2.4.2. Chromatofocusing

1. HPLC-1D column (250 × 2.1 mm) (Beckman Coulter, Fullerton, CA).
2. Polybuffer CF start buffer: 25 mM bis-Tris propane (Sigma), 6 M urea and 0.1% *n*-OG, pH 8.5.
3. CF elution buffer: 7% v/v polybuffer 74 (Amersham Pharmacia, Piscataway, NJ) and 3% polybuffer 96 (v/v), 6 M urea, and 0.1% *n*-OG, pH 4.0.
4. 0.2- μ L nylon low extraction membrane filter (500 mL) (Corning Incorporated, Corning, NY).
5. 1 M NaCl solution.
6. Isopropanol (Sigma).
7. HPLC/FPLC system running at ambient temperature and at a flow rate of 0.2 mL/min, with fraction collector, UV detector (280 nm).
8. pH electrode/flow cell (Lazar Research Laboratories, Los Angeles, CA).

20.2.5. Nonporous-Reverse Phase-HPLC Separation

1. Nonporous-reverse phase (NPS-RP)-HPLC column: 33 × 4.6 mm Micra Platinum column, ODSII (recommended) (Eprogen).
2. Column heater.
3. Deionized water was purified using a Millipore RG system (Bedford, MA).
4. Acetonitrile (ACN).
5. Trifluoroacetic acid (TFA).

20.2.6. Protein Microarrays

20.2.6.1. Printing

1. Nanoplotter 2.0 (GeSiM).
2. Superamine and Superepoxy glass slides (TeleChem International), Nitrocellulose slides (Whatman, Inc.).
3. Printing buffer: 62.5 mM Tris-HCl (pH 6.8), 1% w/v sodium dodecyl sulfate (SDS), 5% w/v dithiothreitol (DTT), and 1% glycerol. Prepare immediately before reconstituting dried-down sample. Make fresh each time it is needed.
4. Speedvac centrifuge (Thermo, Waltham, MA).

20.2.6.2. Hybridization of Slides

Glycoprotein Microarrays with Multilectin Detection

1. Blocking buffer: 1% w/v BSA in PBS-T.
2. Primary detection solution: For Concanavalin A (ConA), Aleuria aurentia (AAL), Maackia amurensis (MAL), and Peanut agglutinin (PNA), 5 µg/mL biotinylated lectin solution in PBS-T (0.1% Tween-20) and for Sambuccus Nigra (SNA), 10 µg/mL lectin solution in PBS-T (0.1% Tween-20). All biotinylated lectins purchased from Vector Laboratories (Burlingame, CA).
3. Washing solution: PBS-T (0.1% Tween-20).
4. Secondary detection solution: 1:1,000 solution of 1 mg/mL Streptavidin conjugated to Alexafluor555 (Invitrogen) in 0.5% BSA and 1× PBS-T (0.1% Tween-20).

Phosphoprotein Analysis with Antibodies

1. Blocking buffer: 1% w/v bovine serum albumin (BSA) in PBS-T (phosphate buffered saline and 0.1% v/v Tween-20).
2. Probe buffer: 5 mM magnesium chloride, 0.5 mM DTT, 0.05% v/v TritonX-100, 5% glycerol, and 1% BSA in 1× PBS.
3. Primary antibody solution: 1:500 solution of 1 mg/mL goat polyclonal antiphosphoserine and goat polyclonal antiphosphothreonine (Zymed laboratories) and mouse monoclonal antiphosphotyrosine (Upstate signaling) in probe buffer.
4. Secondary antibody solution: 1:1,000 solution of 1 mg/mL donkey antigoat conjugated to Cy5 in probe buffer and 1:1,000 solution of 1 mg/mL donkey antimouse conjugated to Cy5 in probe buffer.

Phosphoprotein Analysis with ProQ Diamond

1. Blocking buffer: 1% BSA in PBS-T (0.1% Tween-20).

2. ProQ Diamond dye (Invitrogen).
3. ProQ Diamond destaining solution (Invitrogen).
4. 1× PBS.

20.2.6.3. Slide Scanning

1. Axon 4000A scanner (Molecular Devices, Sunnyvale, CA).

20.2.7. Protein Identification by Peptide Sequencing

1. SpeedVac concentrator (Thermo).
2. Ammonium bicarbonate (Sigma-Aldrich).
3. TPCK modified sequencing grade porcine trypsin (Promega, Madison, WI).
4. HPLC grade water.
5. Paradigm HPLC pump (Michrom Bio Resources, Inc., Auburn, CA).
6. Nanoreverse phase column (0.075 mm × 150 mm, C18).
7. Peptide nanotrap (0.15 × 50 mm, C18).
8. A 75-μm metal spray tip (Michrom).
9. Finnigan LTQ mass spectrometer (Thermo electron).
10. SEQUEST algorithm incorporated in Bioworks software (Thermo).

20.2.8. Protein Glycosylation Analysis by Mass Spectrometry

20.2.8.1. Protein Deglycosylation Using Pngase F

1. PNGase F (QA-Bio, Palm Desert, CA).
2. 5× Reaction Buffer 7.5 for PNGase F: 250 mM sodium phosphate, pH 7.5.
3. Denaturation Solution: 2% SDS, 1 M β-mercaptoethanol. 4. Triton X-100: 15% solution.

20.2.8.2. Glycan Purification

1. Glycoclean R cartridges (Prozyme).
2. 3–5 mL Syringes.
3. Acetic acid and methanol.
4. Black graphitized carbon cartridge (150 mg, 0.5 mL column volume) (Alltech, Deerfield, IL).
5. Trifluoroacetic acid (TFA).

20.2.8.3. Glycan Profiling by LC-MS

1. TSKgel Amide-80 resin (5 μm, 80 Å, Tosoh Biosciences, Montgomeryville, PA).
2. Capillary pump (Ultra-Plus II MD, Micro-Tech Scientific, Vista, CA).
3. ESI-TOF (LCT premier, Micromass/Waters, Milford, MA).
4. Solvent: ACN, formic acid, deionized water.

20.2.8.4. Permethylation of Oligosaccharides

1. Anhydrous dimethyl sulfoxide (DMSO).
2. Sodium hydroxide powder (NaOH).
3. Chloroform (CHCl₃).
4. Methyl iodide (CH₃I).

20.2.8.5. Glycan Structure Analysis

1. MALDI-QIT (Shimadzu Biotech, Manchester, UK).
2. 2',4',6'-Trihydroxyacetophenone monohydrate (THAP) (Sigma-Aldrich).
3. 2,5-dihydroxybenzoic acid (DHB).

20.2.9. Protein Phosphorylation Analysis Using Mass Spectrometry

20.2.9.1. Enzymatic Dephosphorylation

1. Calf alkaline phosphatase.
2. Zip Tips (Millipore).
3. MALDI-TOF MS (Micromass, Inc., TOFSpec2E).

20.3. Methods

20.3.1. Sample Preparation

20.3.1.1. Cell Lysate of Cell Lines and Tissue Sample—Cell lysis is a process that extracts proteins from cells. In order to keep extracted proteins intact in solution, the proteases are inhibited by protease inhibitors and the pH of the lysis buffer is controlled close to physiological pH. Maintaining low ionic strength is a key issue for the subsequent isoelectric focusing or chromatofocusing experiment. Nonionic detergents, reducing agents, and chaotropic agents are used to enhance the solubility of focused proteins.

1. Tissue or cell line samples were quickly thawed and immediately lysed with 2 mL of lysis buffer.
2. Tissue samples could be homogenized on ice using a tissue grinder, mechanical homogenizer, or mini-bead beater according to manufacturer's instruction.
3. The sample with lysis buffer should be vortexed frequently over a period of 1 h at room temperature and then centrifuged at approximately $23,000 \times g$ for 70 min at 4°C.
4. After the supernatant was collected, buffer exchange was conducted using a PD-10 G-25 column. The PD 10 column was equilibrated with the equilibration buffer of the subsequent experiment and then used to exchange the cell lysate from the lysis buffer to the above buffer according to the manufacturer's protocol.
5. Protein concentration in each sample was quantified by the Bradford assay (16). The buffer exchanged protein mixtures were stored in a -80°C freezer until further use.

20.3.1.2. Preparation of Serum Sample

1. 40 cm³ of blood provided by each patient were permitted to sit at room temperature for a minimum of 30 min (and a maximum of 60 min) to allow the clot to form in the red top tubes.

2. The samples were centrifuged at $1,300 \times g$ at 4°C for 20 min. The serum was removed, transferred to a polypropylene capped tube, and frozen. The frozen samples were stored at -70°C until assayed.

20.3.2. Removing High Abundance Proteins Using IgY Antibody Column (For Serum Sample)

The serum proteome is dominated by a few highly abundant proteins which constitute about 90% of the total protein content of the serum. These proteins severely interfere with the quantification and identification of proteins of lower abundance. Most of the high abundance proteins in serum are glycosylated proteins (12) and the presence of these high abundance glycoproteins masks the detection of glycoproteins of lower abundance. Although albumin is not a glycoprotein, it binds to many other glycoproteins. Since many important marker proteins are detected in low concentration in biological samples, removing the high abundance proteins may be a critical strategy for serum biomarker discovery. Human serum was depleted using the ProteomeLab IgY-12 proteome partitioning kit. This antibody column removes albumin, IgG, α 1-antitrypsin, IgA, IgM, transferrin, haptoglobin, α 1-acid glycoprotein, α 2-macroglobin, HDL (apolipoproteins A-I & A-II), and fibrinogen using an affinity column based on avian antibody (IgY)–antigen interactions.

1. Depletion was performed according to the standard operating protocol of this kit. Before setting up the column onto the LC, the HPLC system is flushed with dilution buffer for 20 min at a flow rate of 2 mL/min to make sure that the tube is filled with dilution buffer. Then the antibody column is equilibrated with dilution buffer for 30 min at a flow rate of 2 mL/min to ensure a flat baseline (*see* Note 1).
2. 125- μL serum is diluted in 500 μL dilution buffer and briefly centrifuged using a 0.45 μm spin filter for 1 min at $9,200 \times g$ to remove sample particulates (*see* Note 2).
3. 625- μL diluted sample is injected to the antibody column. The column is washed with dilution buffer at 0.5 mL/min for 30 min. The unbound fraction, which is the top-twelve protein-depleted fraction, flows out at ~ 8 min as shown in Fig. 20.1 and the fraction is collected. The final volume of depleted fraction in dilution buffer is approximately 15 mL. The sample is concentrated and buffer exchanged with equilibration buffer of next step (lectin binding buffer for glycoprotein microarray experiment) using a 5 kDa MW cutoff Amicon filter.
4. The bound fraction, which is the top-twelve protein fraction, is eluted with stripping buffer for 18 min at 2 mL/min and then the neutralization buffer is applied immediately for 10 min at a flow rate of 2 mL/min to regenerate the column. Finally, the column is re-equilibrated with dilution buffer for 10 min at 2 mL/min.
5. 1/10 volume of $10\times$ neutralization buffer is added to the top-twelve protein fraction.
6. Protein assays are performed by the Bradford assay method. The result indicates that around 5–10% of serum proteins are recovered in the depleted fraction.

20.3.3. Lectin Affinity Glycoprotein Extraction

In proteomics studies, fractionation of samples can help to reduce their complexity and to enrich specific classes of proteins for subsequent downstream analyses. One method for

¹Do not freeze this column, store it in 4°C with $1\times$ dilution buffer containing 0.02% (w/v) sodium azide.

²The Proteome Lab IgY-12 columns for different sample capacity (20 μL , 50 μL , 250 μL) are also available from Beckman Coulter.

fractionation and enrichment of glycoproteins is lectin affinity chromatography. Lectins are proteins that are able to specifically and reversibly bind carbohydrates. Lectin affinity chromatography has been used for enrichment of different classes of glycoproteins by many groups (17,18). In this protocol, ConA and WGA lectins are used for extraction of most N-linked glycoproteins. ConA recognizes α -linked mannose including high mannose-type and mannose-core structures. WGA lectin can interact with some glycoproteins via sialic acid residues and it also binds oligosaccharides containing terminal *N*-acetylglucosamine. Almost all N-linked glycoproteins could be extracted since these two lectins can interact with high mannose, complex, and hybrid type glycans.

1. 350- μ L agarose-bound WGA and 250 μ L agarose-bound ConA were packed into disposable screw end-cap spin column with filters at both ends (*see* Note 3).
2. For serum sample, protease inhibitor stock solution is prepared by dissolving one complete EDTA-free Protease inhibitor cocktail tablet in 1 mL H₂O. The stock solution is added to binding buffer and elution buffer at a ratio of (v/v) 1:50. Nondepleted serum sample (~50 μ L) is diluted with binding buffer to make the final volume of 500 μ L. Depleted serum sample (250–600 μ L) is buffer-exchanged into 500 μ L binding buffer using Amicon 5000 MWCO filter.
3. The column is first equilibrated with 500- μ L binding buffer by centrifuging the spin column at 500 rpm for 2 min three times.
4. Diluted sample is loaded onto the column and incubated for 15 min. The column is centrifuged for 2 min at 500 rpm to remove the nonbinding fraction (*see* Note 4).
5. The column is washed with 600- μ L binding buffer twice to wash off the nonspecific binding. The captured glycoproteins are released with 250- μ L elution buffer and the eluted fraction is collected by centrifugation at 500 rpm for 2 min. This step is repeated twice and the eluted fractions are pooled. Proteins bound with WGA are eluted by 0.3 M *N*-acetyl-glucosamine and proteins bound with ConA are eluted by 0.3 M Methyl- α -d-mannopyranoside. The protein assay suggests that around 5–10% of the protein content is extracted from nondepleted serum by the lectin affinity columns.

20.3.4. First Dimension: Fractionation of Proteins Based on pI

20.3.4.1. Fractionation with Microliquid-Phase IEF Cell—For a complex sample like the nondepleted human serum, a one-dimensional separation might not be sufficient to fractionate glycoproteins, especially due to the many high abundance proteins that include glycoproteins. In addition, glycoproteins can exist in multiple glycoforms. Multiple dimension separation is necessary for nondepleted serum glycoprotein mixtures. Here glycoproteins extracted from nondepleted serum are separated based on their pI in the first dimension. The liquid-phase IEF separation is performed using a commercial apparatus called the Rotofor (Bio-Rad). The protocol describes a standard experiment for the micro-Rotofor including materials and step-by-step separation procedures. The Rotofor separates proteins in solution based on their isoelectric point (pI). A protein is positively charged when the pH of the environment is lower than its pI and negatively charged when the pH of the environment is higher than its pI. Upon application of an electric potential, positively charged proteins move toward the anode and negatively charged proteins move toward the cathode until they are neutral. Thus, the proteins are focused at their respective pIs. The

³This spin column is for up to ~50 μ L nondepleted serum sample; larger scale column could be packed if higher amount of sample is needed.

⁴The flow through fraction could be reloaded to the lectin column. Higher recovery could be achieved while nonspecific binding also increases.

typical pH gradient range is 3–10. For the separation of glycoproteins, a Rotofor sample buffer for formation of pH gradient 3–6 is used for a better resolution in the acidic range since most of the glyco-proteins have pI lower than 6. Glycoproteins from human serum can be fractionated into ten pI fractions in this pH range using the microscale Rotofor. Figure 20.2 shows the pH distribution of the ten wells collected from the IEF separation of glycoproteins enriched from nondepleted human serum. The pH of most fractions fall in the range of 3–6 except fraction 1, 9, and 10. The pH of the fractions in the middle increases smoothly from 3 to 6 with a pH difference down to 0.2.

1. Ion exchange membrane is equilibrated in appropriate electrolyte solution overnight before experiment (anode membrane in 0.1 M H₃PO₄ and cathode membrane in 0.1 M NaOH) (*see* Note 5).
2. The pH gradient (3–10) is formed by the 2% Biolyte 3/10 ampholyte. The acidic pH gradient (3–6) is formed by 2% Biolyte ampholyte composed of 10% 3/10 ampholyte, 45% 3/5 ampholyte, and 45% 4/6 ampholyte in the buffer. Chaotropic agent, detergent, and reducing agent are added to the buffer for denaturing and solubilizing the proteins. For separation of lectin-enriched glycoproteins, a 5 k MW cutoff Amicon filter is used to remove the lectin elution buffer from the sample and Rotofor sample buffer is added to the concentrated sample to make the final volume of 2 mL.
3. The focusing chamber is assembled according to the manufacturer's manual. Slowly load the sample through the centermost loading port of the focusing chamber. Make sure that all compartments are filled and no bubbles remain. Seal the loading port with tape (*see* Note 6).
4. Place the focusing assembly into the focusing station. Add 0.1 M H₃PO₄ through the vent hole of anode (red) assembly and 0.1 M NaOH through the cathode (black) assembly. Close the cooling block and switch to cooling setting I (10°C). Equilibrate the system to the setting temperature for 15 min.
5. Set power at a constant of 1 W. The run is complete when the voltage stabilizes. It usually takes 2.5–3 h to complete the run.
6. Turn off power; remove the seal from loading ports. Harvest the fractions by applying vacuum through the vacuum chamber. Ten fractions are collected in the harvest tray.
7. The proteins are separated to ten fractions. The proteins with pI higher than 6 will be focused in the last fraction and the one with pI lower than 3 will be in the first fraction. The pH of collected fractions can be tested with a pH electrode.

20.3.4.2. Protein Fractionation by Chromatofocusing—CF is a column-based chromatographic method that uses a weak anion exchange column to generate a descending linear pH gradient. The exchanger is titrated using ampholytes resulting in a pH separation where fractionation can be performed down to <0.1 pH units. A second dimension such as NPS-RP-HPLC can then be used to analyze each pI fraction. This two-dimensional column technology is available in the Beckman ProteomeLab protein fractionation system kit. In the CF process, accurate pH is measured online by a postdetector pH electrode/cell with low dead volume. Each pH range can be automatically collected using a fraction collector. Up to

⁵The membrane should be stored in electrolyte or distilled water between runs to avoid cracks which cause electrolyte to leak into the focusing chamber.

⁶Adding sample slowly until all of the compartments are filled. Aspirating the sample and reload it if bubble exists. Air bubbles will disrupt the electric field and lead to poor separation.

30 mg proteins could be separated using this technique. The result is a two-dimensional UV map of protein expression, with high reproducibility and excellent fractionation of complex samples such as a cell lines or tissue lysates.

1. CF separation is performed on a HPCF-1D column (250 × 2.1 mm) (Beckman Coulter). The pH of the start buffer is adjusted to pH 8.5 and the pH of elution buffer is adjusted to pH 4.0 with saturated iminodiacetic acid solution and concentrated ammonium hydroxide. Prepared buffer is filtered using a 0.2 µm low extractable membrane filter to remove particles. Prior to sample loading, the column is equilibrated with the start buffer for ~2 h at a flow rate of 0.2 mL/min.
2. After equilibration, 5 mg proteins in start buffer are loaded onto the CF column (*see* Note 7).
3. Elution is achieved by applying a pH 4.0 elution buffer at a flow rate of 0.2 mL/min. A linear pH gradient is generated at the outlet of the column in ~1 h. As a result, proteins elute off the column sequentially according to their pI. Separation is monitored at 280 nm using a UV detector. Fractions are collected every 0.2 pH unit change from pH 8.5 to pH 4.0.
4. The column is washed with 1 M NaCl for 1 h followed by H₂O for 1 h at flow rate of 0.2 mL/min.
5. The column could be washed with isopropanol for 1 h at 0.2 mL/min for removal of the residual proteins and storage of the column (*see* Note 8).

20.3.5. Second Dimension: Fractionation of Proteins Based on Hydrophobicity

Separation in reverse phase (RP)-HPLC is based on hydrophobicity. Particle size, pressure, recovery, stability, sample capacity, and efficiency need to be balanced when selecting the right column for a separation. This protocol employs 1.5-µm ODS (C18) nonporous silica (NPS) particles. NPS columns provide a faster high-resolution separation with improved protein recovery than porous columns since they eliminate issues of macromolecular pore adsorption and stagnant mobile phase in the intraparticle void volume associated with porous stationary phases (19,20). As a result, higher column efficiency is obtained with much improved resolution. NPS packing materials are generally more stable than their porous counterparts at elevated temperatures. In addition, small (1.5-µm) silica particles compensate for the decreased surface area, hence the sample capacity associated with nonporous materials. The high efficiency of NPS 1.5-µm packing allows for reproducible, high-resolution, high-speed protein separations. The enriched glycoproteins or the collection from the pI fractionation (CF or Rotofor) is separated by NPS-RP-HPLC in the second dimension. The LC separation chromatogram of lectin-extracted glycoproteins from depleted serum is shown in Fig. 20.3a and the two-dimensional UV map of micro-Rotofor fractionated glycoproteins (pH 3–6) from nondepleted serum is shown in Fig 20.3b. It is observed that the depletion step effectively simplifies the glycoprotein content in serum (Fig. 20.3a) and most of the glycoproteins in serum are focused in ~pH 5 (Fig. 20.3b). For phosphorylation analysis, the proteins from whole cell lysate are separated by CF as the first dimension and by NPS-RP-HPLC as the second dimension.

1. Sample in aqueous buffer is loaded onto NPS-RP-HPLC. High separation efficiency is achieved by using an ODSII (4.6 × 33 mm) column (*see* Note 9).

⁷Make sure the sample is properly desalted. High ionic strength will interfere with the protein binding. After injecting sample, the pressure may rise rapidly, wait until the pressure decreases and is stable for another injection.

⁸Between washing with isopropanol and 1 M NaCl, the column must be equilibrated with water for at least 1 h. NaCl will precipitate in isopropanol which causes the pressure to increase immediately.

2. To collect purified proteins from NPS-RP-HPLC, the reversed-phase separation was performed at 0.5 mL/min and monitored at 214 nm using a Beckman 166 Model UV detector. Proteins eluting from the column are collected. The reversed phase column is heated to 60°C by a column heater.
3. Both mobile phase A (water) and B (ACN) contain 0.1% v/v TFA. The gradient profile used is as follows: 5–15% B in 1 min, 15–25% B in 2 min, 25–30% B in 3 min, 30–41% B in 15 min, 41–47% B in 4 min, 47–67% B in 5 min and 67–100% B in 2 min.

20.3.6. Protein Microarrays

Protein microarrays are a convenient method to simultaneously scan multiple proteins for a specific property. When separated proteins are exchanged into an appropriate printing buffer, the proteins can be deposited on a covalently modified glass slide or nitrocellulose slide where they bind either covalently or by strong noncovalent interactions. The resulting protein microarray can then be probed with a variety of agents to determine reactivity of proteins. To determine phosphorylated proteins, the slides can be incubated with ionic phosphodies or antiphosphoprotein antibodies. In contrast, the same slides can be probed with multilectins through a biotin–streptavidin detection scheme to elucidate the glycosylation status of the same proteins. Five lectins are used for detection of glycoproteins with different structures. AAL recognizes fucose linked (α -1,6) to *N*-acetylglucosamine or to fucose linked (α -1,3) to *N*-acetylglucosamine. ConA recognizes α -linked mannose including high mannose-type and mannose-core structures. Both MAL and SNA recognize sialic acid on the terminal branches, while SNA binds preferentially to sialic acid attached to terminal galactose in an (α -2,6) and to a lesser degree, an (α -2,3) linkage (21). MAL could detect glycans containing NeuAc-Gal-GlcNAc with sialic acid at the 3 position of galactose (22). In contrast, PNA binds desialylated exposed galactosyl (β -1,3) *N*-acetylglucosamine. In fact, sialic acid in close proximity to the PNA receptor sequence will inhibit its binding. Use of the combination of these five lectins should be highly successful in covering a majority of N-glycan types reported and differentiating them according to their specific structures. When such methods are used in a sandwich assay format, high sensitivity with low limits of detection can be obtained. Studies with standard proteins established the limits of detection to be in the 2.5–5-fmol range with high lectin selectivity (10).

Figure 20.4 shows a glycoprotein microarray section highlighting differences in glycosylation in H factor 1 and alpha acid glycoprotein as determined by processing with five different lectins. When processing human samples, protein concentrations may vary between individuals. In order to eliminate concentration effects all data is normalized with respect to peak heights of the data being analyzed. The bar graph under array images show the extent to which the abundance or specific glycan structures change with respect to disease. For example, it is evident that mannosylation is more abundant in pancreatitis compared to normal serum (as seen in response with ConA) while fucosylation is more abundant in normal compared to pancreatitis serum (as seen in response with AAL).

Figure 20.5 shows sections of a protein microarray probed with ProQ Diamond phosphoprotein gel stain. The Sum52PE cell line was studied and the effect of an inhibitor on protein phosphorylation events in the FGFR2 pathway was studied (15). Both Sum52PE cell lines are separated by CF followed by RP-HPLC. It can be seen that important proteins

⁹For lectin-enriched fraction, the sample may need to be concentrated with 10 k MW cutoff centrifuge filter (Millipore) and rediluted with deionized water since the high concentration of sugar in the lectin elution buffer may cause the column pressure to increase rapidly and shorten the column lifetime.

such as Eps15, STAT3, and MAPK7 interacting proteins were dephosphorylated when the cell line was treated with the inhibitor.

20.3.6.1. Printing

1. Fractionated proteins in 96-well plates are dried down completely using a speedvac centrifuge.
2. The dried fractions are stored at -80°C until needed.
3. Prior to printing, all fractions are resuspended in 15- μL printing buffer and left on a gel shaker at 4°C , overnight.
4. The printing program is set to print five droplets of 500 pL each per spot. All spots are printed 0.6 mm apart and the spot diameter is about 400 μm .
5. At the end of printing all slides are stored in a dessicator cabinet at 4°C .

20.3.6.2. Hybridization of Slides

Hybridization of Slides: Glycoprotein Analysis with Multilectins

1. Printed slides are blocked overnight by placing each slide in a 50 mL tube containing 40 mL of blocking buffer.
2. In the morning the slide is incubated in primary incubation solution. Each slide is placed in a plastic pouch (made in-house using Kpak sealPAK pouches and a heat sealer) containing 4 mL of primary incubation solution. The slides are incubated for 2 h on a rotator at 4°C .
3. After primary incubation, the slides are washed five times for 5 min each in washing solution. Washing is done by placing all slides on a slide rack in a container with probe buffer such that all slides are completely submerged followed by shaking it on a gel shaker.
4. The slides are subsequently incubated in secondary incubation solution using the same procedure as primary incubation. However, the incubation time is only 1 h.
5. After secondary incubation, the slides are again washed five times for 5 min each in washing solution.
6. The slides are then spin-dried using a high-speed centrifuge.
7. It is recommended that all slides be scanned immediately after processing for best results. However, if stored, they should be kept at 4°C in a dessicator cabinet.

Hybridization of Slides: Phosphoprotein Analysis with Antibodies

1. Printed slides are blocked overnight by placing each slide in a 50-mL tube containing 40 mL of blocking buffer.
2. The tubes are attached to a Clay Adam's brand nutator and left with shaking at 4°C .
3. In the morning the slide is incubated in primary incubation solution. Each slide is placed in a plastic pouch (made in-house using Kpak sealPAK pouches and a heat sealer) containing 4 mL of primary incubation solution. The slides are incubated for 2 h on a rotator at 4°C .
4. After primary incubation, the slides are washed five times for 5 min each in probe buffer. Washing is done by placing all slides on a slide rack in a container with probe buffer such that all slides are completely submerged followed by shaking it on a gel shaker.

5. The slides are subsequently incubated in secondary incubation solution using the same procedure as primary incubation. However, the incubation time is only 1 h.
6. After secondary incubation, the slides are again washed five times for 5 min each in probe buffer.
7. Once all antibodies are washed off, the slides are spin-dried using a high-speed centrifuge.
8. It is recommended that all slides be scanned immediately after processing for best results. However, if stored, they should be kept at 4°C in a dessicator cabinet.

Hybridization of Slides: Phosphoprotein Analysis with ProQ Diamond

1. Printed slides are blocked overnight by placing each slide in a 50-mL tube containing 40 mL of blocking buffer (*see* Note 10).
2. The slides are then incubated for 1 h in ProQ Diamond phosphoprotein gel stain with shaking at room temperature. A tube filled with ProQ dye can be used for two slides.
3. After incubation, the slides are destained by washing with destaining solution three times for 10 min each.
4. Once destaining is complete the slides are rinsed with PBS to wash off any residual solutions after which the slides are spun dry using a high-speed centrifuge.
5. It is recommended that all slides be scanned immediately after processing for best results. However, if stored, they should be kept at 4°C in a dessicator cabinet.

20.3.6.3. Slide Scanning

1. Dried, processed slides are scanned on an Axon 4000A scanner.
2. Cy3, Alexa 555 and ProQ treated slides are scanned in the green channel; while Cy5 and Alexa 647 are scanned in the red channel.
3. The photomultiplier tube (PMT) gain should be adjusted until best signal without saturation is seen. Typically, this value ranges from 400 to 700 on our instrument.

20.3.7. Protein Identification by Peptide Sequencing

The purified protein with alterations as detected by protein microarray was collected and subjected to trypsin digestion. The tryptic-digested peptides were further separated by microscale or nanoscale liquid chromatography and analyzed using mass spectrometry. Liquid chromatography-tandem mass spectrometry is a rapid, sensitive method for protein identification. This method allows identification of proteins at low fmol levels and enables identification of numerous proteins in protein mixtures. High-quality MS/MS spectra that contain peptide sequence information could be generated by tandem mass spectrometer. b and y ions are predominantly generated by collision induced dissociation (CID) where peptides were fragmented by collision with nonreactive gases. The commercial available SEQUEST computer program is used to match the sequence information in the spectra to a database of known protein sequences. High-throughput protein identification is enabled by the constantly expanding protein sequence information in various databases.

¹⁰It is possible to probe glass slides with both ProQ Diamond dye as well as antibodies. In this case, the slides should be probed with ProQ dye first followed by incubation with the antibodies. Combining both experiments on one slide reduces the number of slides that need to be used as well as provides valuable information about the type of phosphorylation present on the protein.

20.3.7.1. Tryptic Digestion

1. Fractions obtained from NPS-RP-HPLC (~1–2 µg) are concentrated down to ~20 µL using a SpeedVac concentrator operating at 45°C.
2. 20 µL of 100–300 mM ammonium bicarbonate was then mixed with each concentrated sample to obtain a pH value of ~7.8.
3. 0.5 µL of TPCK modified sequencing grade porcine trypsin was added and vortexed prior to a 12–16 h incubation at 37°C on an agitator.

20.3.7.2. Peptide Separation by Capillary RP-HPLC and Sequencing by Tandem Mass Spectrometry

1. Before separation, peptide samples were dried down and redissolved in 10 µL HPLC grade water.
2. The peptide mixtures were separated using a reverse phase column attached to a Paradigm HPLC pump. For nano-LC-ESI-MS/MS experiments, a nanotrap platform was set up prior to the electrospray source. It includes a peptide nanotrap (0.15 × 50 mm) and a separation column (0.075 mm × 150 mm, C18). The peptide sample was injected and first desalted on the trap column with 5% solvent B (0.3% formic acid in 98% ACN) at 50 µL/min for 5 min.
3. The peptides were eluted using a 45 min gradient from 5 to 95% B at a flow rate of 0.25 µL/min, where solvent A was 0.3% formic acid in 2% ACN.
4. A Finnigan LTQ mass spectrometer equipped with a nano-ESI source was used to acquire spectra. A 75-µm metal spray tip was used and the spray voltage was set at 3 kV. The capillary temperature was 200°C, spray voltage was 4.2 kV, and capillary voltage was 30 V.
5. The instrument was operated in data-dependent mode with dynamic exclusion enabled. The MS/MS spectra on the five most abundant peptide ions in full MS scan were obtained. The high abundant solvent or contamination peaks were placed into an exclusion list to avoid being selected for fragmentation. The normalized collision energy was set at 35% for MS/MS.

20.3.7.3. Database Searching—All MS/MS spectra were searched against the human protein database from SwissProt using SEQUEST algorithm incorporated in Bioworks software and the Swiss-Prot human protein database. The sequence-database search tool was set to expect the following variable modifications: Methionine oxidation (+18 Da), carboxymethylated cysteines if the alkylation was preformed (+57 Da). Trypsin was used as a specific protease with two missed cleavages allowed. Searches were repeated without specification of a protease to search for nonspecific cleavage products. Positive protein identification was accepted for a peptide with X_{corr} of greater than or equal to 3.0 for triply-, 2.5 for doubly- and 1.9 for singly charged ions.

20.3.8. Protein Glycosylation Analysis by Mass Spectrometry

The glycoproteins of interest were collected from LC separation and further characterized using mass spectrometry. The molecular weights and the composition of the enzyme-released glycans could be obtained by mass mapping; a further stage of fragmentation is needed to obtain the sequence or linkage information. The experimentally observed masses were matched with the theoretical value by adding the residue masses of the monosaccharides, the reducing terminal derivative, the mass of any additional groups, and the mass of the adduct. Analyzing the glycosyl moieties on proteins usually involves four

steps (a) glycan release and purification; (b) separating (profiling) the glycans in the pool; (c) derivatization of the glycans; (d) glycan structure analysis using mass spectrometry.

20.3.8.1. Protein Deglycosylation Using Pngase F—Intact N-linked glycans can be released from glycoproteins or glycopeptides enzymatically with peptide *N*-glycosidase F. This enzyme is an amidase that cleaves the linkage between the core GlcNAc and the Asn residue of all asparagine-linked complex, hybrid or high mannose oligosaccharides except alpha(1–3) core fucosylated glycoproteins which is commonly found on plants and can be cleaved by endoglycosidase A. The optimum working pH for this enzyme is 6–10 with the optimum ~7.5. Steric hindrance slows or inhibits the action of PNGase F in some cases. Denaturation of the glycoprotein by heating with SDS and β -mercaptoethanol (β -ME) greatly increases the rate of eglycosylation.

1. Add 10 μ L 250 mM sodium phosphate pH 7.5 and 2.5 μ L 20 $\times$ denaturation solution. Heat the solution at 100°C for 5 min.
2. Cool down the solution and add 2.5 μ L of Triton X-100.
3. Add 2.0 μ L (0.01 U) of PNGase F to the reaction. Incubate overnight at 37°C.

20.3.8.2. Purification of Enzyme-Released Glycans—N-glycanase reaction may contain salt, detergents, proteins (including the enzyme), and released glycans in the reducing form. The glycan sample needs to be purified for further analysis. A reverse phase cartridge could be used to remove the proteins/peptides and detergent since the oligosaccharides were more hydrophilic and will not retain on the cartridge. The glycans could be desalted using a graphitized carbon column as the oligosaccharides are more hydrophobic than salt and will bind this material. The columns or cartridges packed with hydrophilic material could also be used for cleaning the glycan samples.

1. Wash the reverse phase cartridge (e.g., Glycoclean R cartridges from Prozyme) with 5 mL methanol followed by 5 mL 5% acetic acid.
2. The sample should be in aqueous solution with organic solvent less than 5%. Use syringe to pass the sample through the cartridge and collect the eluent. Washed the cartridge with 1 mL 2.5% acetic acid three times and pool the eluent.
3. Wash the graphitized carbon column with ACN and water.
4. Pass the eluent from the RP cartridge through the column and wash the column with 1 mL deionized water three times.
5. The glycan sample was eluted with 2 mL 25% ACN and 2 mL 40% ACN in 0.05% TFA and the eluent was pooled.

20.3.8.3. Profiling the Enzyme-Released Oligosaccharides—Separation of oligosaccharides is challenging because of the complexity of structures. A method commonly used for liquid separation of complex carbohydrate mixtures is HPLC followed by online ESI-MS detection (23–25). Fragile groups, such as sialic acid, which are unstable in MALDI due to in-source decay, are retained using this soft ionization technique (26). For liquid-based separation of glycans, analytical scale normal phase chromatography following fluorescent labeling (27,28) and graphitized carbon chromatography are generally used (29,30). Normal phase separations are based upon hydrophilic interactions of the oligosaccharide hydroxyl groups with the stationary phase. Samples applied in high concentrations of organic solvent adsorb to the column surface and are eluted with an aqueous gradient so that glycans are resolved on the basis of hydrophilicity. Typically, larger oligosaccharides are more hydrophilic than smaller ones, and require higher

concentrations of aqueous solvent to elute. Isocratic structures can be separated using the high resolution of this technique. As shown in Fig. 20.6, N-linked carbohydrates were separated by μ normal phase (0.2×120 mm) LC and detected online by ESI-TOF MS. All the proposed glycan structures are shown on the corresponding doubly charged peaks. Elution positions of the glycans could be expressed as glucose units (GU) by comparison with the elution times of the glucose ladder. The contribution of individual monosaccharides to the overall GU value of a given glycan can be calculated, and these values can be used to predict the structure of an unknown sugar from its GU value.

1. The capillary normal phase column ($200 \mu\text{m} \times 120$ mm) was packed in-house using TSKgel Amide-80 resin ($5 \mu\text{m}$, $80 \text{ \AA}$, Tosoh). A capillary pump was used for separation. The capillary column was directly mounted to a microinjector with a 500-nL internal sample loop. The flow from the solvent delivery pump was split precolumn to generate a flow rate of $\sim 3 \mu\text{L}/\text{min}$.
2. Before injection, purified complex N-glycan mixtures were dried down and redissolved in 80% ACN. The glycans were separated in a 45 min linear gradient which is 100% B to 40% B in 45 min. Solvent A is 50 mM formic acid with pH 4.5 adjusted by ammonium hydroxide. Solvent B is 20% solvent A in ACN.

20.3.8.4. Permethylation of Oligosaccharides—Commonly used derivatization reagents include 2-aminobenzamide, *p*-aminobenzoic acid ethyl ester 2-aminopyridine, etc. This type of derivatization facilitates chromatographic purification and enhances the formation of reducing-end fragment ions in MS experiments. Permethylation is the most popular full derivatization method. Most of the functional groups are converted into their methylated derivatives as permethylation stabilizes the sialic acid residues in acidic oligosaccharides, yielding more predictable ion products when subjected to MS/MS experiments. In addition, methylated sugars seem to ionize more efficiently. Figure 20.7 shows the MS/MS spectrum of a permethylated monosialylated biantennary structure where detailed structure information is yielded by CID fragmentation of permethylated glycans.

1. For permethylation, the glycan sample ($\sim 1 \mu\text{g}$) was dried and add $2 \mu\text{L}$ (this volume should be scaled up if more sample are analyzed) of freshly vortexed DMSO/NaOH suspension (0.2 g NaOH in $500 \mu\text{L}$ anhydrous DMSO) were added. The DMSO/NaOH reagent may be stored in desiccators at room temperature indefinitely but must be revortexed before each use.
2. After incubation for 30 min, one and half volume of iodomethane was added and the mixture was allowed to sit for another 30 min. The same volumes of DMSO/NaOH and CH_3I were added again and were left to stand as above for 60 min.
3. were added and the sample was then Five volumes of CHCl_3 washed with two volumes of water. The water wash was repeated until the aqueous layer was no longer basic. The organic phase was dried in a centrifugal evaporator and the permethylated glycans were redissolved with 50% methonal for MALDI analysis.

20.3.8.5. Glycan Structure Analysis Using Tandem Mass Spectrometry—Fragmentation of glycans occurs by cleavage between the sugar rings (glycosidic cleavage) and cleavage across the rings. Glycosidic cleavage gives information on composition, sequence, and sometimes branching. Cross-ring cleavages provide linkage information. Glycosidic cleavages are the most commonly observed in the collision-induced dissociation. Cross-ring cleavage ions are usually minor peaks in the spectra of sugars ionized as sodium or potassium adducts by low-energy collisions. Abundant cross-ring cleavage ions can be produced in instruments capable of producing high-energy collisions. Glycan structural isomers could be revealed by their specific fragments. A sialic acid is easily lost in positive-

ion MS/MS spectra. In the negative-ion mode, however, sialic acid is relatively stable (31,32). MALDI has proved to be the most effective method to ionize N-link carbohydrates since it does not require the carbohydrates to be derivatized (33). Ion trap instruments have the capability to perform multiple successive stages of fragmentation, which allows probing the details of carbohydrate structure. The interface of MALDI to the quadrupole iontrap-ToF provides a means of performing multiple stages of CID with high mass accuracy and resolution (34–37). This unique instrument is used to study the derivatized or underivatized glycans from the target protein where the carbohydrate could be studied with fragmentation up to MS4 (38,39). The MS/MS spectrum of a premethylated glycan with one sialic acid and a high mannose glycan were shown in Figs. 20.7 and 20.8. It was observed that $[M + Na]^+$ ion were generated in positive mode and the b and y ions from glycosidic cleavage dominate the spectrum.

1. MS and MS² spectra of glycan samples were acquired on a Axima QIT MALDI quadrupole ion trap-TOF (MALDI-QIT). 25 mg/mL 2,5-dihydroxybenzoic acid (DHB) was prepared in 50% ACN with 0.1% TFA.
2. For positive mode analysis, the oligosaccharide solution was applied to the MALDI probe followed by 0.01 M NaCl (0.5 μ L). After the sample was dried in air, 0.5 μ L matrix was added. For negative mode, matrix solution was prepared by dissolving 5 mg/mL 2',4',6'-Trihydroxyacetophenone monohydrate (THAP) and 10 mM ammonium citrate in 50% ACN.
3. Acquisition and data processing were controlled by Launch-pad software (Karatos). A pulsed N₂ laser light (337 nm) with a pulse rate of 5 Hz was used for ionization. Each profile results from two laser shots. Argon was used as the collision gas for CID and helium was used for cooling the trapped ions.
4. TOF was externally calibrated using 500 fmol/ μ L of bradykinin fragment 1–7 (757.40 *m/z*), angiotensin II (1,046.54 *m/z*), P14R (1,533.86 *m/z*), and ACTH (2,465.20 *m/z*) (Sigma-Aldrich) for both positive and negative mode.

20.3.9. Protein Phosphorylation Analysis Using Mass Spectrometry

20.3.9.1. Enzymatic Dephosphorylation—Calf intestine alkaline phosphatase has been widely utilized to dephosphorylate both proteins and nucleic acids. Alkaline phosphatase has a broad pH profile with an optimum between pH 8.0 and 8.5. Calf intestine alkaline phosphatase effectively dephosphorylates proteins containing phosphoserine and phosphothreonine residues, which together account for >97% of the protein-bound phosphate in eukaryotic cells. This enzyme can be utilized for in vitro dephosphorylation of proteins under conditions that do not denature the substrate proteins. In this approach, tryptic digested mixtures from intact and dephosphorylated protein are mass mapped by MALDI-TOF for detection of the shifted mass due to the dephosphorylation. In Fig. 20.9a–d, the MALDI spectra of Rab13 interacting protein and heterogeneous nuclear ribonucleoprotein H before and after dephosphorylation by CAP were shown. The observed mass shifts clearly indicate the phosphorylation on this peptide.

1. After completion of the proteolytic cleavage, the samples were divided into two equal parts. The enzymatic dephosphorylation step was performed by treating one part with five units calf alkaline phosphatase reconstituted in 25 mM NH₄HCO₃ buffer (pH 8.0).
2. The mixture was incubated at 37°C for 2 h, and 2.5% TFA was added to stop the enzymatic reaction. The other part was treated as a control.

3. In preparation for MALDI-MS, the samples were first aspirated using Zip Tips, and then 1 μ L of the eluent was mixed with an equal volume of CHCA matrix solution prepared in 60% ACN/0.1% TFA and spotted on a MALDI plate.
4. Once the spot dried, 1 μ L of 9:1 THAP/DAC matrix solution prepared in 60% ACN/0.1% TFA was applied on top. The spot was allowed to dry slowly afterward.
5. MALDI-TOF MS (Micromass) was used to generate peptide masses of analyzed protein. To verify and locate the phosphorylation sites on proteins, the MALDI-MS spectrum of the phosphorylated (control) and the dephosphorylated samples were compared.

20.3.9.2. Detection of Phosphorylation Site by Neutral Loss Experiment—The low abundance of phosphopeptide coupled with the higher acidity increases the complexity of their analysis by mass spectrometry in positive ESI mode. In tandem mass spectrometry, phosphopeptide precursor ions typically exhibit a prominent neutral loss of phosphate group (98 Da) during fragmentation. To overcome inadequate peptide fragmentation and diagnostic sequence ion information typical in phosphopeptide analysis, a data-dependent experiment could be carried out on a linear ion trap mass spectrometer (LTQ). It selectively triggers MS3 scans on only the MS/MS fragment ions where a specific neutral loss ion (−98 Da) was detected. The increased ion capacity and improved trapping efficiency of the linear ion trap significantly enhances the quality and abundance of ions for the MS3 data that is essential for identification of phosphorylated proteins.

Acknowledgments

This work was supported in part by the National Cancer Institute under grant R01CA106402 (D.M.L.), the National Institute of Health under grant R01GM49500 (D.M.L.), and a Michigan Economic Development Grant MEDC03–622 (D.M.S.). Support was also generously provided by Eprogen, Inc. and Beckman-Coulter. We thank Bio-Rad for the gift of the micro-Rotofor device.

References

1. Rudd PM, Elliott T, Cresswell P, Wilson IA, Dwek RA. Glycosylation and the immune system. *Science* 2001;291:2370–2376. [PubMed: 11269318]
2. Block TM, Comunale MA, Lowman M, et al. Use of targeted glycoproteomics to identify serum glycoproteins that correlate with liver cancer in woodchucks and humans. *Proc Natl Acad Sci USA* 2005;102:779–784. [PubMed: 15642945]
3. Zhao J, Simeone DM, Heidt D, Anderson MA, Lubman DM. Comparative serum glycoproteomics using lectin selected sialic acid glycoproteins with mass spectrometric analysis: Application to pancreatic cancer serum. *J Proteome Res* 2006;5:1792–1802. [PubMed: 16823988]
4. Kobata A, Amano J. Altered glycosylation of proteins produced by malignant cells, and application for the diagnosis and immunotherapy of tumours. *Immunol Cell Biol* 2005;83:429–439. [PubMed: 16033539]
5. Robinson DR, Wu YM, Lin SF. The protein tyrosine kinase family of the human genome. *Oncogene* 2000;19:5548–5557. [PubMed: 11114734]
6. Gschwind A, Fischer OM, Ullrich A. The discovery of receptor tyrosine kinases: Targets for cancer therapy. *Nat Rev Cancer* 2004;4:361–370. [PubMed: 15122207]
7. Cohen P. The role of protein phosphorylation in neural and hormonal control of cellular activity. *Nature* 1982;296:613–620. [PubMed: 6280056]
8. Orzechowski R, Hamelinck D, Li L, et al. Antibody microarray profiling reveals individual and combined serum proteins associated with pancreatic cancer. *Cancer Res* 2005;65:11193–11202. [PubMed: 16322270]

9. Yan F, Sreekumar A, Laxman B, Chinnaiyan AM, Lubman DM, Barder TJ. Protein microarrays using liquid phase fractionation of cell lysates. *Proteomics* 2003;3:1228–1235. [PubMed: 12872223]
10. Patwa TH, Zhao J, Anderson MA, Sime-one DM, Lubman DM. Screening of glycosylation patterns in serum using natural glycoprotein microarrays and multi-lectin fluorescence detection. *Anal Chem* 2006;78:6411–6421. [PubMed: 16970316]
11. Zhu K, Zhao J, Lubman DM, Miller FR, Barder TJ. Protein pI shifts due to posttranslational modifications in the separation and characterization of proteins. *Anal Chem* 2005;77:2745–2755. [PubMed: 15859589]
12. Zhao J, Chang AC, Li C, et al. Comparative proteomic analysis of Barrett's metaplasia and esophageal adenocarcinomas using 2-D liquid mass mapping. *Mol Cell Proteomics* 2007;6:987–999. [PubMed: 16829691]
13. Zhao J, Zhu K, Lubman DM, et al. Proteomic analysis of estrogen response of premalignant human breast cells using a 2-D liquid separation/mass mapping technique. *Proteomics* 2006;6:3847–3861. [PubMed: 16767785]
14. Yoo C, Zhao J, Pal M, et al. Automated integration of monolith-based protein separation with on-plate digestion for mass spectrometric analysis of esophageal adenocarcinoma human epithelial samples. *Electrophoresis* 2006;27:3643–3651. [PubMed: 16927349]
15. Pal M, Moffa A, Sreekumar A, et al. Differential phosphoprotein mapping in cancer cells using protein microarrays produced from 2-D liquid fractionation. *Anal Chem* 2006;78:702–710. [PubMed: 16448042]
16. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248–254. [PubMed: 942051]
17. Rademacher TW, Parekh RB, Dwek RA. Glycobiology. *Annu Rev Biochem* 1988;57:785–838. [PubMed: 3052290]
18. Yang Z, Hancock WS, Chew TR, Bonilla L. A study of glycoproteins in human serum and plasma reference standards (HUPO) using multilectin affinity chromatography coupled with RPLC-MS/MS. *Proteomics* 2005;5:3353–3366. [PubMed: 16052617]
19. Banks JF, Gulcicek EE. Rapid peptide mapping by reversed-phase liquid chromatography on nonporous silica with on-line electrospray time-of-flight mass spectrometry. *Anal Chem* 1997;69:3973–3978. [PubMed: 9322433]
20. Hanson M, Unger KK, Mant CT, Hodges RS. Optimization strategies in ultrafast reversed-phase chromatography of proteins. *Trans Anal Chem* 1996;15:102–110.
21. Shibuya N, Goldstein IJ, Broekaert WF, Nsimba-Lubaki M, Peeters B, Peumans WJ. Fractionation of sialylated oligosaccharides, glycopeptides, and glycoproteins on immobilized elderberry (*Sambucus nigra* L.) bark lectin. *Arch Biochem Biophys* 1987;254:1–8. [PubMed: 3579290]
22. Wang WC, Cummings RD. The immobilized leucoagglutinin from the seeds of *Maackia amurensis* binds with high affinity to complex-type Asn-linked oligosaccharides containing terminal sialic acid-linked alpha-2,3 to penultimate galactose residues. *J Biol Chem* 1988;263:4576–4585. [PubMed: 3350806]
23. Thomsson KA, Karlsson H, Hansson GC. Sequencing of sulfated oligosaccharides from mucins by liquid chromatography and electrospray ionization tandem mass spectrometry. *Anal Chem* 2000;72:4543–4549. [PubMed: 11028608]
24. Kawasaki N, Itoh S, Ohta M, Hayakawa T. Microanalysis of N-linked oligosaccharides in a glycoprotein by capillary liquid chromatography/mass spectrometry and liquid chromatography/tandem mass spectrometry. *Anal Biochem* 2003;316:15–22. [PubMed: 12694722]
25. Wührer M, Deelder AM, Hokke CH. Protein glycosylation analysis by liquid chromatography-mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* 2005;825:124–133.
26. Harvey DJ, Martin RL, Jackson KA, Sutton CW. Fragmentation of N-linked glycans with a matrix-assisted laser desorption/ionization ion trap time-of-flight mass spectrometer. *Rapid Commun Mass Spectrom* 2004;18:2997–3007. [PubMed: 15536626]
27. Naka R, Kamoda S, Ishizuka A, Kinoshita M, Kakehi K. Analysis of total N-glycans in cell membrane fractions of cancer cells using a combination of serotonin affinity chromatography and normal phase chromatography. *J Proteome Res* 2006;5:88–97. [PubMed: 16396498]

28. Merry AH, Neville DC, Royle L, et al. Recovery of intact 2-aminobenzamide-labeled O-glycans released from glycoproteins by hydrazinolysis. *Anal Biochem* 2002;304:91–99. [PubMed: 11969192]
29. Schulz BL, Packer NH, Karlsson NG. Small-scale analysis of O-linked oligosaccharides from glycoproteins and mucins separated by gel electrophoresis. *Anal Chem* 2002;74:6088–6097. [PubMed: 12498206]
30. Hashii N, Kawasaki N, Itoh S, Hyuga M, Kawanishi T, Hayakawa T. Glycomic/glycoproteomic analysis by liquid chromatography/mass spectrometry: Analysis of glycan structural alteration in cells. *Proteomics* 2005;5:4665–4672. [PubMed: 16281179]
31. Deguchi K, Takegawa Y, Ito H, et al. Structural assignment of isomeric 2-aminopyridine-derivatized monosialylated biantennary N-linked oligosaccharides using negative-ion multistage tandem mass spectral matching. *Rapid Commun Mass Spectrom* 2006;20:1480–1481.
32. Deguchi K, Ito H, Takegawa Y, Shinji N, Nakagawa H, Nishimura S. Complementary structural information of positive- and negative-ion MSⁿ spectra of glycopeptides with neutral and sialylated N-glycans. *Rapid Commun Mass Spectrom* 2006;20:741–746. [PubMed: 16456804]
33. Harvey DJ. Matrix-assisted laser desorption/ionization mass spectrometry of carbohydrates. *Mass Spectrom Rev* 1999;18:349–450. [PubMed: 10639030]
34. Fountain ST, Lee H, Lubman DM. Ion fragmentation activated by matrix-assisted laser desorption/ionization in an ion-trap/reflectron time-of-flight device. *Rapid Commun Mass Spectrom* 1994;8:407–416. [PubMed: 7517726]
35. Ding L, Kawatoh E, Koichi T, Smith AJ, Kumashiro S. High efficiency MALDI-QIT-ToF mass spectrometer. *Proc Int Soc Optical Eng* 1999;3777:144.
36. Chien BM, Michael SM, Lubman DM. Enhancement of resolution in matrix-assisted laser-desorption using an ion-trap storage reflectron time-of-flight mass spectrometer. *Rapid Commun Mass Spectrom* 1993;7:837–843.
37. Doroshenko VM, Cotter RJ. A quadrupole ion trap/time-of-flight mass spectrometer with a parabolic reflectron. *J Mass Spectrom* 1998;33:305–318. [PubMed: 9597767]
38. Demelbauer UM, Zehl M, Plematl A, Allmaier G, Rizzi A. Determination of glycopeptide structures by multistage mass spectrometry with low-energy collision-induced dissociation: Comparison of electrospray ionization quadrupole ion trap and matrix-assisted laser desorption/ionization quadrupole ion trap reflectron time-of-flight approaches. *Rapid Commun Mass Spectrom* 2004;18:1575–1582. [PubMed: 15282782]
39. Ojima N, Masuda K, Tanaka K, Nishimura O. Analysis of neutral oligosaccharides for structural characterization by matrix-assisted laser desorption/ionization quadrupole ion trap time-of-flight mass spectrometry. *J Mass Spectrom* 2005;40:380–388. [PubMed: 15712371]

Fig. 20.1.

UV Chromatogram of three 125-μL serum depletion from three patients by IgY antibody column to remove 12 high abundant proteins. During the binding process, the fraction flowing through was collected as top-twelve depleted serum fraction and the top-twelve retained protein fraction was collected during elution. The absorption was set at 280 nm.

Fig. 20.2.

The pH of the ten fractions collected from micro-Rotofor run was measured. The glycoproteins enriched from 750 μ L nondepleted human serum by ConA affinity column were separated by the pH gradient 3–6 formed by mixing of the Biolyte ampholyte 3/10, 3/5, and 4/6.

Fig. 20.3.

a NPS-RP-HPLC separation of ~30 μ g glycoproteins extracted from depleted serum by ConA-WGA lectin column. The UV wavelength is 214 μ m. Flow rate is 0.5 mL/min. **b** Two-dimensional UV map generated from the NPS-RP-HPLC separation of glycoproteins extracted from nondepleted human serum. Rotofor fractions 2–9 (pH 3–6) are shown. The *x*-axis is the fraction number corresponding to Fig. 20.2 and the *y*-axis is the retention time. The chromatogram shown in *left* is the lane #4 (fraction #5).

Fig. 20.5.

Phosphoprotein microarray images of two cell lines highlighting phosphorylation differences between the two. On the *left* is the Sum52PE cell line while on the *right* is the Sum52PE treated with inhibitors of a key signaling pathway. Notice how some proteins are dephosphorylated in this cell line (Proteins of interest are highlighted in *blue boxes* and their IDs are indicated by *arrows*.) (Reproduced from ref. ¹⁵ with permission from ACS).

Fig. 20.6. N-linked carbohydrates were separated by μ normal phase LC and detected online by ESI-TOF MS. The combined mass spectra of retention time 28–29 min are shown. All of the peaks presented here are doubly charged. The spectrum for cancer sample is shown above the normal sample. The proposed glycan structures are shown on the corresponding peaks. *Triangle*, fucose; *star*, N-acetyl neuraminic acid (sialic acid); *square*, HexNAc; *circle*, mannose; *diamond*, galactose.

Fig. 20.7. Fragmentation spectra of permethylated sialylated oligosaccharide $[M + Na]^+$ obtained from MALDI-QIT MS in positive mode. The structure is monosialylated bi-antennary structure with fucose: $(Hex)_2 (HexNAc)_2 (NeuAc)_1 (Fuc)_1 + (Man)_3 (GlcNAc)_2$ at m/z 2,604.95.

Fig. 20.8.

Fragmentation spectra of high mannose oligosaccharide $[M + Na]^+$ obtained from MALDI-QIT MS in positive mode. The composition of this structure is $(\text{Hex})_5 (\text{HexNAc})_2 + (\text{Man})_3(\text{GlcNAc})_2$ at m/z 1,257.0 (reproduced from ref. ¹⁰ with permission from ACS).

Fig. 20.9.
a MALDI spectrum of Rab13 interacting protein (MIRab13) (MICAL-like protein 1) obtained before (*bottom*) and after (*top*) dephosphorylation by CAP. **b** MALDI spectrum of heterogeneous nuclear ribonucleoprotein H (hnRNP H) obtained before and after dephosphorylation (reproduced from ref. ¹⁵ with permission from ACS).

Scheme 20.1.

Flow chart of the strategies introduced in this protocol for the protein glycosylation and phosphorylation screening. The *dashed line* frame is an optional step in the procedure.