Published in final edited form as:

Calcif Tissue Int. 1980 ; 31(2): 173–181.

Transcapillary Exchange and Retention of Fluoride, Strontium, EDTA, Sucrose, and Antipyrine in Bone

Gerard J. Lemon, David R. Davies, Sean P.F. Hughes, James B. Bassingthwaighe, and Patrick J. Kelly

Mayo Clinic and Mayo Foundation, Rochester, Minnesota 55901, USA

Summary

The extractions of $^{85}\text{Sr}^{2+}$, $^{18}\text{F}^-$, sucrose- ^{14}C , EDTA- ^{51}Cr , and antipyrine- ^{14}C in bone were determined by the multiple indicator-dilution method. Fluoride and strontium extractions were 18 to 70% during a single transcapillary passage, and those of EDTA and sucrose were from 11 to 59%, whereas extraction of antipyrine was 87%. Injections of $^{85}\text{Sr}^{2+}$ and $^{18}\text{F}^-$ made when perfusion was done alternately with blood and plasma resulted in similar fractional extractions. When flow and extraction were measured simultaneously, extraction was related inversely to flow.

Keywords

Bone blood flow; Capillary permeability; Tracer exchange

Introduction

The primary objective of this study was to ascertain the transcapillary extractions, E , of fluoride, strontium, sucrose, and EDTA during a single passage through the capillary bed of tibial bone. The tracers fluoride and strontium are of particular interest in bone because of their known uptake and deposition in the crystalline matrix of bone. Because strontium and other tracers are only partially extracted in the capillary bed of bone [1,2], a particular provocation for our examination of fluoride uptake was the report by Wootton [3], who used fluoride deposition as a measure of regional bone blood flow on the assumption that its extraction was complete during a single pass through the capillary. Fluoride tracer has been combined with EDTA- ^{51}Cr as a clinical method for measuring skeletal blood flow. The assumption behind this technique is that $^{18}\text{F}^-$ is completely extracted in a single transcapillary passage. The EDTA- ^{51}Cr function is assumed to be equivalent to that part of the $^{18}\text{F}^-$ function that reflects tracer movement into the extracellular fluid but not into bone.

The tracer antipyrine is a lipid-soluble molecule that has been shown in previous studies to be freely diffusible across the capillary barrier [4]. Since the maximum extraction of water, for example, in the lung, is not 100% [5], it would seem unlikely that even antipyrine would be totally extracted. For this reason, comparison of antipyrine with fluoride allows a useful qualitative comparison of a lipophilic substance and a hydrophilic ion.

A final justification for the study is the wide interest in fluoride as a possible pharmacologic agent in the treatment of osteoporosis and the osteopenia of multiple myeloma [6,7].

Materials and Methods

Six sets of experiments were done (Table 1). The tibial nutrient artery and ipsilateral femoral vein of adult dogs were cannulated for introduction of radioisotopes and collection of venous samples, as described previously [1,2]. The femoral vein is cannulated for outflow collection. The blood will include that from tissues other than bone, but this merely dilutes the tracers but does not change their amounts or their relative concentrations. Our past studies indicate that the fractional recovery rate of a bone-seeking isotope in such an experimental setup is 99% [8]. Using venous outflow from the limb maximizes the amount of blood collected and gives more accurate results than collecting nutrient vein blood [1]. In the first three sets of experiments, the nutrient artery was tied off proximally. Nutrient artery cannulation does reduce the flows somewhat; earlier studies showed a decrease in diaphyseal ^{85}Sr clearance by one-third [1]. In the last three sets of experiments, the tibial capillary bed was perfused at a chosen constant blood flow by the use of a Holter pump (model A175, 110V, Extracorporeal Medical Specialties). The flow was nonpulsatile.

Albumin labeled with ^{51}Cr or $^{99\text{m}}\text{Tc}$ (Squibb) was used as a non-permeating intravascular reference tracer. The permeant test tracers were carrier-free ^{18}F as the fluoride ion in isotonic saline (Medi+Physics and Phoenix Memorial Hospital, University of Michigan), carrier-free $^{85}\text{SrCl}_2$ (Squibb), ^{14}C -labeled antipyrine in isotonic saline (New England Nuclear), and ^{51}Cr -labeled EDTA (New England Nuclear). The injection mixture consisted of 15 or 60 μCi of $^{18}\text{F}^-$, 15 μCi of albumin- ^{51}Cr or 15 μCi of albumin- $^{99\text{m}}\text{Tc}$, 5 μCi of sucrose- ^{14}C , 20 μCi of EDTA- ^{51}Cr , and 5 μCi of antipyrine- ^{14}C or 5 μCi of ^{85}Sr . In the first three sets of experiments, isotonic saline was injected over a period of 1 min, and in the second and third sets over 47 s, with the use of an infusion pump (Harvard Model 932) for the injection. In the last three sets of experiments, a 1- to 2-s injection of 0.25 ml of injectate was followed by a short flush of 0.1 ml. In the first three sets of experiments, the tracer was injected as an extended bolus at a constant rate into the distal segment; this was done because the blood flow through the tibial diaphysis is collateral flow of about 2.5 ml/100 g/min, estimated from clearance values corrected by net extraction [1] or estimated by microspheres in standing, conscious dogs [9]. Further, we wished to keep injectate flow to a modest fraction of the total in order to avoid disturbing the rate of flow or the perfusion pressure. Venous outflow dilution curves were obtained from multiple sequential samples from the femoral vein. In collecting the total femoral vein flow, 18 or more sequential samples, each of 5 or 10 s duration, were obtained, beginning with the start of the injection.

For sets four, five, and six, the experimental setup was modified to provide continuous perfusion of the tibial nutrient artery with a Holter pump. The dog was perfused with its own blood, between 200 to 300 ml of which had been removed and replaced with a similar volume of Rheomacrodex. Hematocrits were known in all experiments, and in set five, we perfused the bone alternately with plasma and with blood having a hematocrit of 20%. Multiple dilution curves were obtained in some dogs (Table 1). In the dogs in which multiple curves were obtained, pre-injection standards were taken from the venous blood to correct for residual isotopic activity. They were very low. The volume of blood was noted at frequent intervals so that accurate estimates of flow could be determined. At the end of the experiments, the tibia was removed, cleaned of soft tissue, and weighed. Since hematocrits were known, plasma flow, F_s (ml/100 g bone per min), could be calculated when blood was used as the perfusate.

Tracer Activity of Samples

In these experiments, activity was measured in samples of blood, plasma, and bone. Because an appreciable amount of $^{18}\text{F}^-$ could enter the red blood cells [10], particularly in the minutes between collecting samples and taking aliquots for counting, 1 ml of whole blood was taken from each sample, and the activities of $^{18}\text{F}^-$ and ^{51}Cr were determined in a well-type scintillation detector (Nuclear Chicago model C 120-1). The $^{18}\text{F}^-$ was allowed to decay out, and then the 1-ml samples of whole blood were recounted. This allowed calculation of the activity due to $^{18}\text{F}^-$ and ^{51}Cr . From aliquots that were centrifuged, the activities of ^{51}Cr and ^{14}C were determined in 0.2 ml of plasma by an automatic liquid-scintillation system (Nuclear Chicago Mark II) with the techniques of sample preparation and calculation of activity due to each isotope which have been described previously [2]. Samples of blood were used in the place of plasma in those experiments in which antipyrine- ^{14}C was given; these were incubated with 60% perchloric acid and 30% hydrogen peroxide before the addition of the liquid scintillator.

In experiments in which $^{18}\text{F}^-$ and $^{85}\text{Sr}^{2+}$ were used together counting was performed on a Beckman 310 instrument, and before the experiment, the windows to be used were determined with the use of $^{22}\text{Na}^+$, a positron emitter like $^{18}\text{F}^-$. Before the experiment, mixtures of the three isotopes were examined on a multichannel analyzer. The efficacy of counting $^{18}\text{F}^-$ at its main peak (0.5 meV) was determined and found to be satisfactory ($\approx 24\%$). The windows were set to count $^{99\text{m}}\text{Tc}$, the main peaks of $^{85}\text{Sr}^{2+}$ and $^{18}\text{F}^-$ (both 0.5 meV), and the coincidence peak of $^{18}\text{F}^-$ (1.02 meV). Spillover of $^{18}\text{F}^-$ into $^{85}\text{Sr}^{2+}$ windows and of $^{85}\text{Sr}^{2+}$ into $^{99\text{m}}\text{Tc}$ windows was determined over a wide range of counts. Strontium activity in each sample was determined by correcting for crossover and background, and then again, with improved accuracy, by recounting after the $^{18}\text{F}^-$ and $^{99\text{m}}\text{Tc}$ had been allowed to decay out.

For the comparisons of sucrose- ^{14}C and EDTA- ^{51}Cr , the albumin- $^{99\text{m}}\text{Tc}$ and EDTA- ^{51}Cr counts were determined on a Beckman 310 gamma counter, allowance being made for EDTA- ^{51}Cr spillover into the $^{99\text{m}}\text{Tc}$ window. After $^{99\text{m}}\text{Tc}$ had decayed out, counts were determined by means of a Packard liquid-scintillation system. Quench data, background counts, counting efficiency, and spillover were determined, and polynomial equations were fitted to these by computer. From these data, counts of sucrose- ^{14}C were obtained. Corrections for decay are made for all tracers; this is especially needed in the instances of ^{18}F and $^{99\text{m}}\text{Tc}$.

Computations

The first stage was to calculate, from the outflow sample counting, the form of the dilution curve. The fraction of the injected dose of tracer appearing in the femoral vein per second, $h(t) \text{ s}^{-1}$, was determined as previously described [1,2], from the outflow concentration-time curve $C(t)$, in counts per minute per milliliter. Because the injection was prolonged, $h(t)$ had the same dimensions as the curve of the probability density function of transit times but not the same shape; in this study, $h(t)$ is not the impulse response but rather the response to a pulse of finite duration. A mixture of two or more tracers is injected into the cannulated tibial nutrient artery. One of these is the intravascular reference tracer. The other tracers are the permeant tracers. By a sampling of venous outflow, it is possible to measure the fraction of the injected dose of each tracer that appears in the outflow per second, $h_D(t)$. The femoral vein is cannulated for outflow collection. Although this blood may include blood from tissue other than bone, this merely dilutes the tracers and does not change their amounts or relative concentrations. Previous studies have indicated that 99% of the tracer can be accounted for in this experimental setup [10]. In the last three sets of experiments, the injection was nearly an impulse injection.

The instantaneous fractional extraction, $E(t)$, for each test tracer is calculated from

$$E(t) = \frac{h_R(t) - h_D(t)}{h_R(t)} \quad 1a$$

in which the subscript R indicates the intravascular reference tracer and D denotes a diffusible permeating tracer. When the injection is a long pulse, only the earliest values of $E(t)$ will provide a measure of the unidirectional flux from blood, just as is true for the impulse response. Mathematically, it has been shown [11] that the extent of intravascular dispersion, either in inflow before the capillary bed or in outflow downstream from the exchange region, should have a negligible influence on the observed $E(t)$, even though the forms of the recorded indicator-dilution curves are markedly changed. Thus the interpretation of $E(t)$ and in particular of the highest value, E_{max} , which occurs fairly early, is not changed by the form of the injection used in these experiments. Choosing the appropriate extraction with which to estimate flux of ions across the capillary barrier is difficult. The form of the curves is seen in Figure 1 for the prolonged injections when the tibia was autoperfused (sets one, two, and three) and in Figure 2 for the short injection into the pump-perfused nutrient artery (sets four, five, and six). For the autoperfused tibias, E_{max} was the highest value of the smoothed curve of $E(t)$ and always occurred on the upslope of the reference curve, as previously described [1,2].

The curves in the pump-perfused dogs for which the short, small volumes of injections were used were more sharply peaked (Fig. 2). From a computer analysis of multiple indicator-dilution curves in the heart (Levin, Holloway, Bassingthwaighite, unpublished data), with the use of the cell-uptake model of Rose et al. [12], and with flow heterogeneity estimated by means of micro-sphere deposition, it is possible to derive a manual method for obtaining the representative E from $E(t)$ from the early part of the dilution curve.

Each curve of $E(t)$ was smoothed by fitting a smooth curve through the data from the 15th to the 40th second; the value E was taken from this curve at the sample time nearest the time when half of $h_R(t)$ had passed, which is at either the peak or one sample beyond the peak of the reference curve. These instantaneous extractions are termed E_{Sr} , E_F , E_{Suc} , or E_{EDTA} .

The tracers strontium, fluoride, and EDTA were retained in the tissue for a long time in comparison with albumin and sucrose. A measure of this retention at any specified time, T , after injection, an integral or net extraction, is given by

$$E_{net}(T) = \frac{\int_0^T [h_R(t) - h_D(t)] dt}{\int_0^T h_R(t) dt} \quad 1b$$

After all the reference tracer has washed out, $E_{net}(T)$ becomes identical to the residue

function for the permeating tracer, $R(t) = 1 - \int_0^T h(t) dt$, which is the fraction of the injectate retained by the tissue.

Results

Initial Experiments on Fluoride Extractions

The first set of experiments, on eight dogs, was designed for estimating E , the extraction for fluoride; the injectate contained albumin, fluoride, and, as an inert but permeating reference, sucrose- ^{14}C .

One set of dilution curves of the sort shown in Figure 1 was obtained from each dog. There was first an appreciable delay before the appearance of any tracer. This delay represents the shortest transit time and is the sum of transit time through the nutrient arterial inflow and that through the bone capillary bed, the outflowing veins, and the collecting catheters. The difference in height between the test indicator curve and the reference indicator curve, relative to the height of the reference indicator curve, is the fractional extraction, $E(t)$, at that instant (Eq. 1b).

Accordingly, the curves of $E(t)$ (lower panel, Fig. 1) increase rapidly to a peak (E_{max}) and then decline because of the back diffusion of permeant tracer from the extravascular fluid compartment to the capillary. In eight such sets of dilution curves, the observed E_{max} for $^{18}\text{F}^-$ was 0.70 ± 0.08 (SD), and the E_{max} for sucrose- ^{14}C was 0.47 ± 0.13 .

Effects of Pre-equilibration with Erythrocytes on Fluoride Extraction

If fluoride entered red blood cells before or during transcapillary passage, the effective flow, F_s , would be different from the plasma flow. Both F_s and the rate at which fluoride enters or leaves red cells would affect its extraction. The second set of experiments along with set five were therefore designed to test whether the presence of red cells affected the extraction of fluoride either in absolute terms or relative to tracers that are known to be carried in plasma, that is, $^{85}\text{Sr}^{2+}$, sucrose- ^{14}C , and EDTA- ^{51}Cr .

Two injections were given to each of three dogs, with an interval of 15 min between them. One of each pair of injection mixtures was equilibrated with whole blood for 5 min before injection. In these particular studies, venous samples were obtained from a small catheter (Intramedic PE240) that was inserted into the femoral vein via a tributary. A pump (Holter RL 175) was used to give a venous sampling rate of about 30 ml/min. Results are summarized in Table 2.

Comparisons of Antipyrine and Fluoride

This third set of experiments concerned the extractions of fluoride and antipyrine. The antipyrine served two purposes. One was to provide an estimate of the maximum extraction that might occur with a flow-limited indicator. Previously it was shown that the washout of antipyrine is apparently flow limited and is not permeability or diffusion limited [4,13]. The other purpose was to compare $^{18}\text{F}^-$ with a tracer that equilibrates essentially instantaneously with erythrocytes, for the effective F_s for antipyrine is the flow of whole blood, F_B . (Actually, it is slightly less, since at equilibrium the concentration of antipyrine in erythrocyte water is 1.1 times that in plasma water [14], but there is only 0.74 ml/ml water in red blood cells compared with 0.96 ml/ml in plasma, so that $F_{AP} = F_B [(1 - \text{hematocrit}) + 1.1 \text{ hematocrit} (0.74/0.96)]$.)

One nutrient artery injection of albumin, $^{18}\text{F}^-$, and antipyrine was made in each of eight dogs. A set of outflow dilution curves is shown in Figure 3. For the eight dogs, the E_{max} was 0.65 ± 0.13 for $^{18}\text{F}^-$ and 0.87 ± 0.05 for antipyrine.

Pump Perfusion Experiments with Strontium and Strontium and Fluoride

The fourth set of experiments represents a group of experiments that were performed to establish the method of perfusion (Table 3 and Fig. 4). The fifth set of experiments provides comparison between the two ions F^- and Sr^{2+} in the same dog. We perfused the bone alternately with plasma and with blood having a hematocrit of 20% (Table 3).

The injections were of about 1-s duration; a set of dilution curves is shown in Figure 2. In each of two dogs, five sets of dilution curves were obtained—three with plasma perfusion

and two with blood perfusion—giving a total of 10 comparisons. Of the 10 curves, 8 showed $E(t)$ s that increased continuously through the main part of $h(t)$, from about the 15th to the 40th second.

Values for E are given in Table 3. There was no systematic influence of the presence of red cells on E .

Values for $E_{net}(t)$, calculated by Eq. 1b and illustrated in Figure 2, showed that a greater fraction of fluoride was retained than of strontium at times up to 3 min. The paired values of E_{net} (2 min)—dogs H159 and H236, Table 3—show 29% retention for F^- and 34% for Sr^{2+} at this time when the albumin had nearly completely washed out. Values for E_F , E_{Sr} , and E_{net} for these experiments are illustrated in Figure 4.

Pump Perfusion Experiments with Sucrose- ^{14}C and EDTA- ^{51}Cr

The final set of experiments, set six, was a comparison between two molecules of similar molecular weight—EDTA- ^{51}Cr (molecular weight 336) and sucrose- ^{14}C (molecular weight 342). EDTA is a positively charged molecule and sucrose is without charge. Their volumes of distribution in canine tibial cortical bone are known; that of EDTA is approximately twice that of sucrose or inulin, tracers employed by us to identify the extracellular fluid space of bone [15,16].

The results are summarized in Table 4 and Figure 5. $E(t)_{sucrose}$ and $E(t)_{EDTA}$, at either the peak of the reference curve or their net extractions at 2 min, are inversely related to flow.

Discussion

The main observations are that none of the ions or molecules is completely extracted during a single passage. A substantial fraction of fluoride, strontium, and EDTA is retained in the large extravascular volume of distribution of these substances by the end of 2 min. This is less so in the instance of sucrose.

The estimates for extraction of antipyrine, 0.87, are consistent with the maximum extraction for other tracers in situations in which the exchange is entirely flow limited—for example, water in the pulmonary circulation [5]—and what we would expect in view of our earlier observations [4], that iodoantipyrine is flow limited in its washout from the tibia. The relatively low extraction for fluoride compared with iodoantipyrine suggests that its exchange should be barrier-limited, presumably at the capillary membrane.

It might be argued that the extraction of antipyrine is greater than that of ^{18}F because of its lipophilic qualities so that it might avidly bind to fat marrow. However, this would be the case only if the oil:water solubility ratio of antipyrine were high. This is the case with regard to the tracer xenon-133, and for that reason we have used a lipophilic tracer, antipyrine, which has a much lower oil:water solubility ratio than xenon-133 [17–19]. Recent experiments indicate that the blood-cortical bone partition coefficient, λ_c , is 0.27 and the blood-marrow partition coefficient, λ_m , is 0.17. These values were derived from two previously reported studies [20,21]. They are consistent with the water content of cortical bone determined by steady-state experiments with tritiated water [22] and the water content of tibial marrow determined by desiccation [13].

Regarding the radionuclides ^{85}Sr and ^{18}F , it is conceivable that these tracers might concentrate in areas of bone formation or low density, sometimes referred to as “hot spots.” Autoradiographic studies [23,24] have shown that during the first 10 min after injection of radionuclides, tracer uptake is seen about the surface where vessels are known to traverse, as

illustrated in Figures 1 and 2 of the article of Pasternak et al. [24], and not about areas of low density.

Problems of methodology include the use of albumin as a reference tracer, the choice of the appropriate extraction value, and the role marrow tissue might have in extraction of tracer.

The appropriateness of albumin as a reference is dependent on its escape from the capillary bed being less than 2% on single passage. Its use is not at all negated by Owen and Triffitt's observation [25] in young rabbits that the clearance of albumin is about once per hour; this translates to $(PS/V_{ISF})_{alb} = 1 \text{ h}^{-1} = 2.8 \times 10^{-4} \text{ s}^{-1}$, which is low compared with that estimated for sucrose from this study—about $4.3 \times 10^{-3} \text{ s}^{-1}$. This value was derived as follows:

$$PS/V_{ISF} = (PS/F_s) \cdot (F_s/V_{ISF}) = -F_s \log_e(1-E_{max})/V_{ISF} = -0.008^1 (\text{ml g}^{-1} \text{min}^{-1}) \log_e(1-0.44)/0.018^2 \text{ ml g}^{-1} = 4.3 \times 10^{-3} \text{ sec}^{-1}$$

Considering that Owen and Triffitt emphasize that permeation of albumin is about 10 times greater in young than in adult rabbits (see Fig. 8 in ref. 25), we can anticipate that in our experiments the maximum extraction of albumin could not be more than 2%, which would not invalidate its use as a reference tracer.

When first planning this study of fluoride and strontium extraction, we recognized that fluoride might enter erythrocytes coming from the collateral vessels to mix with the injectate before transcapillary passage. This meant that we would not know whether there was a “red blood cell carriage effect” [26], whether fluoride exchange with red cells was rapid, or whether the flow, F_s , for fluoride was the same as for sucrose or somewhat greater; thus we knew that information on these points should be obtained before comparison of extraction of fluoride with flow.

From the data of Tosteson [27] on beef erythrocytes, one would have thought that fluoride might enter erythrocytes within seconds. However, our perfusion experiments with and without red blood cells do not support this idea. The answer may be that the red blood cells of dogs are quite different from those of cows and sheep. This would not be wholly unexpected, because the potassium concentration in the red blood cells of dogs is only 10 mM, compared with 136 mM for humans [28]. Also, Dalmark [29] observed that, of all the halides, fluoride produces the least inhibition (both competitive and noncompetitive) of chloride transport in human red blood cells, a finding that suggests that fluoride transport by the chloride carrier may be minimal. It would be desirable to have a direct experiment on fluoride uptake in dog red blood cells, preferably with a rapid-reaction apparatus.

If there is rapid entry of fluoride, the value F_s for fluoride would be increased to about 1.12 F_{plasma} ; that is, in the 20% hematocrit experiments, it would be $F_{plasma} 0.8 (0.65 \times 0.7 \times 0.2 + 0.8)$, in which 0.65 is the water fraction of red blood cells, 0.7 is an expected Donnan factor, and 0.2 is the hematocrit. Such an occurrence would not change the relationships observed in Figure 4. Although there may be increased solubility of F^- for endothelial cells and the cells of bone, this ion is not freely diffusible in the sense that antipyrine is, and therefore there is some limitation of diffusion at the capillary barrier.

It would appear that there is a disadvantage in employing fluoride tracer similar to that in using strontium tracer—namely, that $^{18}F^-$ measures clearance and not flow. At flows similar to those we have observed in the tibial diaphysis of the conscious dog—2.5 ml/100 g/min—net extraction is about 0.30 (Fig. 4).

Concerning marrow effects on tracer uptake, the microvasculature of the tibial diaphysis is composed of the capillaries of the Haversian canals. The marrow of mature dogs is a fatty marrow and serves mainly as a supporting structure for conduit vessels entering and leaving the cortex [15]. Cortical bone forms the major fraction of the mass of the tibial diaphysis, 0.90 [1]. Furthermore, uptake of ^{45}Ca by dog tibial marrow after perfusion into the nutrient artery is small and is only 0.4% of total bone uptake [30]. It is recognized that a tracer injected into the nutrient artery will reach the capillary beds of marrow tissue as well as bone tissue. However, studies with microspheres indicate that the percentage of blood flow to bone tissue in a mature dog tibia is approximately 89%—that is, flow in the tibia is mainly to bone [31]. Thus we do not believe that the presence of blood flow to marrow influences our interpretation.

Studies in this laboratory by Morris et al. [16,22] have shown that the volume of distribution for EDTA- ^{51}Cr is twice that for sucrose- ^{14}C —0.0758 ml/ml compared with 0.0429 ml/ml. EDTA does not enter the cell, and it seems unlikely that the chromium-51 label separates from EDTA in competition with calcium [32]. The stability constant for CrEDTA is $10^{23.4}$, compared with $10^{10.6}$ for CaEDTA [33–35]. Furthermore, studies by both Lopez-Curto et al. [15] and Morris et al. [16] indicated that the concentration of ^{51}Cr in blood was the same in dogs with ligated kidneys at 3 and 4 h.

Therefore, the lower net extraction for EDTA- ^{51}Cr compared with sucrose- ^{14}C is likely due to a larger volume of distribution for EDTA, resulting in less back diffusion of tracer. In view of these observations, plus the close agreement between the volume of distribution of inulin (0.0420 ml/ml) and that of sucrose (0.0429 ml/ml) in bone, we are led to view with skepticism the use of EDTA as an extracellular fluid space marker of bone [36].

Acknowledgments

This investigation was supported in part by Research Grants AM-15980 and HL-19139 from the National Institutes of Health, Public Health Service. The authors are grateful to Glenn Christensen for technical help and Rose M. Garmers for preparation of the manuscript.

References

1. Cofield RH, Bassingthwaighe JB, Kelly PJ. Strontium-85 extraction during transcapillary passage in tibial bone. *J. Appl. Physiol* 1975;39:596–602. [PubMed: 1104562]
2. Davies DR, Bassingthwaighe JB, Kelly PJ. Transcapillary exchange of strontium and sucrose in canine tibia. *J. Appl. Physiol* 1976;40:17–22. [PubMed: 1248978]
3. Wootton R. The single-passage extraction of ^{18}F in rabbit bone. *Clin. Sci. Mol. Med* 1974;47:73–77. [PubMed: 4855089]
4. Kelly PJ, Yipintsoi T, Bassingthwaighe JB. Blood flow in canine tibial diaphysis estimated by iodoantipyrine- ^{125}I washout. *J. Appl. Physiol* 1971;31:38–47. [PubMed: 5556961]
5. Perl W, Silverman F, Delea AC, Chinard FP. Permeability of dog lung endothelium to sodium, diols, amides, and water. *Am. J. Physiol* 1976;230:1708–1721. [PubMed: 937558]
6. Kyle RA, Jowsey J, Kelly PJ, Taves DR. Multiple-myeloma bone disease: the comparative effect of sodium fluoride and calcium carbonate or placebo. *N. Engl. J. Med* 1975;293:1334–1338. [PubMed: 1102987]
7. Riggs BL, Hodgson SF, Hoffman DL, Kelly PJ, Johnson KA, Taves D. Treatment of primary osteoporosis with fluoride and calcium: clinical tolerance and fracture occurrence. *J.A.M.A.* (in press).
8. Hughes SPF, Davies DR, Bassingthwaighe JB, Knox FG, Kelly PJ. Bone extraction and blood clearance of diphosphonate in the dog. *Am. J. Physiol* 1977;232:H341–H347. [PubMed: 402827]
9. Morris MA, Kelly PJ. Use of tracer microspheres to measure bone blood flow. *Trans. 25th Annu. Meet. Orthop. Res. Soc* 1979;4:16.

10. Carlson CH, Armstrong WD, Singer L. Distribution, migration and binding of whole blood fluoride evaluated with radiofluoride. *Am. J. Physiol* 1960;199:187–189. [PubMed: 13807751]
11. Bassingthwaite JB. A concurrent flow model for extraction during transcapillary passage. *Circ. Res* 1974;35:483–503. [PubMed: 4608628]
12. Rose CP, Goresky CA, Bach GG. The capillary and sarcolemmal barriers in the heart: an exploration of labeled water permeability. *Circ. Res* 1977;41:515–533. [PubMed: 902358]
13. Kelly PJ. Comparison of marrow and cortical bone blood flow by ^{125}I -labeled 4-iodoantipyrine (I-Ap) washout. *J. Lab. Clin. Med* 1973;81:497–505. [PubMed: 4696183]
14. Effros RM, Chinard FP. The in vivo pH of the extravascular space of the lung. *J. Clin. Invest* 1969;48:1983–1996. [PubMed: 4898722]
15. Lopez-Curto JA, Bassingthwaite JB, Kelly PJ. Anatomy of the microvasculature of adult dog tibial diaphysis. *J. Bone Joint Surg. [Am]*. (in press).
16. Morris MA, Kelly PJ, Bassingthwaite JB. Fluid spaces in canine cortical bone. *Fed. Proc* 1978;37:315. (abst.).
17. Lindbjerg IF. Disappearance rate of $^{133}\text{Xenon}$, 4-iodo-antipyrine- ^{131}I and $^{131}\text{I}^-$ from human skeletal muscles and adipose tissue. *Scand. J. Clin. Lab. Invest* 1967;19:120–128.
18. Munck O, Andersen AM, Binder C. Clearance of 4-iodoantipyrine-125-I after subcutaneous injection in various regions. *Scand. J. Clin. Invest. [Suppl. 99]* 1967;19:39–45.
19. Renkin EM. Capillary and cellular permeability to some compounds related to antipyrine. *Am. J. Physiol* 1953;173:125–130. [PubMed: 13050774]
20. Lopez-Curto JA, Hughes SPF, Kelly PJ, Bassingthwaite JB. Interstitial fluid space in bone. *Fed. Proc* 1976;35:850. (abst.).
21. Hughes SPF, Lemon GJ, Davies DR, Bassingthwaite JB, Kelly PJ. Extraction of minerals after experimental fractures of the tibia in dogs. *J. Bone Joint Surg. [Am]* 1979;61:857–866.
22. Morris MA, Day JB, Bassingthwaite JB, An K-N, Kelly PJ. Potassium and fluid spaces of canine cortical bone. *Physiologist* 1979 August;22:90. (abst.).
23. Kelly, PJ.; Weinman, DT.; Owen, CA, Jr. Determination of blood flow in bone with particular reference to the use of bone-seeking isotopes; Proceedings of the Conference on Aseptic Necrosis of the Femoral Head; St. Louis: National Institutes of Health, United States Public Health Service; 1964. p. 129-142.
24. Pasternak HS, Kelly PJ, Owen CA Jr. Metabolic rate in tibia of normal and thyrotoxic dogs. *J. Lab. Clin. Med* 1968;71:240–246.
25. Owen M, Triffitt JT. Extravascular albumin in bone tissue. *J. Physiol. (Lond.)* 1976;257:293–307. [PubMed: 950595]
26. Goresky CA, Bach GG, Nadeau BE. Red cell carriage of label: its limiting effect on the exchange of materials in the liver. *Circ. Res* 1975;36:328–351. [PubMed: 1090390]
27. Tosteson DC. Halide transport in red blood cells. *Acta Physiol. Scand* 1959;46:19–41.
28. Widdowson, EM.; Dickerson, JWT. Chemical composition of the body. In: Comar, CL.; Bronner, F., editors. *Mineral Metabolism: An Advanced Treatise*. Vol. Vol. 2. New York: Academic Press; 1964. p. 140
29. Dalmark M. Effects of halides and bicarbonate on chloride transport in human red blood cells. *J. Gen. Physiol* 1976;67:223–234. [PubMed: 1255128]
30. Bosch WJ. Plasma ^{45}Ca clearance by the tibia in the immature dog. *Am. J. Physiol* 1969;216:1150–1157. [PubMed: 5797296]
31. Morris MA, Kelly PJ. Use of tracer microspheres to measure bone blood flow in conscious dogs. *Calcif. Tissue Int.* (in press).
32. Foreman, H. The pharmacology of some useful chelating agents. In: Seven, MJ., editor. *Metal-Binding in Medicine*. Philadelphia: J.B. Lippincott Company; 1960. p. 82-94.
33. Pecsok RL, Shields LD, Schaefer WP. Complexes of chromium(II) and (III) with ethylenediamine-tetraacetic acid. *Inorganic Chem* 1964;3:114–116.
34. Schwarzenbach G, Gut R, Anderegg G. Komplexe XXV: Die polarographische Untersuchung von Austauschgleichgewichten; Neue Daten der Bildungskonstanten von Metallkomplexen der

Äthylendiamin-tetraessigsäure und der 1,2-Diaminocyclohexan-tetraessigsäure. *Helv. Chim. Acta* 1954;37:937–957.

35. Schwarzenbach G, Ackermann H, Komplexone V. Die Äthylendiamin-tetraessigsäure. *Helv. Chim. Acta* 1947;30:1798–1804.
36. Wootton R, Reeve J, Veall N. The clinical measurement of skeletal blood flow. *Clin. Sci. Mol. Med* 1976;50:261–268. [PubMed: 1261207]

Fig. 1.
Outflow dilution curves of $^{18}\text{F}^-$, albumin- ^{51}Cr , and sucrose- ^{14}C injected in saline into nutrient artery of adult dog tibia (*upper panel*) and extraction curves for sucrose and fluoride (*lower panel*)

Fig. 2.

Outflow dilution in dog H-236, run 4, $h(t)$ (upper panel), and extraction curves, $E(t)$ (lower panel), for albumin, fluoride, and strontium in pump-perfused dog tibial diaphysis. Impulse injection was made into nutrient artery. Curves of $E(t)$ slope upward from $t = 15$ to $t = 40$ s and then diminish. Extraction continues longer for F^- than for Sr^{2+} ; $E_{\text{Sr}}(t)$ crosses zero, or $h_{\text{Sr}}(t)$ becomes greater than $h_{\text{alb}}(t)$, at $1\frac{1}{2}$ min, whereas $E_{\text{F}}(t)$ remains positive for 3 min. E_{max} is taken near peak of reference curve; $E_{\text{F}} = 0.41$, $E_{\text{Sr}} = 0.36$

Fig. 3.
Outflow dilution curves of $^{18}\text{F}^-$, albumin- ^{51}Cr , and antipyrine- ^{14}C (upper panel).
Instantaneous extraction of antipyrine- ^{14}C is considerably higher than that of ^{18}F but is still less than unity (lower panel). There is evidence of recirculation at end of curve

Fig. 4.

Data from experiments in sets 4 and 5, Table 3, E_F or E_{Sr} . Instantaneous extraction near peak of reference curve is plotted against flow (*upper panel*) and against net extraction (*lower panel*). Extraction decreases with flow

Fig. 5.
Data from experimental set 6. Instantaneous and net extraction are plotted against flow

Table 1

Experimental methods

Experimental set	Tracers(dose)	No. of dogs	No. of dilution curves	Objectives
1 ^a	¹⁸ F ⁻ ; Sucrose- ¹⁴ C (15 μCi) (5 μCi)	8	8	Comparative extractions of ¹⁸ F ⁻ and sucrose- ¹⁴ C
2 ^a	¹⁸ F ⁻ ; Sucrose- ¹⁴ C (15 μCi) (5 μCi)	3	6	Extraction when ¹⁸ F ⁻ is pre-equilibrated in blood
3 ^a	¹⁸ F ⁻ ; Antipyrine- ¹⁴ C (15 μCi) (5 μCi)	8	8	Comparison of freely diffusible molecule (antipyrine) with hydrophilic ion (fluorine)
4 ^b	⁸⁵ Sr ²⁺ (5 μCi)	7	7	Comparison of extraction to known flow
5 ^b	⁸⁵ Sr ²⁺ ; ¹⁸ F ⁻ (5 μCi) (60 μCi)	2	10	Comparison of extraction of Sr ²⁺ with that of F ⁻ with known flow
6 ^b	EDTA- ⁵¹ Cr; Sucrose- ¹⁴ C (20 μCi) (10 μCi)	3	6	Comparison of extraction of two molecules with similar molecular weight but different volume of distribution

^a Autoperfused (collateral flow), flow not measured; reference tracer, albumin-⁵¹Cr (15 μCi)

^b Perfusion of capillary bed by Holter pump, flow measured; reference tracer, albumin-^{99m}Tc (15 μCi)

Table 2

Influence of pre-equilibration with erythrocytes

Experiment no.	Injectate	E_{max}	
		^{18}F	Suc
1	Saline	0.53	0.35
	Blood	0.63	0.47
2	Saline	0.62	0.31
	Blood	0.61	0.31
3	Saline	0.73	0.33
	Blood	0.62	0.26

Table 3

Extractions for $^{85}\text{Sr}^{2+}$ and for $^{85}\text{Sr}^{2+}$ and $^{18}\text{F}^-$ injected simultaneously

Experiment	Hematocrit, %	F_{ss} , ml/100 g/min	E_F $^{18}\text{F}^-$	E_{Sr} $^{85}\text{Sr}^{2+}$	E_{net} (2 min), $^{18}\text{F}^-$ $^{85}\text{Sr}^{2+}$
G783	0.43	0.65		0.68	0.68
G705	0.44	0.92		0.61	0.45
G662	0.38	1.46		0.35	0.35
G794	0.50	1.69		0.54	0.42
G862	0.35	1.03		0.52	0.52
G816	0.40	1.07		0.75	0.44
H 80	0.44	1.03		0.66	0.57
H159 1	0	3.46	0.42	0.42	0.30 0.35
2	0	3.25	0.18	0.18	0.15 0.18
3	20	1.73	0.28	0.28	0.18 0.22
4	0	2.85	0.42	0.42	0.36 0.35
5	20	1.25	0.35	0.35	0.29 0.28
H236 1	0	2.81	0.52	0.52	0.43 0.49
2	0	2.66	0.31	0.32	0.27 0.31
3	20	1.68	0.31	0.31	0.30 0.30
4	0	3.44	0.41	0.41	0.32 0.36
5	20	1.20	0.29	0.29	0.30 0.29

Table 4

Extractions for EDTA-⁵¹Cr and sucrose-¹⁴C

Experiment	Hematocrit, %	F_{ss} , ml/100 g/min	E_{EDTA}	$E_{sucrose}$	E_{net} (2 min)	
					EDTA	Sucrose
G983 1	0.20	1.62	0.14	0.07	0.13	0.03
2	0.06	1.52	0.22	0.08	0.17	0.01
G843 1	0.36	0.82	0.33	0.26	0.27	0.20
2	0.36	0.79	0.30	0.20	0.29	0.19
G799 1	0.38	1.06	0.26	0.17	0.31	0.22
2	0.09	1.56	0.23	0.11	0.24	0.15