Cardiologists' Charting Varied by Risk Factor, and was Often Discordant with Patient Report

Shannon Gravely-Witte, MSc¹, Donna E. Stewart, MD, FRCPC^{1,2}, Neville Suskin, MBChB, MSc, FRCPC, FACC³, Lyall Higginson, MD, FRCPC⁴, David A. Alter, MD, FRCPC^{5,6,7}, and Sherry L. Grace, PhD^{1,8}

¹ University Health Network Women's Health Program

² The University of Toronto

³ London Health Sciences Centre and University of Western Ontario

⁴ University of Ottawa Heart Institute

⁵ Institute for Clinical and Evaluative Sciences

⁶ St. Michael's Hospital

⁷ Toronto Rehabilitation Institute, Toronto, ON

⁸ York University

Abstract

Objective—To assess the completeness of cardiac risk factor documentation by cardiologists, and agreement with patient report.

Study Design and Setting—A total of 68 Ontario cardiologists and 789 of their ambulatory cardiology patients were randomly selected. Cardiac risk factor data were systematically extracted from medical charts, and a survey was mailed to participants to assess risk factor concordance.

Results—With regard to completeness of risk factor documentation, 90.4% of charts contained a report of hypertension, 87.2% of diabetes, 80.5% of dyslipidemia, 78.6% of smoking behavior, 73.0% of other comorbidities, 48.7% of family history of heart disease, and 45.9% of body mass index or obesity. Using Cohen's κ , there was a concordance of 87.7% between physician charts and patient self-report of diabetes, 69.5% for obesity, 56.8% for smoking status, 49% for hypertension and 48.4% for family history.

Conclusion—Two of 4 major cardiac risk factors (hypertension and diabetes) were recorded in 90% of patient records, however arguably the most important reversible risk factors for cardiac disease (dyslipidemia and smoking) were only reported 80% of the time. The results suggest that physician chart report may not be the criterion standard for quality assessment in cardiac risk factor reporting.

Keywords

Medical charts; risk factors; cardiologist; concordance; completeness

What is New?

Key Findings

1. Less than 20% of the outpatient charts completely denoted all major cardiac risk factors. Most frequently, charts were missing 1 risk factor.
2. Modifiable risk factors for cardiac disease such as dyslipidemia and smoking were reported 80% of the time.
3. Agreement between chart and patient report of cardiac risk factors ranged from substantial to moderate, while concordance for comorbid conditions was poor.

What this paper adds: Neither the medical record nor patient report were necessarily the “gold standard” for risk factor documentation, and each source has distinct advantages and disadvantages for specific risk factors.

Implications: Initiatives such as electronic patient records and standardized reports should be explored as avenues to improve chart reporting, and potentially patient risk factor management.

Medical records are customarily used as the criterion standard to assess quality of care in the healthcare setting. Accurate and complete medical record documentation by physicians is essential to ensure appropriate treatment and optimal continuity of care. Missing information in medical charts can lead to medication errors, poorer quality patient management, and may have a negative effect on patient outcomes [1]. The poor quality of patient health records has been repeatedly documented in hospital settings [1–4]. An alternative to medical records as a quality assessment tool, patient self-report surveys are increasingly being used, and shown to be valid and accurate [5–9]. It is important in both clinical practice and research to identify patient treatment plans and clinical history; however, it is often difficult to obtain a complete and accurate patient profile using one data source alone as the standard.

To our knowledge, there is no multi-site study that has examined the quality of physician charting and patient self-report of cardiac risk factors in a large sample of ambulatory cardiac outpatients. Given that cardiovascular disease is the leading cause of death in the developed world, and there are major reversible risk factors that are directly related to atherosclerotic disease progression, total risk assessment is essential in order to ensure better patient care, improve disease prognosis and outcomes, and to aid in the informed decision-making process. The current study aimed to assess the completeness of reporting of cardiac risk factors within cardiologists’ outpatient charts, and concordance with patient report of diabetes, hypertension, smoking history, family history, and obesity. Patient and physician characteristics related to degree of chart completeness were also examined.

Methods

Design and Procedure

This study represents a cross-sectional component of a larger longitudinal observational study on access to cardiac rehabilitation. Upon receiving ethics approval from participating institutions, a sample of Ontario-based cardiologists was generated through a national physician registry, CMD Online, and basic sociodemographic data were extracted. Consent to participate was solicited via mail, and included a brief survey. Subsequently, the research assistant performed on-site screening on a retrospective, sample of 20 of the cardiologists' most recent patients with coronary artery disease (CAD).

With informed patient consent, clinical and risk factor data was recorded from charts, and patients were mailed a self-report survey assessing cardiac risk factors. Patient and chart report data were entered by different research assistants in order to minimize bias.

Chart Extraction

Charts of patients that had been seen by the cardiologist in the outpatient clinic between 2004 and 2006 were eligible for review. After patient consent, demographic data, cardiac risk factors (e.g., diabetes, hypertension, smoking status, family history, obesity, and dyslipidemia), cardiac medications, and disease severity indicators were extracted from charts using a standardized form. Following training, the charts were systematically reviewed by the first author prior to patient report of risk factors (i.e., blind). The complete medical record of every patient was reviewed in order to obtain a comprehensive overview of the medical history and current status. Chart extraction was completed between May 2005 and September 2006.

Participants

Sixty-eight non-pediatric Ontario cardiologists consented to participate, and their characteristics are shown in Table 1. A retrospective sample of 1376 CAD outpatients were mailed to ask for their consent to participate in this study. CAD diagnosis was confirmed based on indication in patient chart of detailed history, focused physical examination, diagnostic ECG changes (i.e., Q waves, and/or ST-T segment changes), troponin levels above the 99th percentile of normal and/or receiving revascularization such as a percutaneous coronary intervention (PCI) or acute coronary bypass (ACB). Patients who had concurrent valve repair/replacement or arrhythmia, or received a diagnosis of heart failure were also eligible. Reasons for ineligibility were based on exclusion criteria for the larger study as follows: lack of English language proficiency (n = 87; 33.5%), inaccurate/outdated contact information (n = 62; 23.8%), orthopedic, neuromuscular, cognitive or vision impairment which would preclude cardiac rehabilitation participation (n = 31; 11.9%), unconfirmed CAD diagnosis (n = 26; 10.0%), index event or treatment prior to 2004 (n = 17; 6.5%), death (n = 14, 5.4%), residence outside the province of Ontario (n = 8, 3.1%), ineligibility for cardiac rehabilitation based on Canadian guidelines [10] (n = 7; 2.7%), previous participation in cardiac rehabilitation (n = 5; 1.9%) and comorbid non-affective psychiatric disorders (n = 3, 1.2%).

Measurements

Clinical and risk factor data extracted from outpatient medical charts included sex, age, date of last outpatient visit, cardiac event or procedure, Canadian Cardiovascular Society (CCS) angina class [10], New York Heart Association class (NYHA) class [11], blood pressure, lipids, smoking status (i.e. current smoker or non-smoker [inclusive of those that quit smoking]), family history of CAD, body mass index or obesity status, and comorbid conditions including diabetes. Hypertension and dyslipidemic status were recorded, and where values were presented cut-off values of 140/90 for systolic and diastolic blood pressure and lipid levels above reference ranges [10, 12] were used.

In the self-report survey, patients were asked through forced-choice options to report if they had been diagnosed with hypertension, diabetes, whether they had a family history of CAD, and their current smoking status. Body mass index (BMI) was computed based on self-reported height and weight, with overweight/obesity defined as a BMI (kg/m^2) > 25 [10, 13]. Patients also completed the Duke Activity Status Index (DASI) [14] to determine functional capacity.

Statistical Analyses

During data entry in SPSS 14.02, all missing fields were coded as such. To assess completeness of outpatient charts, frequencies were used to enumerate missing risk factor data. Using one-way ANOVA, t-tests and chi-square as appropriate, differences in the number of risk factors missing were compared based on patient characteristics. Differences in risk factor charting based on cardiologist characteristics were analyzed using linear mixed modelling with a random intercept for physician in R [15, 16].

With regard to concordance, Cohen's κ was used to assess agreement of risk factor information between physician charts and patient self-report. For the Cohen's κ statistic, the classification system suggested by Fleiss (1973) [17, 18] was applied, where κ less than 0.40 represents poor to fair agreement, 0.40 to 0.60 represents moderate agreement, 0.60 to 0.8 represents substantial agreement, and 0.80 to 1.00 represents almost perfect agreement. Sensitivity, specificity, positive predictive values (PPV), and negative predictive values (NPV) were computed using the medical record review as the criterion standard.

Results

A total of 789 patients agreed to participate ($789/(1376-260) = 71\%$ response rate). This represents a mean of 11.6 patients per cardiologist, with a range from 6 to 18. Participating patient characteristics are shown in Table 2. The mean number of days between last outpatient visit or index event and the date the survey was completed was 208.11 ± 143.17 (approximately 7 months).

Chart Completeness

With regard to completeness of risk factor documentation, Table 3 depicts the risk factor, clinical and disease severity information that was denoted in patient charts. Of diabetes, hypertension, dyslipidemia, obesity, family history and smoking, charts were missing a

mean of 1.64 ± 1.35 risk factors. Figure 1 presents the number of missing risk factors in each chart. One hundred and fifty-three (19.4%) charts were complete with all risk factors denoted. Most frequently ($n=270$; 34.2%) charts had 1 risk factor missing. Over three-quarters ($n=625$; 79.2%) of the charts had less than 3 risk factors missing. Data were split by cardiologist. The lowest mean number of missing risk factors was 0.32 ± 0.48 , and the highest was 3.9 ± 1.6 .

Cardiologist and patient correlates of mean number of missing risk factors were computed. With regard to the former, there was no significant relationship between number of missing risk factors and cardiologist sex, subspecialty, graduation year, location of medical school, and whether they held a university appointment. However, cardiologists who had lower self-reported weekly volume of patients had significantly less missing data in their charts ($F=6.23$, $p=.01$ in adjusted hierarchical regression). With regard to patient characteristics, there was no significant difference in the number of missing risk factors in the charts based on sex, whether or not they had a previous ACB, and the total number of self-reported comorbid conditions ($p > .05$). Based on Pearson's correlation, there was significantly less missing risk factor data for older patients ($p < .001$), and those with lower activity status ($p=.02$).

Chart Accuracy

Table 4 reports the concordance between physician chart-report and patient self-report of cardiac risk factors and comorbid conditions. According to the classification system, there was almost perfect agreement between the two sources for diabetes, and substantial agreement for obesity. There was moderate agreement for smoking status, hypertension and family history. There was poor concordance for the presence of one or more comorbid conditions (illness other than diabetes).

Discussion

Independent risk factors for the development and prognosis of CAD are well established. Total risk assessment and subsequent treatment is essential to optimize disease prognosis [19]. Treatment-to-target can only be achieved through charting risk factors to monitor response to therapies.

Chart Completeness

In the current study, established cardiac risk factors were fairly consistently recorded in cardiologists' outpatient charts. Overall, one-fifth of charts had more than two key risk factors missing. Results revealed that two major cardiac risk factors, hypertension and diabetes, were recorded in 90% of patient records. However, arguably important reversible risk factors for cardiac disease, dyslipidemia and smoking, were only reported 80% of the time. Notation of family history and overweight/obesity were reported in less than 50% of patient charts. Finally, CCS Angina and NYHA Class were reported in approximately 30% and 20% of charts respectively. In addition to cardiac risk factor profiles, clinical histories of other diseases (comorbid conditions) were reported in 73% of the outpatient charts. These

results suggest that physician chart report may not be the criterion standard for quality assessment in cardiac risk factor reporting and clinical history assessment.

Reasons for under-reporting of some data in the charts may be somewhat explicable. For instance, CCS and NYHA class may have been less consistently reported if the patients were stable and asymptomatic at the time of assessment. Family history and obesity may have been less frequently reported as they are not, or are less, amenable to treatment respectively [20, 21]. With regard to serum cholesterol tests, physicians often do not copy results to other providers [22], thus leading to over or under-testing and chart inconsistency. Finally, smoking status is paramount to patient assessment, and initiatives should be taken to ensure higher rates of chart notation.

With regard to correlates of chart completeness, perhaps fittingly there was less missing data where patients were older and had lower activity status. This indicates that cardiologists document more completely with patients for whom there likely are more risk factors and greater overall risk. Secondly, cardiologists with higher patient volumes had lower risk factor reporting. This suggests that time is a key commodity for complete charting. Means to facilitate charting within busy practices should be explored and exploited, including use of electronic records, standardized forms and office support.

Chart Accuracy

It remains undetermined in the literature as to which tool, chart or self-report, is the criterion standard to estimate risk factor burden. Some studies have found the medical record to be more accurate than patient self-report [23, 24] while other studies have found the medical record to be unreliable [25, 26]. Further studies have shown that one source may be better at indicating certain conditions than others [27]. In this study, there was large variation in concordance between patient surveys and chart-report of cardiac risk factors and comorbid conditions. These findings raise key methodological challenges considering self-report data is generally perceived as inferior to chart report, and indeed the latter is often utilized to verify self-report data. Overall, there was significant under-reporting or over-reporting of traditional cardiovascular risk factors in both the medical record and patient self-report despite the fact that both methods utilize primary data collection. Reasons for discordance between outpatient charts and patient report likely vary by risk factor.

Similar to much of the existing literature, the results showed almost perfect agreement between patient self-report and the medical chart for diabetes. The literature shows that there is very good agreement for diabetes in various patient populations [6–8, 24, 27]. The excellent agreement between the two sources can be explained by the fact that diabetes mellitus is a chronic condition requiring repeated healthcare visits and has concrete diagnostic criteria. PPV and specificity values showed that where there was slight discordance such that patients under-reported diabetes compared to their medical charts. This could potentially occur if the physician minimized the results of borderline blood glucose tests. Secondly, if patients require only dietary changes rather than pharmacological management for diabetes, they may be unaware of their diagnosis.

There was substantial agreement for overweight/obesity. However, considering that notation of overweight/obesity was missing from over 50% of the charts, this rate of agreement may be inflated.

Smoking cessation can be considered the most effective lifestyle modification in the management of patients with CAD [28] as it can significantly reduce the risk of cardiovascular events [29, 30]. The results in this study found only moderate agreement between patient and chart report for smoking status. PPV and sensitivity values showed medical charts under-reported smoking status compared to self-report. It is concerning that there is a lack of agreement for such an important modifiable risk factor, and that the root of the problem may stem from patients' reluctance to discuss their smoking status with their cardiologist. It is important for not only cardiologists, but also all health care professionals to open the lines of communication to discuss smoking status and subsequently promote smoking cessation.

Hypertension is an asymptomatic condition, which may result in low awareness among patients. The preponderance of literature shows good agreement for hypertension between the medical record with patient self-report [7, 8, 27, 31], however this study revealed only moderate agreement. The hypertension PPV value showed that patients reported hypertension less often than what was recorded in their medical chart. Some studies within the literature support our findings [6, 24]. Reasons for discordance include patient perception that they no longer have hypertension if it is controlled pharmacologically, or the changing thresholds for defining hypertension. Cardiologists must communicate blood pressure status to their patients to support risk-reducing behaviours such as medication adherence and sodium reduction.

Moderate agreement was also found for family history of CAD. Finally, with regard to comorbid conditions, there was poor agreement between charts and patient report. Based on the PPV value, patients reported more health problems than were recorded in their charts. Previous literature has found that patients and physicians may differ in their report on comorbid illnesses [27] and that there tends to be a wide variation in agreement between questionnaires and medical records by specific diagnosis. Patients in our study may have more often reported minor illnesses (i.e. muscle/joint pain or allergies), while cardiologists more often focused on serious conditions (i.e. cancer or neurological problems). Moreover, comorbid conditions which are diagnostically complex or require clinical judgment in addition to laboratory testing may be more vulnerable to error in self-report [32]. This may have partly accounted for the high discordance.

There are other potential explanations for chart and patient-report discordance which are not specific to any risk factor. For example, discordance could reflect lack of assessment of a risk factor or condition by the cardiologist, while the patient is aware of their status through communication from another physician. The intermittent nature of diabetes symptoms or those from other comorbid conditions may also affect the awareness and accuracy of patient self-report. Moreover, risk factors and chronic conditions which have diagnostic values that fluctuate over time such as dyslipidemia and hypertension, may not be clearly communicated to the patient, therefore yielding lower agreement rates [31]. Finally,

discordance could be due to lack of communication of risk factor status to the patient, or perhaps the information was reported to the patient but the patient did not comprehend or retain the information. Future research should be conducted to explore what factors account for risk factor discordance.

Based on these results and those in the literature, it may be implied that neither chart nor patient report can serve as the criterion standard as also suggested by Dendukuri et al. [23]. Arguably both patient self-report and medical records should be used in combination to produce a complete and valid patient profile. Indeed, depending upon one's purposes, both forms of report have distinct advantages and disadvantages. For example, to determine whether a patient meets diagnostic criteria for a given risk factor or condition, the medical chart would be the criterion standard. However, for an exhaustive list of comorbid conditions or for smoking status, the patient would likely be the preferred source.

Caution is warranted when interpreting results. First, this study did not incorporate a second extractor to examine a random subsample of charts for coding consistency. However, as only one RA examined the patient charts across all sites, reporting should be highly consistent. Second, where we reported a lack of agreement in risk factor reporting, this discordance between physician chart and patient report may be due not to poor quality charting but to lack of patient awareness or the failure to disclose the risk factor to the patient. Therefore the accuracy of risk factor charting data should be interpreted with caution.

In conclusion, cardiac risk factors are consistently reported in outpatient charts, and most risk factors charted were at least moderately concordant with patient report. However, neither the medical record nor patient report were necessarily the "gold standard" for risk factor documentation, and indeed each source has distinct advantages and disadvantages for specific risk factors. Initiatives such as electronic patient records and standardized reports should be explored as avenues to improve chart reporting, and potentially patient risk factor management.

Acknowledgments

This research is funded by the Canadian Institutes of Health Research, grant # MOP-74431. Dr. Grace is supported by the Ontario Ministry of Health and Long-Term Care. We also acknowledge Sheena Kayaniyl for data entry. We are thankful to the Editor for thoughtful comments on this paper.

References

1. Ministry of Health and Long-Term Care. physician documentation expert panel. 2006. A guide to better physician documentation.
2. Roukema J, Los RK, Bleeker SE, van Ginneken AM, van der Lei J, Moll HA. Paper versus computer: Feasibility of an electronic medical record in general pediatrics. *Pediatrics*. 2006; 117:15–21. [PubMed: 16396855]
3. Martin CA. Improving the quality of medical record documentation. *J Healthc Qual*. 1992; 14:16–23.
4. Ashkenazi A, Cooper M, Chemke J, et al. The quality assurance committee in a general hospital: Its use in improvement of the medical record. *Isr J Med Sci*. 1992; 28:714–717. [PubMed: 1399501]

5. Schootman M, Jeffe DB, West MM, Aft R. Self-report by elderly breast cancer patients was an acceptable alternative to surveillance, epidemiology, and end results (SEER) abstract data. *J Clin Epidemiol.* 2005; 58:1316–1319. [PubMed: 16291477]
6. Bush TL, Miller SR, Golden AL, Hale WE. Self-report and medical record report agreement of selected medical conditions in the elderly. *Am J Public Health.* 1989; 79:1554–1556. [PubMed: 2817172]
7. Skinner KM, Miller DR, Lincoln E, Lee A, Kazis LE. Concordance between respondent self-reports and medical records for chronic conditions: Experience from the veterans health study. *J Ambul Care Manage.* 2005; 28:102–110. [PubMed: 15923944]
8. Okura Y, Urban LH, Mahoney DW, Jacobsen SJ, Rodeheffer RJ. Agreement between self-report questionnaires and medical record data was substantial for diabetes, hypertension, myocardial infarction and stroke but not for heart failure. *J Clin Epidemiol.* 2004; 57:1096–1103. [PubMed: 15528061]
9. Simpson CF, Boyd CM, Carlson MC, Griswold ME, Guralnik JM, Fried LP. Agreement between self-report of disease diagnoses and medical record validation in disabled older women: Factors that modify agreement. *J Am Geriatr Soc.* 2004; 52:123–127. [PubMed: 14687326]
10. Canadian Association of Cardiac Rehabilitation. Canadian guidelines for cardiac rehabilitation and cardiovascular disease prevention: Enhancing the science, refining the art. Winnipeg, MB: CACRC; 2004.
11. The Criteria Committee of the New York Heart Association. Nomenclature and Criteria for Diagnosis of Diseases of the Heart and Great Vessels. Boston: Little, Brown & Co; 1994.
12. American Association for Clinical Chemistry. Cardiac Risk Assessment. Accessed 03/07, 2007 Available at: http://www.labtestsonline.org/understanding/analytes/cardiac_risk/glance.html
13. World Health Organization. Obesity: Preventing and managing the global epidemic -report of a WHO consultation on obesity. Geneva: 1997. WHO/NUT/NCD/98.1
14. Hlatky MA, Boineau RE, Higginbotham MB, et al. A brief self-administered questionnaire to determine functional capacity (the duke activity status index). *Am J Cardiol.* 1989; 64:651–654. [PubMed: 2782256]
15. R Development Core Team. R: A language and environment for statistical computing. 2006 Available at: URL <http://www.R-project.org>
16. Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. Nlme: Linear and nonlinear mixed effects models. 2006. p. 3.1-79.
17. Fleiss, JL. Statistical Methods for Rates and Proportions. New York: Wiley; 1973.
18. Landis JR, Koch GG. An application of hierarchical kappa-type statistics in the assessment of majority agreement among multiple observers. *Biometrics.* 1977; 33:363–374. [PubMed: 884196]
19. Graham IM. The importance of total cardiovascular risk assessment in clinical practice. *Eur J Gen Pract.* 2006; 12:148–155. [PubMed: 17127600]
20. Belle SH, Berk PD, Courcoulas AP, et al. Safety and efficacy of bariatric surgery: Longitudinal assessment of bariatric surgery. *Surg Obes Relat Dis.* 2007; 3:116–126. [PubMed: 17386392]
21. Thirlby RC, Randall J. A genetic “obesity risk index” for patients with morbid obesity. *Obes Surg.* 2002; 12:25–29. [PubMed: 11868292]
22. Grace SL, Krepostman S, Brooks D, et al. Referral to and discharge from cardiac rehabilitation: Key informant views on continuity of care. *J Eval Clin Pract.* 2006; 12:155–163. [PubMed: 16579824]
23. Dendukuri N, McCusker J, Bellavance F, et al. Comparing the validity of different sources of information on emergency department visits: A latent class analysis. *Med Care.* 2005; 43:266–275. [PubMed: 15725983]
24. Merkin SS, Cavanaugh K, Longenecker JC, Fink NE, Levey AS, Powe NR. Agreement of self-reported comorbid conditions with medical and physician reports varied by disease among end-stage renal disease patients. *J Clin Epidemiol.* 2007; 60:634–642. [PubMed: 17493523]
25. Luck J, Peabody JW, Dresselhaus TR, Lee M, Glassman P. How well does chart abstraction measure quality? A prospective comparison of standardized patients with the medical record. *Am J Med.* 2000; 108:642–649. [PubMed: 10856412]

26. Cradock J, Young AS, Sullivan G. The accuracy of medical record documentation in schizophrenia. *J Behav Health Serv Res*. 2001; 28:456–465. [PubMed: 11732247]
27. Harlow SD, Linet MS. Agreement between questionnaire data and medical records. the evidence for accuracy of recall. *Am J Epidemiol*. 1989; 129:233–248. [PubMed: 2643301]
28. Daly LE, Mulcahy R, Graham IM, Hickey N. Long term effect on mortality of stopping smoking after unstable angina and myocardial infarction. *Br Med J (Clin Res Ed)*. 1983; 287:324–326.
29. Critchley J, Capewell S. Smoking cessation for the secondary prevention of coronary heart disease. *Cochrane Database Syst Rev*. 2004; (1):CD003041. [PubMed: 14974003]
30. Health Canada. Smoking and Heart Disease. Accessed 10/28, 2007 Available at: http://hc-sc.gc.ca/hl-vs/tobac-tabac/body-corps/disease-maladie/heart-coeur/index_e.html
31. Haapanen N, Miilunpalo S, Pasanen M, Oja P, Vuori I. Agreement between questionnaire data and medical records of chronic diseases in middle-aged and elderly finnish men and women. *Am J Epidemiol*. 1997; 145:762–769. [PubMed: 9126003]
32. Colditz GA, Martin P, Stampfer MJ, et al. Validation of questionnaire information on risk factors and disease outcomes in a prospective cohort study of women. *Am J Epidemiol*. 1986; 123:894–900. [PubMed: 3962971]

Figure 1.
Total number of cardiac risk factors missing in outpatient charts.

Table 1

Characteristics of participating cardiologists

Characteristics	Participants (N=68)
Sex (% female)	11 (16.2%)
Graduation year – medical degree (mean $\pm$ SD)	1982 $\pm$ 8.3
Location of medical school (% Ontario)	40 (58.8%)
University appointment (% yes)	28 (42.4.0%)
Subspecialty (% internists/no subspecialty)	44 (64.7%)
Self-reported volume of patients/week (mean $\pm$ SD)	51.4 $\pm$ 33.02

Table 2

Sociodemographic and clinical characteristics of patient sample (N=789)

Characteristics	Mean $\pm$ SD or n (%)
Age	66.46 $\pm$ 11.35
Sex (% female)	224 (28.4%)
BMI [†]	27.39 $\pm$ 5.33
Marital status [†] (% married)	550 (69.7%)
Ethnocultural background [†] (% minority)	109 (13.8%)
Education [†] (% >high school)	425 (53.9%)
Family income [†] (% \$50,000CAD or more)	345 (43.7%)
Work status [†] (% retired)	422 (53.5%)
Systolic BP mm Hg (mean $\pm$ SD)	131.8 $\pm$ 20.61
Diastolic BP mm Hg (mean $\pm$ SD)	75.48 $\pm$ 10.37
Total Cholesterol/HDL Ratio	3.74 $\pm$ 1.50
HDL mmol/L (mean $\pm$ SD)	1.20 $\pm$ 0.37
LDL mmol/L (mean $\pm$ SD)	2.34 $\pm$ 0.93
NYHA Class II–IV (%)	64 (8.1%)
CCS angina class 2–4 (%)	183 (23.2%)
Multi-vessel Disease (>1 diseased coronary arteries)	238 (30.2%)
Duke Activity Status Index [†]	36.23 $\pm$ 16.12
Current or Previous MI	366 (46.4%)
Current or Previous PCI	365 (46.3%)
Current or Previous ACB	233 (29.5%)
Current or Previous HF	140 (17.7%)
Current or Previous Arrhythmia	166 (21.0%)
Current or Previous of Valve repair/replacement	121 (15.3%)

Note: Percentages take into account missing data for some variables.

BMI, Body Mass Index; BP, blood pressure; HDL, high-density lipoprotein; LDL, low-density lipoprotein; CCS, Canadian Cardiovascular Society; NYHA, New York Heart Association. MI, Myocardial Infarction; PCI, Percutaneous Coronary Intervention; ACB, Acute Coronary Bypass; HF, Heart Failure.

[†] denotes data from patient report

Table 3

Presence of cardiac risk factor data in cardiologist charts

Clinical Data	Data Present n (%)
Hypertension	713 (90.4%)
Recent BP Values	667 (84.5%)
Diabetes	688 (87.2%)
Dyslipidemia	635 (80.5%)
Lipid Values	567 (40.1%)
Smoking Status	620 (78.6%)
Comorbidities	576 (73.0%)
FHx of CAD	384 (48.7%)
BMI or Overweight/Obesity	362 (45.9%)
CCS Anginal Class	230 (29.2%)
NYHA Class	148 (18.8%)

Note: BP, blood pressure; CCS, Canadian Cardiovascular Society; NYHA, New York Heart Association; FHx, Family History; BMI, Body Mass Index.

Table 4

Agreement between patient report and medical outpatient chart

Risk Factors	Chart		Totals
	Yes	No	
Diabetes Mellitus			
Patient Survey			
Yes	167	9	176
	24.4%	1.3%	25.7%
No	24	484	508
	3.5%	70.8%	74.3%
Chart Totals	191	493	684
	27.9%	72.1%	100.0%
Sensitivity	87.4%		
Specificity	98.2%		
PPV	94.9%		
NPV	95.3%		
Cohen's κ	0.88		
Obesity			
Patient Survey			
Yes	166	28	194
	46.5%	7.8%	54.3%
No	26	137	163
	7.3%	38.4%	45.7%
Chart Totals	192	165	357
	53.8%	46.2%	100.0%
Sensitivity	86.5%		
Specificity	83.0%		
PPV	85.6%		
NPV	84.0%		
Cohen's κ	0.69		
Current Smoking Status			
Patient Survey			
Yes	61	12	73
	9.9%	2.0%	11.9%
No	59	483	542
	9.6%	78.5%	88.1%
Chart Totals	120	495	615
	19.5%	80.5%	100.0%
Sensitivity	83.6%		
Specificity	97.6%		
PPV	50.8%		
NPV	89.1%		

Risk Factors	Chart		Totals
	Yes	No	
Cohen's κ	0.57		
Hypertension			
Patient Survey			
Yes	321	40	361
	45.8%	5.7%	51.5%
No	137	203	340
	19.5%	29.0%	48.5%
Chart Totals	458	243	701
	65.3%	34.7%	100.0%
Sensitivity	70.1%		
Specificity	83.5%		
PPV	88.9%		
NPV	59.7%		
Cohen's κ	0.49		
Family History of CAD			
Patient Survey			
Yes	213	42	255
	55.8%	11.0%	66.8%
No	45	82	127
	11.8%	21.5%	33.2%
Chart Totals	258	124	382
	67.5%	32.5%	100.0%
Sensitivity	82.6%		
Specificity	66.1%		
PPV	83.5%		
NPV	64.6%		
Cohen's κ	0.48		
Co-morbid Conditions*			
Patient Survey			
Yes	361	93	454
	62.7%	16.1%	78.8%
No	73	49	122
	12.7%	8.5%	21.2%
Chart Totals	434	142	576
	75.3%	24.7%	100.0%
Sensitivity	83.2%		
Specificity	34.5%		
PPV	79.5%		
NPV	40.2%		
Cohen's κ	0.19		

Note: PPV, Positive Predictive Value; NPV, Negative Predictive Value; CAD, Coronary Heart Disease.

* Conditions other than diabetes.