Published in final edited form as:

Am J Alzheimers Dis Other Dement. 2008 ; 23(2): 125–131. doi:10.1177/1533317507307961.

PATIENT CARE AND MANAGEMENT OF FRONTOTEMPORAL LOBAR DEGENERATION

Lauren Massimo and Murray Grossman

Dept of Neurology, University of Pennsylvania School of Medicine

Abstract

Frontotemporal lobar degeneration (FTLD) is a neurodegenerative disease that affects frontal and temporal regions of the brain. Two proteins indicated in the pathology are tau and the recently discovered TDP-43. Major manifestations include progressive aphasia and a disorder of social comportment. The diagnosis of a patient includes a detailed cognitive exam, clinical testing and neuroimaging techniques. The current goal of therapy for FTLD is symptomatic management with medications borrowed from other conditions. Non-pharmacologic management such as behavioral interventions and environmental engineering are also efficacious in optimizing quality of life.

Introduction

Frontotemporal lobar degeneration (FTLD) is a neurodegenerative disease that affects frontal and temporal regions of the brain. FTLD is remarkable for its early onset, typically presenting typically in the fifth or sixth decade of life¹. Although epidemiological studies of FTLD are limited, the prevalence of FTLD is estimated to be comparable to that of Alzheimer's Disease (AD) among individuals with dementia who are younger than 65 years at onset².

Many individuals presenting with an FTLD spectrum disorder have histopathologic disease at autopsy that identifies the condition as a *tauopathy*. Tau, a microtubule-associated protein present in neurons, serves two crucial functions: maintaining the shape of the cell and transporting metabolic substances throughout the neuron. Tauopathies include conditions like dementia with Pick bodies (PiD), corticobasal degeneration (CBD), progressive supranuclear palsy (PSP) and argyrophilic grain disease (AGD)³. Other individuals with FTLD are said to have a tau-negative disease at autopsy. The most common histopathology seen in these patients is frontotemporal lobar degeneration with ubiquitin-positive immunoreactive inclusions (FTLD-U)⁴. Until recently, the ubiquitinated protein was unknown. TDP-43 is now known as the signature pathology in FTLD-U⁵, as well as amyotrophic lateral sclerosis (ALS) and FTLD accompanied by ALS. Other causes of FTLD that are tau-negative include Dementia Lacking Distinctive Histopathology (DLDH) and Neuronal Intermediate Filament Inclusion Disease (NIFID).

Approximately 60% of FTLD cases are sporadic, where there are no other family members with an FTLD spectrum disorder, and relatives do not appear to have an increased risk of developing the disease. However, the remaining 40% of cases appear to have some family history of dementia or neurodegenerative disease⁶, and up to 30% of FTLD is inherited in an autosomal dominant pattern. In a small percentage cases, FTLD is due definitively to an

identifiable mutation⁷. Mutations in the tau gene (*MAPT*) account for 7 to 50% of FTLN cases with an autosomal dominant pattern of inheritance⁸⁻¹⁴. Goldman and colleagues explain that the large variation in occurrence of genetic profiles of FTLN patients with tau mutations is due to differing criteria among centers for a positive family history⁷. Recently, mutations in familial FTLN have been discovered in a gene that codes for progranulin (PGRN). This too is located on chromosome 17, the same chromosome as the tau mutation¹⁵, although there is no known relationship between these two genes on chromosome 17. Mutations in the PGRN gene are associated with FTLN-U pathology¹⁶.

Clinically, FTLN presents with a number of distinctive phenotypes that may be linked to the specific frontal and temporal distributions of disease caused by tau-positive and tau-negative disorders^{17, 18}. For example, patients with damage to ventral and inferolateral temporal lobe structures experience profound deficits in word comprehension and object knowledge. These patients are said to have semantic dementia. In contrast, patients with significant disease burden in left frontal lobe structures experience progressively effortful speech. This is accompanied by a disorder of motor speech that includes articulatory approximations and groping as well as grammatical comprehension difficulty. These patients have a syndrome known as progressive non-fluent aphasia. Patients with damage to the right frontal and temporal lobes, by comparison, show behavioral disinhibition and other changes in personality and social comportment. This includes a disorder of social conduct involving loss of insight, decline in personal hygiene, mental rigidity, distractibility, hyperorality, and perseveration. Neary and colleagues published clinical criteria that delineate these three FTLN syndromes¹⁹.

This great diversity in phenotype makes FTLN difficult to diagnose. Recent clinical-pathological studies suggest that diagnostic accuracy in experienced clinicians ranges from 15% to about 33%²⁰⁻²². Furthermore, there is no single *in vivo* diagnostic test with sufficient specificity to discriminate FTLN from other forms of dementia, although advances are being made in terms of antemortem diagnosis (Bian & Grossman, in press). This diagnostic issue is complicated by the fact that there are also clinical features that overlap with those of more common conditions such as Alzheimer's Disease (AD), small vessel ischemic disease (SVID) and Parkinson's Disease (PD)^{23, 24}. In this paper, we briefly review FTLN from a clinical perspective and discuss treatment options that are currently available.

Diagnosis

The diagnosis of FTLN requires great attention by the physician because of the neurologic and psychiatric features that are shared with other disorders and that can mimic FTLN. For example, serum studies like B12 level, sedimentation rate, and thyroid function are important to obtain to reduce the likelihood of a metabolic cause. An MRI scan will help rule out structural disease that can masquerade as FTLN such as normal pressure hydrocephalus and small vessel ischemic disease. A lumbar puncture is useful to eliminate conditions such as lymphomatous meningitis and other non-bacterial meningitides.

A major obstacle in an accurate diagnosis is distinguishing FTLN from more common neurodegenerative conditions such as AD. A variety of biomarkers are beginning to demonstrate some utility in increasing the reliability of the diagnosis of FTLN (Bian & Grossman, in press). A detailed cognitive and behavioral assessment is an essential element of the neurological exam¹. Neuropsychological assessments are generally quite good at identifying specific patterns of impairment in these patients, but these lengthy evaluations are not very practical. Nevertheless, several studies appear to show specific patterns of neuropsychological impairment in patients with autopsy-proven FTLN^{25, 26}. Brief screening instruments like the Mini Mental State Examination (MMSE)²⁷ do not appear to

be sensitive for capturing patients with FTLT, and FTLT patients often have a normal MMSE when they first present^{28, 29}. However, several brief instruments have been developed recently that attempt to capture disease severity in non-Alzheimer's forms of dementia such as FTLT^{30, 31}. For example, the Philadelphia Brief Assessment of Cognition (PBAC) is a brief instrument that measures five domains of cognition: executive, language, visuospatial, social, and memory functioning. In recent analyses from our lab, the PBAC was useful at distinguishing patients with FTLT from AD and other neurodegenerative conditions such as corticobasal syndrome (Libon, et al., under review).

Abnormalities on imaging studies can contribute to the diagnosis of FTLT³². Each clinical syndrome of FTLT appears to manifest a relatively distinct MRI pattern. The behavioral or social form is associated with atrophy that is most evident in right frontal and temporal regions, including medial orbital frontal cortex³³. In contrast, patients with language disorders tend to have more prominent left hemisphere disease. Atrophy associated with PNFA is found in brain regions supporting speech production such as inferior frontal gyrus, insula, and superior temporal gyrus of the left hemisphere³⁴. Ventral and lateral atrophy of the anterior temporal lobe is frequently described in SD³⁵. Functional neuroimaging with positron emission tomography (PET) or single photon emission computed tomography (SPECT) can be useful if the MRI is within normal limits or the atrophy is too subtle³⁶. Perhaps the most sensitive imaging studies are longitudinal, in large part because of the increased variance in normal MRI volume and normal glucose metabolism found in healthy aging. A longitudinal PET study of 22 patients with a clinical diagnosis of frontal variant FTD showed a significantly lower glucose metabolism in the frontal cortical areas than healthy controls at baseline, for example³⁷. Follow up visits conducted 19.5 (+/-7.5) months later showed significantly lower glucose metabolism in frontal, inferior and middle temporal, and inferior parietal cortical areas as well as the caudate nuclei, the insula and thalamus. Similar findings have been reported in a longitudinal MRI study using a fully automated registration and high-dimensional normalization procedure³⁸.

Other biomarkers also may be useful in diagnosing FTLT. In one large series, CSF tau was found to be significantly low in 34% of individual FTLT patients, an effect that was never seen in AD cases³⁹. A follow-up study confirmed that CSF total tau and the ratio of tau/ β -amyloid are significantly lower in autopsy-confirmed FTLT than in AD (Bian, et al., under review).

Treatment

Medication treatments

The current goal of therapy for FTLT is symptomatic management. Unfortunately, there is no etiologically-based curative treatment available at present that will change the natural history of FTLT, although several labs are actively pursuing the development of medications targeted at the abnormally accumulating proteins responsible for the condition. Some medications may be helpful in managing the clinical manifestations of FTLT, and behavioral treatments also may contribute substantially to patient care.

Social abnormalities are very common in FTLT. Depression, anxiety and obsessive behavior are observed frequently. These symptoms are associated with decreased 5-HT_{1A} receptor activity⁴⁰. There have been a few double blind, randomized, placebo-controlled trials of serotonin-supplementing agents for treatment of FTLT. One trial suggested that patients who receive the serotonin-specific reuptake inhibitor paroxetine at a dosage of 20mg/day show improvement in behavioral symptoms after 14 months of drug therapy⁴¹. However, another trial of paroxetine failed to show any efficacy⁴². A small double-blinded, placebo

controlled study using trazodone up to 300 mg/day demonstrated an improvement in behavioral symptoms⁴³. None of these trials showed an improvement in cognition.

Agitation and psychosis are most often treated with neuroleptic drugs. There have been no studies to date that evaluate the effectiveness of antipsychotics in FTLN. Traditional neuroleptic agents – particularly high potency medications – are frequently avoided in FTLN because of the associated extrapyramidal side effects. Low potency neuroleptics, on the other hand, have significant soporific side effects that limit their usefulness. In a large multi-site, double-blind, placebo-controlled trial with AD patients, atypical antipsychotics drugs were efficacious, and their advantages offset adverse effects in the treatment of psychosis, agitation and aggression⁴⁴. However, other less well controlled studies failed to show any benefit from this class of medication⁴⁵. Clinicians and family members must consider preexisting medical conditions and known side effect profiles before beginning a regimen of medication.

Anticonvulsant therapy may be used alone or in conjunction with antipsychotics to treat aggressive behavior. Medications such as valproic acid have been shown to have efficacy in treating behavioral disturbance in various dementia populations⁴⁶, although this medication can be associated with weight gain and confusion. Carbamazepine is another useful anticonvulsant, but gait disturbance is a worrisome dose-dependent adverse reaction in an already vulnerable population.

Cholinesterase inhibitors have been approved by the FDA to treat mild to moderate Alzheimer's disease in the hope of delaying cognitive decline. Medications in this class such as donepezil (Aricept), rivastigmine (Exelon), and galantamine (Razadyne) are widely used in the treatment of Alzheimer's disease. In the AD population, this class of medications has been shown to be efficacious in slowing the decline in cognition and behavior⁴⁷. There have been few small studies that examine the usefulness of these drugs in FTLN patients. Some authors note that patients will experience a worsening of symptoms⁴⁸. Others report that patients improve their scores on behavioral rating scales⁴⁹. Neither study found improvement in cognitive performance.

Other medications may be useful in the management of specific symptoms. For example, medroxyprogesterone may help calm sexual urges. Dopaminergic supplementing agents may help treat some forms of involuntary movement that can occur in FTLN. Small doses of methylphenidate may contribute to the management of an amotivational syndrome.

Non-pharmacologic management

Non-pharmacologic treatments, such as behavioral or environmental interventions, play a crucial role in the management of FTD. Strategies reviewed below should be discussed with caregivers throughout the course of the illness.

Behavioral symptoms can be quite distressful for loved ones and caregivers of patients. Environmental strategies can help limit unwanted behaviors. For example, avoid situations that provoke the unwanted behavior. If crowded places or large family gatherings provoke confusion and agitation, expose the patient to social interactions only in a calmer and more controlled situation where there are small numbers of visitors. It is also useful to structure the environment in order to limit the burden of decision-making and minimize the confusion associated with multiple choices. This can involve development of a structured curriculum of planned activities every hour throughout the day. A daily schedule will keep the patient occupied and control the sleep-wake cycle so that the patient is not awake agitated in the evening but sleeping during the day. Control access to foods containing large amount of sugar, particularly in patients who are hyperoral.

Impairment in executive function is can also be problematic in patients. Limitations in planning and organization often lead to significant frustration for patients and caregivers alike. Likewise, apathy can be a substantial impediment to health and quality of life. Decreased attention and disinhibition can emerge as perseveration, ritualistic behavior or echolalia²³. Encourage activities while providing a plan for executing the activities. Multi-step activities that involve complex contingencies should be divided into simple steps. Make a list and cross each step off as it is preformed. Provide guidance and redirection to facilitate the performance of activities. Limiting options is not a restriction of freedom, but instead allows individuals to perceive clear choices and optimize the selection of the most desirable choice.

A variety of strategies can be implemented to optimize communicative efficacy. Strategies that promote language comprehension include: speaking in context; speaking at a slower pace; using grammatically simple sentences; speaking redundantly while using different words to articulate the same message; and using gestures. Computer assistive devices and personalized picture vocabulary cards are also helpful when expression is a problem.

Management of physical features of visuospatial functioning is important, particularly in patients with CBD. Limb apraxia is common and can produce awkward, clumsy movement even with simple tasks⁵⁰. Occupational therapy can play an important role in recommending orthotic devices and adaptive clothing to assist with activities of daily living (ADLs) such as dressing and using implements for eating. Bright colored tape can be used to mark spatial problem areas such as stairs and ovens. Remove throw rugs and toys from walkways to reduce the risk of falls. Physical therapy and passage range of motion exercises are useful to maintain joint mobility in the face of a limited range of motion and minimize pain due to contractures.

Aspiration pneumonia is a major source of morbidity and mortality. This may be related to difficulty coordinating swallowing, a reduced gag reflex, or inattention while eating. A swallowing study can help confirm swallowing difficulty. Recommended procedures include modifying the diet to include mechanically ground solid foods (e.g. porridge consistency) or soft foods like scrambled eggs, and thickened liquids that are nectar consistency. Eating should occur slowly to control shoveling food into the mouth, possibly by providing multiple small portions one at a time. Patients should not be distracted while eating. Surgical procedures that facilitate tube feeding without swallowing are often not effective because oral secretions continue to be produced that remain a challenge⁵¹.

Another source of morbidity and mortality is related to reduced mobility. In some patients, gait may be unstable. Physical therapy can be useful in providing a regimen of lower extremity exercises, and identifying devices that can assist gait stability such as a multi-pronged cane or a walker. In other patients who are apathetic, there may be limited postural adjustment. The minimal spontaneous movement can be a risk for the development of pressure sores or decubitus ulcers. Patients should be repositioned frequently during the day, and should be turned in bed in the middle of the night if they cannot role over independently themselves. Specialty beds and mattresses are also available to reduce pressure, but a regimen of turning is still indicated⁵².

As patients become increasingly dependent on others for activities of daily living such as bathing and toileting, the risk of infection increases in the urinary tract and in sacral pressure sores. A regular regimen of bathing should be instituted, and careful skin care is essential following bathing as well as toileting.

FTLD is a challenging disease. Clearly a great deal remains to be learned. Although recent advances help improve diagnostic accuracy, FTLT is often confused with other

neurodegenerative conditions. Medical management is currently aimed at symptomatic treatment and involves mostly medications borrowed from other conditions. As we come to understand more about the underlying pathology from autopsy studies, curative treatments will increasingly become the focus of further research. The combination of a judicious medication regimen, together with effective management of the environment, will optimize the patient's quality of life and minimize the risk factors associated with morbidity and mortality.

References

1. Grossman M. Frontotemporal dementia: A review. *J Int Neuropsych Soc* 2002;8:564–583.
2. Ratnavalli E, Brayne C, Dawson K, Hodges JR. The prevalence of frontotemporal dementia. *Neurology* 2002;58:1615–1621. [PubMed: 12058088]
3. Lee VMY, Goedert M, Trojanowski JQ. Neurodegenerative tauopathies. *Annu Rev Neurosci* 2001;24:1121–1159.
4. Mackenzie IRA, Baker M, Pickering-Brown S, et al. The neuropathology of frontotemporal lobar degeneration caused by mutations in the progranulin gene. *Brain* Nov;2006 129(Pt 11):3081–3090. see comment. [PubMed: 17071926]
5. Neumann M, Sampathu DM, Kwong LK, et al. Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Science* 2006;314:130–133. [PubMed: 17023659]
6. Stevens M, van Duijn CM, Kamphorst W, et al. Familial aggregation in frontotemporal dementia. *Neurology* 1998;50(6):1541–1545. [PubMed: 9633692]
7. Goldman J, Farmer J, Van Deerlin VM, et al. Frontotemporal Dementia: Genetics and genetic counseling dilemmas. *The Neurologist* 2004;10(5):227–234. [PubMed: 15335440]
8. Rosso SM, van Swieten JC. New developments in frontotemporal dementia and parkinsonism linked to chromosome 17. *Curr Opin Neuro Aug;2002* 15(4):423–428.
9. Hutton M, Lendon CL, Rizzu P, et al. Association of missense and 5'-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature* 1998;393:702–705. [PubMed: 9641683]
10. Clark LN, Poorkaj P, Wszolek Z, et al. Pathogenic implications of mutations in the tau gene in pallido-ponto-nigral degeneration and related neurodegenerative disorders linked to chromosome 17. *Proc Natl Acad Sci Oct 27;1998* 95(22):13103–13107. [PubMed: 9789048]
11. Morris HR, Khan MN, Janssen JC, et al. The genetic and pathological classification of familial frontotemporal dementia. *Arch Neurol* 2001;58:1813–1816. [PubMed: 11708988]
12. Rizzu P, van Swieten JC, Joosse M. High prevalence of mutations in the microtubule-associated protein tau in a population study of frontotemporal dementia in the Netherlands. *Am J Hum Genet* 1999;414–421. [PubMed: 9973279]
13. Houlden H, Baker M, Adamson J, et al. Frequency of tau mutations in three series of non-Alzheimer's degenerative dementia. *Ann Neurol* 1999;46:243–248. [PubMed: 10443890]
14. Poorkaj P, Grossman M, Steinbart E, et al. Frequency of tau gene mutations in familial and sporadic cases of non-Alzheimer's dementia. *Arch Neurol* 2001;58:383–387. [PubMed: 11255441]
15. Baker M, Mackenzie IR, Pickering-Brown SM, et al. Mutations in progranulin cause tau-negative frontotemporal dementia linked to chromosome 17. *Nature* Aug 24;2006 442:916–919. print. [PubMed: 16862116]
16. Gass J, Cannon A, Mackenzie IR, et al. Mutations in progranulin are a major cause of ubiquitin-positive frontotemporal lobar degeneration. *Hum Mol Genet Oct 15;2006* 15:2988–3001. [PubMed: 16950801]
17. Grossman M, Lee C, Morris J, Stern MB, Hurtig HI. Assessing resource demands during sentence processing in Parkinson's disease. *Brain Lang* 2002;80:603–616. [PubMed: 11896660]
18. Snowden JS, Neary D, Mann DM. *Fronto-temporal Lobar Degeneration: Fronto-temporal Dementia, Progressive Aphasia, Semantic Dementia*. New York: Churchill Livingstone; 1996.
19. Neary D, Snowden JS, Gustafson L, et al. Frontotemporal lobar degeneration: A consensus on clinical diagnostic criteria. *Neurology* 1998;51:1546–1554. [PubMed: 9855500]

20. Forman MS, Farmer J, Johnson JK, et al. Frontotemporal dementia: Clinicopathological correlations. *Ann Neurol* 2006;59:952–962. [PubMed: 16718704]
21. Knopman DS, Boeve BF, Parisi JE, et al. Antemortem diagnosis of frontotemporal lobar degeneration. *Ann Neurol* 2004;57:480–488. [PubMed: 15786453]
22. Hodges JR, Davies RR, Xuereb J, et al. Clinicopathological correlates in frontotemporal dementia. *Ann Neurol* 2004;56:399–406. [PubMed: 15349867]
23. Farmer J, Grossman M. Frontotemporal Dementia: An overview. *Alzheimer's Care Quarterly* 2005;6(3):225–232.
24. Johnson JK, Head E, Kim R, Starr A, Cotman CW. Clinical and pathological evidence for a frontal variant of Alzheimer disease. *Arch Neurol* 1999;56:1233–1239. [PubMed: 10520939]
25. Rascovsky K, Salmon DP, Ho GJ, et al. Cognitive profiles differ in autopsy-confirmed frontotemporal dementia and AD. *Neurology* 2002;58:1801–1808. [PubMed: 12084880]
26. Grossman M, Libon DJ, Forman MS, et al. Distinct antemortem profiles in pathologically defined patients with frontotemporal dementia. *Arch Neurol*. 2007 in press.
27. Folstein MF, Folstein SF, McHugh PR. "Mini Mental State." A practical method for grading the cognitive state of patients for the clinician. *J Psych Res* 1975;12:189–198.
28. Gregory CA, Hodges JR. Frontotemporal dementia: Use of consensus criteria and prevalence of psychiatric features. *Neuropsychiatry Neuropsychol Behav Neurol* 1996;9:145–153.
29. Miller BL, Cummings JL, Villanueva-Meyer J. Frontal lobe degeneration: clinical, neuropsychological, and SPECT characteristics. *Neurology* 1991;41:1374–1382. [PubMed: 1891084]
30. Dubois B, Slachevsky A, Litvan I, Pillon B. The FAB: A frontal assessment battery at bedside. *Neurology* Dec 12;2000 55:1621–1626. [PubMed: 11113214]
31. Mathuranath PS, Nestor PJ, Berrios GE, Rakowicz WP, Hodges JR. A brief cognitive test battery to differentiate Alzheimer's disease and frontotemporal dementia. *Neurology* 2000;55:1613–1620. [PubMed: 11113213]
32. Knibb JA, Kipps CM, Hodges JR. Frontotemporal dementia. *Curr Opin Neuro Dec*;2006 19(6): 565–571.
33. Eslinger PJ, Moore P, Troiani V, et al. Oops! Resolving social dilemmas in frontotemporal dementia. *J Neurol Neurosurg Psychiatry*. 2007 in press.
34. Gorno-Tempini ML, Ogar JM, Brambati SM, et al. Anatomical correlates of early mutism in progressive nonfluent aphasia. *Neurology* Nov 28;2006 67(10):1849–1851. see comment. [PubMed: 16931509]
35. Mummery CJ, Patterson K, Price CJ, Hodges JR. A voxel-based morphometry study of semantic dementia: Relationship between temporal lobe atrophy and semantic memory. *Ann Neurol* 2000;47:36–45. [PubMed: 10632099]
36. McNeill R, Sare GM, Manoharan M, et al. Accuracy of single-photon emission computed tomography in differentiating frontotemporal dementia from Alzheimer's disease. *J Neurol Neurosurg Psychiatry* Apr;2007 78(4):350–355. [PubMed: 17158559]
37. Diehl-Schmid J, Grimmer T, Drzezga A, et al. Decline of cerebral glucose metabolism in frontotemporal dementia: a longitudinal 18F-FDG-PET-study. *Neurobiol Aging* Jan;2007 28(1): 42–50. [PubMed: 16448722]
38. Avants B, Grossman M, Gee JC. Frontotemporal dementia induced annualized gray matter loss using diffeomorphic morphometry. *Alzheimer Dis Assoc Disord* 2004;19:S25–S28. [PubMed: 16317254]
39. Grossman M, Farmer J, Leight S, et al. Cerebrospinal fluid profile distinguishes frontotemporal dementia from Alzheimer's disease. *Ann Neurol* 2005;57:721–729. [PubMed: 15852395]
40. Blier P, Ward NM. Is there a role for 5-HT1A agonists in the treatment of depression? *Biol Psychiatry* Feb 1;2003 53(3):193–203. [PubMed: 12559651]
41. Moretti R, Torre P, Antonello RM, Cazzato G, Bava A. Frontotemporal Dementia: Paroxetine as a Possible Treatment of Behavior Symptoms. *Eur Neurol* 2003;49:13–19. [PubMed: 12464713]

42. Deakin JB, Rahman S, Nestor PJ, Hodges JR, Sahakian BJ. Paroxetine does not improve symptoms and impairs cognition in frontotemporal dementia: a double-blind randomized controlled trial. *Psychopharmacology* Apr;2004 172:400–408. [PubMed: 14666399]
43. Lebert F, Stekke W, Hasenbroekx C, Pasquier F. Frontotemporal dementia: A randomised, controlled trial with trazodone. *Dement Geriatr Cogn Disord* 2004;17:355–359. [PubMed: 15178953]
44. Schneider LS, Tariot PN, Dagerman KS, et al. Effectiveness of Atypical Antipsychotic Drugs in Patients with Alzheimer's Disease. *N Engl J Med* Oct 12;2006 355:1525–1538. [PubMed: 17035647]
45. Caballero J, Hitchcock M, Scharre D, Beversdorf D, Nahata MC. Cognitive effects of atypical antipsychotics in patients with Alzheimer's disease and comorbid psychiatric or behavioral problems: a retrospective study. *Clin Ther* Oct;2006 28(10):1695–1700. [PubMed: 17157125]
46. Herrmann N. Valproic acid treatment of agitation in dementia. *Can J Psych* Feb;1998 43(1):69–72.
47. Doody RS, Stevens JC, Beck C, et al. Practice parameter: management of dementia (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* May 8;2001 56(9):1154–1166. [PubMed: 11342679]
48. Mendez MF, Shapira JS, McMurtry A, Licht E. Preliminary findings: behavioral worsening on donepezil in patients with frontotemporal dementia. *Am J Geriatr Psychiatry* Jan;2007 15(1):84–87. [PubMed: 17194818]
49. Moretti R, Torre P, Antonello RM, Cattaruzza T, Cazzato G, Bava A. Rivastigmine in frontotemporal dementia: an open-label study. *Drugs Aging* 2004;21(14):931–937. [PubMed: 15554751]
50. Soliveri P, Piacentini S, Girotti F. Limb apraxia in corticobasal degeneration and progressive supranuclear palsy. *Neurology* Feb 08;2005 64:448–453. [PubMed: 15699373]
51. Finucane TE, Christmas C, Travis K. Tube feeding in patients with advanced dementia: a review of the evidence. *JAMA* Oct 13;1999 282(14):1365–1370. see comment. [PubMed: 10527184]
52. Garcia AD, Thomas DR. Assessment and management of chronic pressure ulcers in the elderly. *Med Clin North Am* Sep;2006 90(5):925–944. [PubMed: 16962850]