Published in final edited form as:

Prev Med. 2008 October ; 47(4): 452–453. doi:10.1016/j.ypmed.2008.06.007.

Meta-analysis of the effects of smokefree laws on acute myocardial infarction: An update

Stanton A. Glantz

University of California, San Francisco, CA 94143-1390, USA

Stanton A. Glantz: glantz@medicine.ucsf.edu

Keywords

Tobacco smoke pollution; Secondhand smoke; Environmental tobacco smoke; Acute myocardial infarction

There have been 8 published studies (Sargent et al., 2004; Barone-Adesi et al., 2006; Bartecchi et al., 2006; Cronin et al., 2007; Juster et al., 2007; Khuder et al., 2007; Cesaroni et al., 2008; Lemstra et al. 2008) evaluating the immediate effects of smokefree policies on hospital admissions for acute myocardial infarction since the first report of such a drop in Helena, Montana (Sargent et al. 2004). This brief report updates an earlier meta-analysis of the first four studies (Dinno and Glantz, 2007).

Pooling all the available estimates of this effect using the Stata 9.2 metan procedure in a random effects meta-analysis yields a pooled estimate of an immediate 19% (95% CI 14% to 24%) reduction on AMI admission rates associated with these laws (Fig. 1).

The fact that the studies from Italy (Barone-Adesi et al., 2006; Cesaroni et al., 2008) and Ireland showed smaller drops in AMI admissions than in the American locations may reflect lower levels of compliance with the law than in the US. In Italy and Ireland, implementation of the law was associated with 64% (Valente et al., 2007) and 69% (Mulcahy et al., 2005) reductions, respectively, in levels of secondhand smoke exposure, compared to an 84% reduction in the US (Travers et al., 2004).

In addition to these 8 published studies, this analysis does not include two studies for which confidence intervals are not available: a small study of Monroe County, Indiana (Seo and Torabi, 2007) which found a significant drop in AMI and a study of Scotland (Moss, 2007) presented at a conference that reported a 17% drop, but which has not yet been published.

The fact that many studies from so many locations around the world provide consistent findings of a substantial drop in AMI's associated with the implementation of smokefree laws increases the confidence that we can have that smokefree policies have immediate and substantial benefits in terms of reducing acute myocardial infarctions.

Acknowledgments

This work was supported by the National Cancer Institute (CA-61012).

© 2008 Published by Elsevier Inc.

Conflict of interest statement

The sponsor played no role in the conduct of the research or preparation of the manuscript. The author has no conflict of interest.

References

- Barone-Adesi F, Vizzini L, Merletti F, Richiardi L. Short-term effects of Italian smoking regulation on rates of hospital admission for acute myocardial infarction. *Eur. Heart J.* 2006; 27(20):2468–2472. [PubMed: 16940340]
- Bartecchi C, Alsever RN, Nevin-Woods C, et al. Reduction in the incidence of acute myocardial infarction associated with a citywide smoking ordinance. *Circulation.* 2006; 114(14):1490–1496. [PubMed: 17000911]
- Cesaroni G, Forastiere F, Agabiti N, Valente P, Zuccaro P, Perucci CA. Effect of the Italian smoking ban on population rates of acute coronary events. *Circulation.* 2008; 117(9):1183–1188. [PubMed: 18268149]
- Cronin, E.; Kearney, P.; Kearney, P.; Sullivan, P. Impact of a national smoking ban on the rate of admissions to hospital with acute coronary syndromes. *European Society of Cardiology Congress;* 2007.
- Dinno A, Glantz S. Clean indoor air laws immediately reduce heart attacks. *Prev. Med.* 2007; 45(1):9–11. [PubMed: 17499350]
- Juster HR, Loomis BR, Hinman TM, et al. Declines in hospital admissions for acute myocardial infarction in New York state after implementation of a comprehensive smoking ban. *Am. J. Pub. Health.* 2007; 97(11):2035–2039. [PubMed: 17901438]
- Khuder SA, Milz S, Jordan T, Price J, Silvestri K, Butler P. The impact of a smoking ban on hospital admissions for coronary heart disease. *Prev. Med.* 2007; 45(1):3–8. [PubMed: 17482249]
- Lemstra M, Neudorf C, Opondo J. Implications of a public smoking ban. *Can. J. Pub. Health.* 2008; 99(1):62–65. [PubMed: 18435394]
- Moss, L. Heart attack rates drop by fifth since smoke ban introduced *The Scotsman*, 11. Edinburgh: The Scotsman; 2007 Sept.
- Mulcahy M, Evans DS, Hammond SK, Repace JL, Byrne M. Secondhand smoke exposure and risk following the Irish smoking ban: an assessment of salivary cotinine concentrations in hotel workers and air nicotine levels in bars. *Tob. Control.* 2005; 14(6):384–388. [PubMed: 16319361]
- Sargent RP, Shepard RM, Glantz SA. Reduced incidence of admissions for myocardial infarction associated with public smoking ban: before and after study. *BMJ.* 2004; 328(7446):977–980. [PubMed: 15066887]
- Seo DC, Torabi MR. Reduced admissions for acute myocardial infarction associated with a public smoking ban: matched controlled study. *J. Drug. Educ.* 2007; 37(3):217–226. [PubMed: 18047180]
- Travers M, Cummings K, Hyland A, et al. Indoor air quality in hospitality venues before and after implementation of a clean indoor air law – Western New York, 2003. *MMWR.* 2004; 53(44): 1038–1041. [PubMed: 15538318]
- Valente P, Forastiere F, Bacosi A, et al. Exposure to fine and ultrafine particles from secondhand smoke in public places before and after the smoking ban, Italy 2005. *Tob. Control.* 2007; 16(5): 312–317. [PubMed: 17897989]

Fig. 1.

Summary of studies of the effects of smokefree laws on acute myocardial infarction in Helena, Montana (Sargent et al., 2004), Pueblo, Colorado (Bartecchi et al., 2006), Piedmont, Italy (Barone-Adesi et al., 2006), Bowling Green, Ohio (Khuder et al., 2007), New York State (Juster et al., 2007), Ireland (Cronin et al., 2007), Saskatoon, Canada (Lemstra, Neudorf and Opondo 2008), and Rome, Italy (Cesaroni et al., 2008).