Published in final edited form as:

Biol Psychiatry. 2010 September 15; 68(6): 521–527. doi:10.1016/j.biopsych.2010.04.023.

Wnt2 Expression and Signaling is Increased by Different Classes of Antidepressant Treatments

Hideki Okamoto¹, Bhavya Voleti¹, Mounira Banasr, Maysa Sarhan, Vanja Duric, Matthew J. Girgenti, Ralph J. DiLeone, Samuel S. Newton, and Ronald S. Duman

Division of Molecular Psychiatry, Abraham Ribicoff Research Facilities, Connecticut Mental Health Center, Yale University School of Medicine, New Haven, Connecticut 06508

Abstract

Background—Despite recent interest in glycogen synthase kinase-3b (GSK-3b) as a target for the treatment of mood disorders, there has been very little work related to these illnesses on the upstream signaling molecules that regulate this kinase, as well as downstream targets.

Methods—Using a focused microarray approach we examined the influence of different classes of antidepressants on Wnt signaling that controls GSK-3b activity as well as the transcription factors that contribute to the actions of GSK-3b.

Results—The results demonstrate that Wnt2 is a common target of different classes of antidepressants, and also show differential regulation of Wnt-GSK-3b signaling genes. Increased expression and function of Wnt2 was confirmed by secondary measures. Moreover, using a viral vector approach we demonstrate that increased expression of Wnt2 in the hippocampus is sufficient to produce antidepressant-like behavioral actions in well-established models of depression and treatment response.

Conclusions—These findings demonstrate that Wnt2 expression and signaling is a common target of antidepressants and that increased Wnt2 is sufficient to produce antidepressant effects.

Keywords

Wnt Signaling; Depression; Antidepressants

Major depression is caused by a combination of genetic and environmental factors, and is one of the top three most widespread and debilitating illnesses world-wide (1). Currently available antidepressants require weeks or months to produce a therapeutic response, and only a third of patients experience improvement with the first drug prescribed (1). Moreover, the mechanisms underlying the therapeutic efficacy of these agents remain largely unclear. Over the past decade intracellular pathways, including the glycogen-synthase kinase-3 (GSK-3) system, have been implicated in the adaptive responses underlying mood disorder medications (2–4). Lithium, which is used for the treatment of bipolar disorder as well as depression, inhibits the activity of GSK-3 β (5). In addition, selective inhibitors of GSK-3

© 2010 Society of Biological Psychiatry. Published by Elsevier Inc. All rights reserved.

Correspondence: Ronald S. Duman, 34 Park Street, New Haven, CT 06508. ronald.duman@yale.edu.

¹These authors contributed equally and are co-first authors.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors reported no biomedical financial interests or potential conflicts pertaining to this work.

have antidepressant efficacy in behavioral models of depression, providing further evidence that this system is a viable target for depression (6,7).

Activation of Wnt signaling is one of the primary pathways leading to inhibition of GSK-3 β activity (8). Wnts are secreted glycoproteins that signal through the frizzled (Fz) receptors (9). In the mouse genome, 19 Wnts and 13 Fzs have been identified, and can activate any of the 3 major intracellular signaling pathways: the canonical or Wnt/ β -catenin, planar cell polarity (PCP), and Wnt/calcium cascades (10). Wnt signaling is crucial for normal embryonic development, as well as neuroplasticity in the adult brain (11–13). Wnt-pathways have been implicated in various physiological functions, such as cell fate determination, cell and tissue polarity (14), synaptogenesis, dendritic morphogenesis (15) and axon remodeling (16,17). Moreover, abnormal Wnt signaling has been implicated in various disorders, ranging from cancer and neurodegeneration to psychiatric illnesses (18–22).

To identify the common signaling targets of different classes of antidepressants, we employed focused microarrays (300bp probes) containing most of the known genes in the WntFz signaling pathway, as well as different classes of neurotrophic factors, growth factors, neuroplasticity-related signaling cascades, and transcription factors (23,24). The different classes of antidepressants tested include selective serotonin (5-HT) reuptake inhibitors (SSRI, citalopram or fluoxetine), a selective norepinephrine (NE) reuptake inhibitor (SNRI, atomoxetine), a dual 5-HT/NE reuptake inhibitor (venlafaxine), and chronic ECS. The results demonstrate that selected Wnt signaling related genes are regulated by different classes of antidepressants. Notably, we identify Wnt2 as a common target and determine the functional consequences of altered Wnt2 expression using a viral expression approach.

MATERIALS AND METHODS

Animals

Male Sprague-Dawley rats (160–180g for antidepressant treatment; 250–300g for adeno-associated virus (AAV) studies) (Charles River Labs, Wilmington, MA) were housed, two or three per cage, under standard illumination parameters (12h light/dark cycle) and were given free access to food and water. All procedures were in strict accordance with the National Institutes of Health *Guide for the Care and Use of Laboratory Animals* and were approved by the Yale University Animal Care and Use Committee (YACUC).

Antidepressant administration

Rats were administered (i.p.) the following agents twice daily for 5 days for subchronic treatment or for 3 weeks for chronic treatment: vehicle (distilled water), citalopram (15 mg/kg), fluoxetine (5mg/kg), venlafaxine (15mg/kg) or atomoxetine (3mg/kg). Bilateral ECS was administered via moistened pads on spring-loaded ear clip electrodes using a pulse generator (ECT Unit 57800–001; Ugo Basile, Comerio, Italy) (frequency, 100 pulses/sec; pulse width, 0.5 msec; shock duration, 0.5 sec; current, 55 mA) as previously described (25).

Microarray analysis of gene expression

Array analysis was performed as previously described (23,26) with minor modifications. Briefly, analysis was performed on a focused array containing ~3000 targets (300 bp open-reading frame PCR products) for growth factor signaling, transcription factors, kinases and genes implicated in neuropsychiatric disorders and drug action. Total RNA (5 μ g) was extracted from hippocampus using the phenol-free total RNA isolation kit, RNAqueous (Ambion, Austin, TX) and was reverse-transcribed into cDNA using oligo-dT primers containing a nucleic acid capture sequence unique for Cy3 or Cy5. Arrays were hybridized

using a 2-step protocol where vehicle and antidepressant-treated samples were hybridized overnight followed by stringent washing and then post-staining using fluorescent dendrimers (Genisphere Inc., Hatfield, PA). Slides were subsequently subjected to washes, then scanned and analyzed using GenePix Pro 6.0 software (Molecular Devices, Sunnyvale, CA, USA).

Statistical Analysis of microarray data

Microarray image files were subjected to statistical analysis as previously described with minor modifications (23,24). Files created by GenePix analysis were imported into expression analysis software GeneSpring GX 7.3.1 (Silicon Genetics, Redwood City, CA) for additional visualization and data mining. In a two-step process, raw values were initially normalized per spot to control channel values, followed by per chip normalization to positive control genes (β -tubulin and cyclophilin). Gene regulation was determined by taking the log ratio of the median experimental channel signal to the median control channel signal. Up- and down-regulated genes were defined as having an average expression ratio of >1.3 or <0.7 , respectively. Lists of regulated genes (antidepressant drug treatment versus vehicle treated) were attained by unpaired t-test. Statistical significance was considered to be $p < 0.05$ following adjustment with Benjamini and Hochberg False Discovery Rate (FDR) multiple testing correction.

Quantitative Real-Time RT-PCR

A sensitive Sybr Green-based protocol that can reliably detect two-fold differences in gene expression was used (23). Complementary DNA was synthesized from 3 μ g of total RNA, isolated from the same tissue sample used in the microarray analysis. All the PCR data were normalized to the housekeeping gene, cyclophilin. A primer set was designed with the Primer3 software (www.genome.wi.mit.edu/cgi-bin/primer/primer3.cgi).

In situ hybridization analysis

In situ hybridization was conducted on coronal brain sections (14 μ) by hybridization with the 35S-labeled Wnt2 riboprobe as previously described (23,25). Wnt2 intensity was measured by analysis of the number of grains in 20 cells per area for total of 100 cells from 5 different regions of dentate gyrus for each animal. For nonspecific binding, grains were counted in 20 cells per adjacent area of molecular layer to each analyzed area of dentate gyrus for total 100 cells. An overall average for cells and for animals was calculated.

Immunoblotting

Briefly, hippocampus was homogenized in ice-cold lysis buffer and lysates were centrifuged at 20,000 rpm for 10 min. Supernatants were resolved in SDS-polyacrylamide gels, and transferred to nitrocellulose. Blots were probed with antibodies to phospho-Ser9-GSK3 β (cat.no. 9336) and total GSK3 β (cat. No. 9332; Cell Signaling Technology, Beverly, MA). Immunoblots were developed using horseradish peroxidase-conjugated goat anti-mouse or goat anti-rabbit IgG (Bio-Rad), followed by detection with enhanced chemiluminescence.

Construction of AAV-Control and AAV-Wnt2 plasmids

pCMV-Wnt2 was generated by first ligating the Wnt2 CDS into pGEM-T Easy vector using pGEM Easy vector systems (Promega) to generate pGEM-Wnt2 plasmid. pGEM-Wnt2 was digested with SacII, filled with C and G by DNA polymerase in order to get the blunt end. SmaI-digested pCMV-MCS and SpeI-digested Wnt2 insert were ligated using Rapid DNA ligation kit (Roche) to generate pCMV-Wnt2. To generate the AAV-Wnt2 plasmid, pAAV-MCS vector, cut with NsiI, ClaI and NdeI, was ligated to pCMV-Wnt2, cut with NdeI and NsiI. The pAAV-Wnt2 plasmid was designed to coexpress DSRed and Wnt2 under the

control of two independent CMV promoters. Control plasmid (AAV-DSRed) expresses only DSRed.

Viral production and purification

The virus was generated using a triple-transfection, helper-free method, and purified with a modified version of a published protocol (27). Briefly, HEK 293 cells were cultured in five 150 × 25 mm cell culture dishes and transfected with pAAV-DSRed/Wnt2, pHelper and pAAV-RC plasmids (Stratagene) using a standard calcium phosphate method and subsequent purification was done as described previously (27).

Stereotaxic surgery and infusions

Rats were anesthetized with xylazine (6 mg/kg, i.m., Lloyd laboratories, Shenandoar, IA) and ketamine (80 mg/kg i.m., Fort Dodge Animal Health, Overland Park, KS). Bilateral viral injections were performed with coordinates −4.3 mm (anterior/posterior), −2.0 mm (lateral), and −4.2 mm (dorsal/ventral) relative to the Bregma (28). A total of 2μl of purified virus was delivered at a rate of 0.1μl/min followed by 5 min of rest. Needles were removed and the scalp incision was closed with wound clips. After behavioral testing, animals were perfused with PBS. One half of the brain was quickly frozen and 14μ sections were cut using cryostat to perform insitu hybridization for Wnt2. Other half of the brain was kept overnight in 4% paraformaldehyde and then transferred to 30% sucrose. 40μ sections were cut using a microtome for DSRed visualisation.

Learned Helplessness (LH) paradigm

The LH procedure was performed in custom built, 2-chambered shuttle boxes (Med Associates, Vermont) as previously described (29,30). Inescapable shock (IES) consisted of 60 unsignaled, randomized foot shocks at an intensity of 0.8 mA. For active avoidance testing, 24h after the IES, animals were exposed to 30 escape trials using an FR1 schedule in which a single crossing terminated the footshock. Numbers of escape were automatically scored. Results are expressed as number of escape failures observed in the 30 trial period. Two sets of animals were used for behavioral testing. Animal groups used for LH paradigm were different from the group used to perform the other behavioral tests.

Sucrose Preference test (SPT)

All behavioral testing was performed during the light cycle as previously described (29,31). Animals were habituated to a sucrose solution (1%, Sigma, St. Louis, MO) for 48h. The sucrose preference test was performed after a 4h water deprivation on the test day. For 1h, the rats were presented with two identical bottles, one filled with the sucrose solution and the other with water. Sucrose and water consumption were determined by measuring the change in volume of fluid.

Novelty Suppressed Feeding (NSF)

NSF was conducted as previously described (29). Animals were food deprived for 12h and on the test day were placed in an open field (76.5 cm × 76.5 cm × 40 cm, Plexiglas) with eight pellets of food in the center. The animals were given 8 min to approach the food and eat. The test was stopped as soon as the animal took the first bite. The latency to eat was recorded in seconds. As a control, food consumption, after the 12h food deprivation, in the home cage was quantified.

Forced swim test (FST)

FST was a 2-day paradigm in which rats were placed for 15 min in a clear cylinder with water (24±1°C, 45 cm depth). 24h after the forced swim, on the test day, rats were placed

again for 10 min in a clear cylinder with water ($24\pm1^{\circ}\text{C}$, 45 cm depth). The sessions were recorded from the side and the latency to immobility and the immobility time during the first 5 min was calculated by a blind observer. Latency to become immobile and immobility times were scored as previously described (29,31,32).

Locomotor Activity (LA)

Ambulatory locomotor activity was assessed in clear plastic boxes fitted to automated activity meters (MED Associates, St. Albans, VT) consisting of two parallel rows of photosensors (16 sensors per row, 2.5 cm apart). Locomotor activity was recorded in 5-min bins for a total of 30 min by using Med PC software.

RESULTS

Influence of different classes of antidepressants on Wnt signaling

To identify the changes in hippocampal gene expression in response to different antidepressants, we performed microarray analysis using a focused array that included multiple classes of signaling molecules. The expression of several genes involved in the Wnt signaling pathway was regulated by chronic antidepressant administration. As seen in Fig. 1a, Wnt2 was upregulated by chronic administration of each of these antidepressants tested (21d). The results also show that Wnt2 expression was increased by chronic ECS (10d). In addition, several other Wnts were upregulated, although these were not consistently increased by all antidepressants (Fig. 1A).

Wnt ligands bind to and signal through Fz receptors and can activate 3 different signaling pathways, including the Wnt/ β -catenin cascade. We found evidence that 1 of the 10 Fz subtypes, Fz9 is upregulated by both atomoxetine and venlafaxine, but not by the other 3 treatments (Fig. 1A). We also found that frizzled-related protein (sFRP3 or Frzb) is upregulated by citalopram, fluoxetine and venlafaxine, but not by atomoxetine or ECS (Fig. 1A). The sFRPs were initially found as Wnt scavengers that bind to Wnts and prevent Wnt-Fz activation, but recent studies have reported direct binding of sFRPs to Fzs resulting in activation (33). We also examined Dvl, a cytoplasmic scaffolding protein involved in Wnt signaling. Dvl1 was significantly upregulated by venlafaxine and atomoxetine, but not the other treatments tested (Fig. 1A). There were no significant effects on expression levels of Dvl2 and Dvl3 (not shown).

Wnt signaling leads to inhibition of GSK-3 β activity. In the absence of Wnt stimulation, GSK-3 β phosphorylates β -catenin and targets it for ubiquitination and proteosomal degradation. Activation of canonical Wnt signaling thereby leads to β -catenin stabilization. We found significant upregulation of β -catenin mRNA in response to citalopram and venlafaxine, but not the other antidepressants tested (Fig. 1C). We also found that expression of Akt1, another kinase that phosphorylates and inactivates GSK-3 β , is increased by venlafaxine (Fig. 1C).

β -catenin can translocate to the nucleus and regulate gene expression via the lymphoid enhancer-binding factor (LEF)/T-cell factor (TCF) family of transcription factors. Levels of TCFL1, TCF15, and LEF1 were significantly upregulated by the different antidepressant treatments (Fig. 1B). There are multiple genes that are regulated by LEF/TCF transcription factors, but recent studies have demonstrated that a crucial target for the Wnt pathway in adult neurogenesis was to activate the transcription factor, NeuroD1 (34,35). In the current study, we found that expression of NeuroD1 was significantly increased by citalopram, fluoxetine, and venlafaxine (Fig. 1B).

Secondary confirmation of Wnt2 gene expression and function

To further validate the findings obtained by microarrays, we performed quantitative RT-PCR. As seen in Fig. 2, Wnt2 was upregulated by all 4 chemical antidepressants tested, confirming the microarray findings, and our previous findings with ECS (25). To determine the effect of sub chronic antidepressant treatment on gene expression, drugs were administered with the same dose and regimen as used for chronic treatment, but for only 5d. For these and the following studies of GSK-3, we chose two representative antidepressants, fluoxetine and venlafaxine. No significant change in the average mRNA fold change with sub chronic (5d) treatment of either fluoxetine (0.92 ± 0.09) or venlafaxine (1.01 ± 0.08) was found indicating that the induction of Wnt2 expression requires chronic treatment.

Since Wnt2 was upregulated by chronic treatment of all antidepressants tested, we also performed in situ hybridization and emulsion autoradiography to determine the expression in subregions of the hippocampus. Fig. 3 shows representative photomicrographs of silver grain distribution for Wnt2 mRNA in dentate gyrus of rat hippocampus. There was a significant increase in the density of the Wnt2 mRNA in the dentate gyrus of antidepressant-treated rats (Fig. 3).

Activation of canonical Wnt signaling results in the destabilization of protein complex composed of Axin, APC, β -catenin and GSK-3 β resulting in an increase in β -catenin levels. Some studies show that the activation of the PI3 kinase/Akt pathway results in an increase in Akt1 catalytic activity which in turn phosphorylates GSK-3 β at Ser-9 in a Dvl1-mediated mechanism resulting in the inhibition of its kinase activity (36,37). Since we found an increase in both β -catenin and Akt1 in response to some of the antidepressant treatments and to determine if the regulation of Wnt signaling by chronic antidepressant treatment affects GSK-3 phosphorylation, levels of phospho-Ser 9-GSK-3 were determined by immunoblot analysis. As seen in Fig. 4, chronic administration of fluoxetine or venlafaxine significantly increased the ratio of phospho-Ser9-GSK3 β /GSK-3 β (lower band) whereas the GSK-3 β protein levels were unaffected (Fig. 4). There was a similar trend for induction of phospho-GSK-3 α (upper band), although this effect was not significant (data not shown).

Behavioral effects of viral Wnt2 expression

To determine the role of Wnt2 in behavioral models of depression/antidepressant response, an AAV vector was designed to express Wnt2 as well as the marker DS Red to allow for detection of the infected neurons. Control AAV (DS Red only) or AAV-Wnt2 was bilaterally injected into the dentate gyrus of adult rat hippocampus. Behavioral testing was done 3 weeks after viral infusion, a time point when AAV expression is optimal (Fig. 5A) (27). Expression (Fig. 5B) was analyzed by in situ hybridization (a,b) and DS-red staining (c). The exact cell types infected are difficult to assess, but since AAV2 selectively infects neurons the localization of signal to the granule cell layer and hilus indicates that neurons in these regions are being targeted. Animals were tested in behavioral models known to be responsive to antidepressant administration, including learned helplessness (LH), novelty suppressed feeding (NSF), sucrose preference test (SPT) and forced swim test (FST) (Fig. 5C–F). Over expression of Wnt2 significantly decreased the number of escape failures in the LH paradigm, an antidepressant-like response (Fig. 5C). There was also a decrease in the latency to feed in NSF, although this effect did not reach significance (Fig. 5D). Animals infused with AAV-Wnt2 also displayed an increase in sucrose consumption relative to AAV control infused animals, with no change in water consumption (Fig. 5E). The levels of sucrose preference (expressed as percent sucrose consumed over total fluid consumed) were 54 ± 3 and 63 ± 5 (mean $\pm$ S.E.M., for AAV-control and AAV-Wnt2 respectively; $p = 0.08$). Finally, AAV-Wnt2 had no significant effect in the FST (Fig. 5F) or locomotor activity

(number of beam breaks for AAV-control and AAV-Wnt2 were 580 ± 75 and 618 ± 50 , respectively; mean $\pm$ S.E.M.; $p = 0.33$).

DISCUSSION

The results demonstrate that chronic antidepressant administration influences hippocampal expression of multiple components of the Wnt/ β -catenin cascade, including regulation of subtypes of Wnts, Fz, β -catenin, and TCF. Notable among these signaling molecules is Wnt2, which was the only gene that was uniformly regulated by all treatments tested, including SSRI, SNRI, a dual reuptake inhibitor, and chronic ECS. Expression of some genes appeared to be more sensitive to a certain class of antidepressants. For example, FrzB, NeuroD1, and β -catenin were induced by SSRI administration, whereas Fz9 and Dvl1 were up-regulated by the SNRI and dual reuptake inhibitor. These findings suggest that certain components of Wnt signaling are differentially regulated by the 5-HT and NE neurotransmitter systems. However, there were also cases where the two SSRIs did not produce the same effects (e.g., TCF15 was induced by citalopram but not by fluoxetine). This could result from differences in the effective doses used or different actions of the two SSRIs tested. Further dose response studies, as well as additional SSRI and SNRI antidepressants will be required to determine the differential effects of these two classes of agents on selected Wnt signaling genes.

The microarray results for up-regulation of Wnt2 were confirmed by RT-PCR and by in situ hybridization, the latter demonstrating increased Wnt2 expression in the dentate gyrus granule cell layer. These findings are in agreement with a previous report that chronic ECS increases Wnt2 expression in the same hippocampal cell layer (25). The requirement for long-term (21d) drug treatment for induction of Wnt2 was confirmed by RT-PCR, as short-term treatment (5d) had no effect. We also found that chronic antidepressant treatment increased the phosphorylation of GSK-3 β , which causes inactivation, consistent with reports that pharmacological inhibition of GSK-3 β produces antidepressant-like behavioral effects (6,38–41). However, the mechanism underlying increased phosphorylation of GSK-3 β is unclear. Canonical Wnt signaling results in inhibition of GSK-3 β , but does so by destabilization of a GSK-3 β protein complex (42,43). There are reports that Wnt/Dvl1 signaling can lead to phosphorylation of GSK-3 β via the PI3 kinase/Akt pathway (Gomez and Arias et al; 2008; Fukumoto et al; 2001), and further studies will be required to determine the mechanism by which antidepressants increase phospho-GSK-3 β (i.e., via Wnt2/Dvl1/Akt or via another pathway such as growth factor receptor activation of PI3 kinase/Akt).

The functional actions at the behavioral level were determined using a viral expression approach. Induction of Wnt2 expression by microinfusion of AAV-Wnt2 into the dentate gyrus granule cell layer resulted in significant antidepressant-like responses in the LH and SPT, but no significant effects in NSF or FST. In the current LH paradigm expression of Wnt2 before exposure to IES assesses the ability of Wnt2 to prevent the development of escape deficits. It would also be interesting to test the influence of Wnt2 in a reversal paradigm, where Wnt2 is expressed after exposure to IES. Another consideration is the possibility that Wnt2 expression alters learning, which could also contribute to the more rapid responding observed in the active avoidance test. Expression of Wnt2 significantly influenced behavior in the SPT, which is typically used as a measure of anhedonia that occurs in response to repeated stress exposure (29). Although, the paradigm used in the current study did not include repeated stress, anhedonic responses were observed in AAV infused control animals (i.e., no preference for sucrose compared to water). This is likely due to the stress resulting from cranial surgery, as well as the stress encountered in the NSF and FST that precede the SPT.

Although not significant, there was a trend for AAV-Wnt2 to produce an effect in the NSF test, which is responsive to chronic antidepressant treatment. It would be interesting in future studies to determine if Wnt2 expression influences behavior in other models of anxiety, such as the elevated plus maze. The lack of effect of Wnt2 on the FST was surprising, but could be explained by several possibilities. First, the FST is a different despair paradigm relative to LH, notably being responsive to acute antidepressants, indicating that different signaling mechanisms are involved. Second, FST/immobility could be influenced by brain regions other than, or in addition to the hippocampus. Third, although expression of Wnt2 in the dorsal hippocampus was sufficient to alter LH and SPT behavior, it is possible that more widespread expression or targeting of other parts of hippocampus (e.g. ventral) would influence FST behavior. Although we have shown that antidepressant drugs decrease immobility in previous studies, a positive control was not included in the current study and it is possible that the animals were unresponsive for an unknown reason. Finally, although the results demonstrate that Wnt2 expression can influence LH and SPT behavior, future studies will be needed to determine if Wnt2 signaling is also necessary for the actions of antidepressant drug treatments. These studies will require the development of a Wnt2 knockdown strategy (e.g., viral expression of Wnt2 shRNA or Wnt2 conditional deletion mutant mice).

Wnt signaling pathways have been implicated in many aspects of neuronal morphology and function that could contribute to antidepressant-like actions, including axon and dendrite remodeling and development, synaptogenesis, neuroplasticity and neurogenesis (11–13,15,17). The actions of antidepressants are thought to occur in part via regulation of similar cellular processes (e.g., increased synaptic plasticity, spine/synapse formation, neurogenesis) and it is possible that these effects occur, in part, via regulation of Wnt signaling. For example Wnt2 signaling plays a critical role in activity dependent formation of dendrites via intracellular β -catenin (44); Wnt7b has been shown to regulate dendritic arborization via Dvl1 (15) and interestingly, we found that Dvl1 is up-regulated, but only by atomoxetine and venlafaxine, not by SSRIs. Regulation of neurogenesis in adult hippocampus is reported to occur via Wnt3a (45). Unfortunately, the signal strength of Wnt3a in the current study was too low to be included in the analysis, and further work will be required to determine if antidepressant treatment regulates the expression of Wnt3a, and alternatively, if Wnt2 regulates neurogenesis. Recent studies demonstrate that ablation of NeuroD1 in the adult neurogenic niche decreases neurogenesis and show that NeuroD1 links Wnt signaling and TCF/LEF activity (34,35). With regard to Wnt receptors, mutation of Fz9 decreases the number of dentate gyrus granule cells and reduces visual and spatial memory processing (46). Dvl1 is the major Dvl subtype in brain and recent studies have shown that Dvl1 accumulates at growing axon tips and is required for axon formation (47).

In summary, the results of this study provide evidence for the regulation of multiple components of Wnt/ β -catenin signaling, both canonical and noncanonical pathways (PCP, and Wnt/calcium cascades), showing that a complex network of molecular changes may contribute to the efficacy of antidepressant treatment. Importantly, the results demonstrate that local expression of Wnt2 in the hippocampus is sufficient to produce antidepressant-like effects in behavioral models. These studies identify specific molecular substrates within the Wnt/ β -catenin pathway that could be targeted for the development of novel therapeutic medications.

Acknowledgments

This work is supported by USPHS grants MH45481 (RSD), 2 P01 MH25642 (RSD), and the Connecticut Mental Health Center (RSD).

REFERENCES

1. Wong M-L, Licinio J. Research and treatment approaches to depression. *Nat Rev/Neurosci*. 2001; 2:343–351.
2. Chuang D, Manji HK. In search of the Holy Grail for the treatment of neurodegenerative disorders: has a simple cation been overlooked? *Biol Psych*. 2007; 62(1):4–6.
3. Mao Y, Ge X, Frank CL, Madison JM, Koehler AN, Doud MK, Tassa C, Berry EM, Soda T, Singh KK, Biechele T, Petryshen TL, Moon RT, Haggarty SJ, Tsai LH. Disrupted in schizophrenia 1 regulates neuronal progenitor proliferation via modulation of GSK3 β /beta-catenin signaling. *Cell*. 2009; 136(6):1017–1031. [PubMed: 19303846]
4. Silva R, Mesquita AR, Bessa J, Sousa JC, Sotiropoulos I, Leão P, Almeida OF, Sousa N. Lithium blocks stress-induced changes in depressive-like behavior and hippocampal cell fate: the role of glycogen-synthase-kinase-3 β . *Neurosci*. 2008; 152(3):656–669.
5. Gould T, Manji HK. The Wnt signaling pathway in bipolar disorder. *The Neuroscientist*. 2002; 8(5):497–511. [PubMed: 12374432]
6. Gould T, Einat H, Bhat R, Manji HK. AR-A014418, a selective GSK-3 inhibitor, produces antidepressant-like effects in the forced swim test. *Int J Neuropsychopharmacol*. 2004; 7(4):387–390. [PubMed: 15315719]
7. Kaidanovich-Beilin O, Milman A, Weizman A, Pick CG, Eldar-Finkelman H. Rapid antidepressant-like activity of specific glycogen synthase kinase-3 inhibitor and its effect on beta-catenin in mouse hippocampus. *Biol Psych*. 2004; 55(8):781–784.
8. Logan CY, Nusse R. The Wnt signaling pathway in development and disease. *Annu Rev Cell Dev Biol*. 2004; 20:781–810.
9. Kikuchi A, Yamamoto H, Kishida S. Multiplicity of the interactions of Wnt proteins. *Cellular Sig*. 2007
10. Li F, Chong ZZ, Maiese K. Vital elements of Wnt-Frizzled signaling pathway in the nervous system. *Curr Neurovasc Res*. 2005
11. Chizhikov V, Millen KJ. Roof plate-dependent patterning of the vertebrate dorsal central nervous system. *Dev Biol*. 2005; 277(2):287–295. [PubMed: 15617675]
12. Ciani L, Salinas PC. WNTs in the vertebrate nervous system: from patterning to neuronal connectivity. *Nat Rev Neurosci*. 2005; 6(5):351–362. [PubMed: 15832199]
13. Ikeya M, Lee SM, Johnson JE, McMahon AP, Takada S. Wnt signalling required for expansion of neural crest and CNS progenitors. *Nature*. 1997; 389(6654):966–970. [PubMed: 9353119]
14. Kiecker C, Niehrs C. A morphogen gradient of Wnt/ β -catenin signalling regulates anteroposterior neural patterning in *Xenopus*. *Development*. 2001; 128:4189–4201. [PubMed: 11684656]
15. Rosso SA, Sussman D, Wynshaw-Boris A, Salinas PC. Wnt signalling through Dishevelled, Rac and JNK regulates dendritic development. *Nat Neurosci*. 2005; 8:34–42. [PubMed: 15608632]
16. Lucas FRS PC. WNT-7a induces axonal remodeling and increases synapsin I levels in cerebellar neurons. *Dev Biol*. 1997; 192:31–44. [PubMed: 9405095]
17. Lyuksyutova AI, et al. Anterior-posterior guidance of commissural axons by Wnt-Frizzled signaling. *Science*. 2003; 302:1984–1988. [PubMed: 14671310]
18. Caricasole A, Copani A, Caruso A, et al. The Wnt pathway, cell-cycle activation and beta-amyloid: novel therapeutic strategies in Alzheimer's disease? *Trends Pharmacol Sci*. 2003; 24:233–238. [PubMed: 12767722]
19. Katsu T, Ujike H, Nakano T, et al. The human frizzled-3 (FZD3) gene on chromosome 8p21, a receptor gene for Wnt ligands, is associated with the susceptibility to schizophrenia. *Neurosci Lett*. 2003; 353:53–56. [PubMed: 14642436]
20. Kozlovsky N, Belmaker RH, Agam G. GSK-3 and the neurodevelopmental hypothesis of schizophrenia. *Eur Neuropsychopharmacol*. 2002; 12:13–25. [PubMed: 11788236]
21. Luo J, et al. Wnt signaling and human diseases: what are the therapeutic implications? *Lab Invest*. 2007; 87(2):97–103. [PubMed: 17211410]
22. Miyaoka T, Seno H, Ishino H. Increased expression of Wnt-1 in schizophrenic brains. *Schizophr Res*. 1999; 38:1–6. [PubMed: 10427605]

23. Newton SS, Collier EF, Hunsberger J, Adams D, Terwilliger R, Selvanayagam E, et al. Gene profile of electroconvulsive seizures: induction of neurotrophic and angiogenic factors. *J Neurosci*. 2003; 23(34):10841–10851. [PubMed: 14645477]
24. Ploski JE, Newton SS, Duman RS. Electroconvulsive seizure-induced gene expression profile of the hippocampus dentate gyrus granule cell layer. *J Neurochem*. 2006; 99(4):1122–1132. [PubMed: 17081142]
25. Madsen T, Newton SS, Eaton ME, Russell DS, Duman RS. Chronic electroconvulsive seizure up-regulates b-catenin expression in rat hippocampus: role in adult neurogenesis. *Biol Psychiat*. 2003; 54:1006–1014. [PubMed: 14625142]
26. Hunsberger JG, Newton SS, Bennett AH, Duman CH, Russell DS, Salton SR, et al. Antidepressant actions of the exercise-regulated gene VGF. *Nat Med*. 2007; 13(12):1476–1482. [PubMed: 18059283]
27. Hommel J, Sears RM, Georgescu D, Simmons DL, DiLeone RJ. Local gene knockdown in the brain using viral-mediated RNA interference. *Nat Med*. 2003; 9(12):1539–1544. [PubMed: 14634645]
28. Paxinos, G.; Watson, C. The rat brain in stereotaxic coordinates. San Diego: Academic; 1998.
29. Banasr M, Duman RS. Glial loss in the prefrontal cortex is sufficient to induce depressive-like behaviors. *Biol Psych*. 2008; 64(10):863–870.
30. Valentine G, Dow AL, Banasr M, Pittman B, Duman RS. Differential effects of chronic antidepressant treatment on shuttle box escape deficits induced by uncontrollable stress. *Psychopharmacology (Berl)*. 2008; 200(4):585–596. [PubMed: 18604599]
31. Banasr M, Valentine GW, Li XY, Gourley S, Taylor J, Duman RS. Chronic stress decreases cell proliferation in adult cerebral cortex of rat: Reversal by antidepressant treatment. *Biol Psych*. 2007a; 62:496–504.
32. Pliakas A, Carlson RR, Neve RL, Konradi C, Nestler EJ, Carlezon WA. Altered responsiveness to cocaine and increased immobility in the forced swim test associated with elevated CREB expression in the nucleus accumbens. *J Neurosci*. 2001; 21:7397–7403. [PubMed: 11549750]
33. Rodriguez J, Esteve P, Weint C, Ruiz JM, Fermin Y, Trousse F, Dwivedy A, Holt C, Bovolenta P. SFRP1 regulates the growth of retinal ganglion cell axons through the Fz2 receptor. *Nat Neurosci*. 2005; 8(10):1301–1309. [PubMed: 16172602]
34. Gao Z, Ure K, Ables JL, Lagace DC, Nave KA, Goebbels S, et al. Neurod1 is essential for the survival and maturation of adult-born neurons. *Nat Neurosci*. 2009; 12(9):1090–1092. [PubMed: 19701197]
35. Kuwabara T, Hsieh J, Muotri A, Yeo G, Warashina M, Lie DC, et al. Wnt-mediated activation of NeuroD1 and retro-elements during adult neurogenesis. *Nat Neurosci*. 2009; 12(9):1097–1105. [PubMed: 19701198]
36. Fukumoto S, Hsieh CM, Maemura K, Layne MD, Yet SF, Lee KH, Matsui T, Rosenzweig A, Taylor WG, Rubin JS, Perrella MA, Lee ME. Akt participation in the Wnt signaling pathway through Dishevelled. *J Biol Chem*. 2001; 276(20):17479–17483. [PubMed: 11278246]
37. Mercado-Gómez O, Hernández-Fonseca K, Villavicencio-Queijeiro A, Massieu L, Chimal-Monroy J, Arias C. Inhibition of Wnt and PI3K signaling modulates GSK-3 β activity and induces morphological changes in cortical neurons: role of tau phosphorylation. *Neurochem Res*. 2008; 33(8):1599–1609. [PubMed: 18461448]
38. Beaulieu J, Zhang X, Rodriguiz RM, Sotnikova TD, Cools MJ, Wetsel WC, Gainetdinov RR, Caron MG. Role of GSK3 β in behavioral abnormalities induced by serotonin deficiency. *PNAS*. 2008; 105(4):1333–1338. [PubMed: 18212115]
39. Gould T, Picchini AM, Einat H, Manji HK. Targeting glycogen synthase kinase-3 in the CNS: implications for the development of new treatments for mood disorders. *Curr Drug Targets*. 2006; 11:1399–1409. [PubMed: 17100580]
40. Gould T, Einat H, O'donnell KC, Picchini AM, Schloesser RJ, Manji HK. Beta-catenin overexpression in the mouse brain phenocopies lithium-sensitive behaviors. *Neuropsychopharmacol*. 2007; 32(10):2173–2183.

41. Rosa A, Kaster MP, Binfaré RW, Morales S, Martín-Aparicio E, Navarro-Rico ML, Martinez A, Medina M, García AG, López MG, Rodrigues AL. Antidepressant-like effect of the novel thiadiazolidinone NP031115 in mice. *Biol Psych*. 2008; 32(6):1549–1556.
42. Adli M, Hollinde DL, Stamm T, Wiethoff K, Tsahuridu M, Kirchheiner J, Heinz A, Bauer M. Response to lithium augmentation in depression is associated with the glycogen synthase kinase 3- β -50T/C single nucleotide polymorphism. *Biol Psych*. 2007; 62(11):1295–1302.
43. Jope R, Roh MS. Glycogen synthase kinase-3 (GSK3) in psychiatric diseases and therapeutic interventions. *Curr Drug Targets*. 2006; 7(11):1421–1434. [PubMed: 17100582]
44. Wayman G, Impey S, Marks D, Saneyoshi T, Grant WF, Derkach V, Soderling TR. Activity-dependent dendritic arborization mediated by CaM-kinase I activation and enhanced CREB-dependent transcription of Wnt-2. *Neuron*. 2006; 50(6):897–909. [PubMed: 16772171]
45. Lie D, Colamarino SA, Song HJ, Desire L, Mira H, Consiglio A, Lein ES, Jessberger S, Lansford H, Dearie AR, Gage FH. Wnt signalling regulates adult hippocampal neurogenesis. *Nature*. 2005; 437:1370–1375. [PubMed: 16251967]
46. Zhao C, Pleasure SJ. Hippocampal and visuospatial learning defects in mice with a deletion of frizzled 9, a gene in the Williams syndrome deletion interval. *Development*. 2005; 132(12):2917–2927. [PubMed: 15930120]
47. Zhang X, et al. Dishevelled promotes axon differentiation by regulating atypical protein kinase C. *Nat Cell Biol*. 2007; 9(7):743–754. [PubMed: 17558396]

Figure 1. Microarray analyses of Wnt signaling components in hippocampus in response to chronic antidepressant administration

Rats were administered vehicle, citalopram (CIT, 15 mg/kg), fluoxetine (FLX, 5 mg/kg), venlafaxine (VNF, 15 mg/kg), or atomoxetine (ATX, 3 mg/kg) twice daily for 3 weeks. ECS was administered once daily for 10 days. Gene expression in the hippocampus was determined by microarray analysis using a focused microarray. A. Wnt/Fz/Dvl subtype expression. B. TCF/LEF subtype and NeuroD1 expression C. β -catenin/Akt1 expression. The results are expressed as the mean ratio of fold change in gene expression, and are the mean $\pm$ S.E.M. of n=6 (vehicle and drugs) or n=4 (sham and ECS). Horizontal line at ratio

of 1 indicates no regulation. * $p < 0.05$ (Student's t-test; $p < 0.05$; no FDR adjustment). # $p < 0.05$ (Student's t-test on samples after adjustment for FDR; $p < 0.05$).

Figure 2. Confirmation of microarray results by RT-PCR

Quantitative RT PCR analysis of Wnt2 expression by chronic (3wks) chemical antidepressant administration (A) in hippocampus. Data in both A and B are normalized to cyclophilin and are the mean ratio (antidepressant:vehicle) of fold change $\pm$ S.E.M. of 5 individual animals. Horizontal line at ratio of 1 indicates no regulation. * $p < 0.05$; compared to vehicle-treated control (Student's t-test)

Figure 3. Chronic antidepressant administration increases Wnt2 mRNA in dentate gyrus of the hippocampus

The expression of Wnt2 mRNA in dentate gyrus was determined by in situ hybridization analysis. (A) Representative images of emulsion-dipped slides at different magnifications: upper; 10X, middle; 100X, lower; 400X. (B) Quantitation of Wnt2 mRNA in the dentate gyrus. Results are presented as the mean average number of grains $\pm$ S.E.M. of 6 replicates. * $p < 0.05$; compared to vehicle-treated control (Student's *t*-test). ML; molecular layer, DG; dentate gyrus, GCL; granular cell layer.

Figure 4. Chronic antidepressant administration increases GSK-3β phosphorylation

Rats were administered vehicle (distilled water, twice daily), fluoxetine (5 mg/kg, twice daily), or venlafaxine (15 mg/kg, twice daily) and whole hippocampus was subjected to western blot analysis. Optical density values were normalized to loading control (Glyceraldehyde-3-Phosphate dehydrogenase (GAPDH)) and expressed as the ratio of phospho-Ser9- GSK-3β/total GSK-3β. Data are expressed as a mean fold change ± S.E.M, compared to vehicle-treated controls (n=6); *p<0.05 (Student's t-test).

Figure 5. Influence of AAV-Wnt2 expression on behavioral models of depression and antidepressant response

AAV-Wnt2 or control (AAV-DS Red) was infused into the dentate gyrus of the hippocampus and tissue was harvested 8 weeks later. (A) Schematic diagram showing details of experimental timeline. (B) In situ hybridization analysis was conducted to determine the expression of Wnt2, and representative autoradiograms were shown for control (a) and AAV-Wnt2 (b). DS-red staining is also shown to further demonstrate localization of AAV spread (c) The influence of AAV-Wnt2 expression on different behavioral models was examined, including (C) escape failures in the learned helplessness (LH) paradigm; * $p=0.02$; (D) latency to feed in the novelty suppressed feeding (NSF) test;

p=0.09; (E) sucrose preference (SPT) test; *p=0.02 for sucrose consumption; p=0.08 for sucrose preference; (F) latency to immobility in the forced swim test (FST); p=0.39; Immobility; p=0.47. Results are the mean $\pm$ SEM of Control (n=8) vs. Wnt2 (n=7). *p < 0.05 compared to AAV-controls (Student's t-test).