Published in final edited form as:

J Abnorm Child Psychol. 2009 January ; 37(1): 45–58. doi:10.1007/s10802-008-9257-1.

Are Oppositional-Defiant and Hyperactive-Inattentive Symptoms Developmental Precursors to Conduct Problems in Late Childhood?: Genetic and Environmental Links

Benjamin B. Lahey,

Departments of Health Studies and Psychiatry, University of Chicago

Carol A. Van Hulle,

Waisman Center, University of Wisconsin

Paul J. Rathouz,

Department of Health Studies, University of Chicago

Joseph Lee Rodgers,

Department of Psychology, University of Oklahoma

Brian M. D'Onofrio, and

Department of Psychological and Brain Sciences, Indiana University

Irwin D. Waldman

Department of Psychology, Emory University

Abstract

Inattentive-hyperactive and oppositional behavior have been hypothesized to be developmental precursors to conduct problems. We tested these hypotheses using a longitudinal sample of 6,466 offspring of women selected from nationally-representative U.S. households. Conduct problems across 8–13 years were robustly predicted by conduct problems at 4–7 years, but also were independently predicted to a small extent by both inattentive-hyperactive and oppositional behaviors at 4–7 years. Longitudinal multivariate behavior genetic analyses revealed that the genetic and environmental factors that influence conduct problems at both 4–7 and 8–13 years also influence the putative precursors at 4–7 years. After genetic and environmental influences on conduct problems at 4–7 years were taken into account, however, inattentive-hyperactive and oppositional behavior at 4–7 years shared causal influences with conduct problems 8–13 years to a negligible extent. These findings suggest that after early conduct problems are controlled, little is gained in terms of prediction or understanding genetic and environmental influences on later child conduct problems by treating early inattentive-hyperactive and oppositional behavior as developmental precursors to later conduct problems.

An important goal of developmental psychopathology is the description of regular developmental progressions from early precursors to more serious problems later in development. This is because it would be both theoretically revealing and practically useful if milder problems (a) typically emerge earlier than more serious problems, and (b) reliably predict later serious problems. In such cases, it should be possible to identify factors that govern the developmental progression from milder to more serious problems and find ways to prevent

serious problems. Therefore, considerable effort has been made to identify the developmental precursors to serious conduct problems in older children and adolescents.

The behaviors that define attention-deficit/hyperactivity disorder (ADHD) and oppositional defiant disorder (ODD) have long been hypothesized to be developmental precursors to serious conduct problems (Lahey, McBurnett, & Loeber, 2000). ADHD refers to developmentally-inappropriate levels of attention problems, motor hyperactivity, and impulsive behavior. ODD is defined by developmentally-inappropriate levels of irritability, defiance, and the intentional annoyance of others.

Do ADHD and ODD Precede Conduct Problems?

By definition, developmental precursors must precede the more serious problem in development. Unfortunately, there appears to be only one study in which the ages of onset of symptoms of ADHD, ODD, and conduct problems were directly compared in the same sample. A clinic study of 7–12 year old males, which retrospectively assessed age of onset of all symptoms that were reported during the clinic assessment, found that most symptoms of ADHD and ODD emerged early in life, with half of their symptoms having median ages of onset by 5 years of age (Loeber, Green, Lahey, Christ, & Frick, 1992). The median ages of emergence for the remaining symptoms of ADHD and ODD were later, however, with the last symptoms of each disorder emerging around 11 years (Loeber et al., 1992). In the same study, most symptoms of conduct disorder (CD) were reported to emerge only slightly later than those of ADHD and ODD, with the median age of onset for half of CD symptoms reported by 6 years. On the other hand, a number of serious conduct problems, such as forced sex, theft by confrontation, running away from home, and truancy, did not emerge until late childhood or adolescence (Loeber et al., 1992). Thus, this study suggested that many symptoms of all three disorders are present before 6 years of age, even though some CD symptoms emerge later than those of ADHD and ODD. This study is of limited value, however, because it is based on retrospective reports from a clinic sample that was not representative. Nonetheless, the ages of onset found in this study for ADHD, ODD, and CD symptoms are similar to those found in other clinic-based (Applegate et al., 1997) and population-based studies (Lahey et al., 1998; Willoughby, Curran, Costello, & Angold, 2000) that did not compare ages of onset of across disorders.

Furthermore, recent longitudinal studies that began during toddler and preschool ages clearly indicate that many young children already exhibit serious conduct problems during the early childhood years when most symptoms of ADHD and ODD are first evident (Keenan et al., 2007; Tremblay, 2004). Thus, although there may be a longer developmental window for the emergence of some serious conduct problems than most ADHD and ODD symptoms, it seems likely that many conduct problems emerge in early childhood concurrently with symptoms of ADHD and ODD. This raises a question about the extent to which early symptoms of ADHD or ODD function as independent developmental precursors to later child conduct problems.

Does Childhood ADHD Predict Later Conduct Problems?

Seminal prospective longitudinal studies (Hechtman, Weiss, & Perlman, 1984; Loney, Kramer, & Milich, 1981) showed that children with ADHD symptoms are at increased risk for exhibiting serious conduct problems in adolescence. These studies did not control for conduct problems in childhood, however. Some more recent longitudinal studies did not support the hypothesis that childhood ADHD symptoms predict later conduct problems when early childhood conduct problems were controlled (Lahey et al., 2000; Lilienfeld & Waldman, 1990; Loeber, 1988; Loeber, Green, Keenan, & Lahey, 1995; Magnusson, 1984; Satterfield & Schell, 1997). Firm conclusions cannot be reached, however, as other longitudinal studies suggest that ADHD predicts later conduct problems even in the apparent absence of high levels of childhood

conduct problems (Lambert, 1988; Mannuzza, Klein, Abikoff, & Moulton, 2004; Mannuzza, Klein, Bessler, Malloy, & LaPadula, 1993; Mannuzza et al., 1991).

Does Childhood ODD Predict Later Conduct Problems?

Several studies suggest that childhood ODD predicts later serious conduct problems. A longitudinal study of 7–12 year old clinic-referred boys found that ODD in the first assessment was a significant predictor of meeting criteria for CD for the first time during the next 7 years (Lahey et al., 2000; Loeber et al., 1995). Another longitudinal study of clinic-referred 6–17 year old males who met criteria for ADHD found that ODD in the first assessment did not predict meeting criteria for CD over the next three years (Biederman et al., 1996). When the prevalence of new cases of CD among males who met criteria for ODD in the baseline assessment was compared to all youth who did not meet criteria for ODD at baseline in this sample (those with only ADHD and the normal controls), however, CD was significantly more likely to emerge at follow-up among youths with ODD (Biederman et al., 1996). Similarly, a longitudinal study of a representative sample reported that meeting criteria for ODD in the first assessment predicted meeting criteria for CD for the first time in later assessments in males but not females (Rowe, Maughan, Pickles, Costello, & Angold, 2002). These studies suggest that meeting criteria for ODD at an early age means that a child is at increased risk for later meeting criteria for CD for the first time. This does not necessarily mean that the emergence of all symptoms of ODD preceded the emergence of all symptoms of CD in development, however. It only refers to the sequence of meeting criteria for the two disorders.

Are There Genetic and Environmental Links Between Conduct Problems and Earlier ADHD and ODD?

If children with high levels of ADHD or ODD symptoms prove to be at high risk for future serious conduct problems, it would be very important to know *why* this occurs. For example, if early ADHD and/or ODD symptoms are early manifestations of the same genetic or environmental influences that will influence conduct problems later in childhood, knowing that is the case should facilitate the design of early interventions to reduce antisocial outcomes. This is clearly a possibility, as many cross-sectional studies suggest that concurrently assessed ADHD, ODD, and conduct problems are largely influenced by the same genetic factors (Dick, Viken, Kaprio, Pulkkinen, & Rose, 2005; Nadder, Rutter, Silberg, Maes, & Eaves, 2002; Waldman, Rhee, Levy, & Hay, 2001) and perhaps the same environmental factors (Burt, McGue, Krueger, & Iacono, 2005).

The critical question, however, is whether early childhood ADHD and ODD share genetic or environmental influences with *later* conduct problems, *over and above* the genetic and environmental influences that later conduct problems share with *earlier* conduct problems. If so, obtaining measures of ADHD and ODD in early childhood would allow prevention researchers to identify children who are at risk for later conduct problems for reasons not revealed by their early conduct problems. If early ADHD and ODD do not share genetic and environmental influences with later conduct problems beyond those shared with early conduct problems, the focus of prevention research probably should be on early childhood conduct problems rather than on ADHD and ODD.

Therefore, we conducted two kinds of analyses of longitudinal data from a large and diverse sample of the offspring of a representative sample of mothers. First, we used regression analyses to test the predictive hypotheses that inattentive-hyperactive behaviors resembling symptoms of ADHD and oppositional behaviors resembling symptoms of ODD at 4–7 years of age predict conduct problems at 8–13 years of age, controlling for demographic factors and conduct problems at 4–7 years of age. Second, we used longitudinal multivariate behavior genetic analyses to test the hypotheses that inattentive-hyperactive and oppositional behavior

at 4–7 years share genetic or environmental influences with those on conduct problems at 8–13 years, over and above the genetic and environmental influences shared by the putative precursors with conduct problems at 4–7 years.

METHOD

Sample

The National Longitudinal Survey of Youth (NLSY) was funded by the Bureau of Labor Statistics to study the U.S. workforce. A nationally-representative household sample of 9,763 14–22 year old male and female youth who were not in the military was selected in 1979 using a complex survey design, with an oversample of African American and Hispanic youth. To date, 4,926 NLSY females (1,472 African American, 977 Hispanic, and 2,477 non-Hispanic European American and other groups) have given birth to children who have participated in the assessments of offspring. Biennial assessments of all biological offspring of each NLSY woman have been conducted during 1986–2004 on the Children of the NLSY (CNLSY) (Chase-Lansdale, Mott, Brooks-Gunn, & Phillips, 1991), with participation in these assessments averaging $\approx 90\%$. In order to conduct longitudinal analyses on the same children at different ages, the present analyses were conducted on 6,994 offspring of 2,256 mothers who were assessed at least once during 4–7 years and at least once during 8–13 years of age. Because total family income is an important covariate in the regression analyses, and the variable with the greatest amount of missing data in survey research, the present regression analyses were limited to the 6,466 offspring of women who reported income data. Descriptive statistics on the sample are presented in Table 1. Because the longitudinal behavior genetic analyses could only be conducted on children with at least one sibling, both of whom were assessed at both 4–7 and 8–13 years, 5,281 sibling pairs were included in these analyses (Table 1).

Because only biological children of NLSY mothers participated, all CNLSY siblings are at least half siblings. Algorithms were developed to classify offspring pairs as either twins, full siblings, or half siblings, with some pairs classified as ambiguous siblings because they share a biological mother, but their sharing of the same biological father is uncertain (Rodgers, Johnson, & Bard, 2005; Rodgers, Rowe, & Li, 1994). The CNLSY kinship links (including for ambiguous siblings) used in the present and previous analyses of this data set (Van Hulle, Rodgers, D’Onofrio, Waldman, & Lahey, 2007) were validated by demonstrating that the heritability of physical height is .90, consistent with twin studies (Rodgers et al., 1994). Because ambiguous siblings are either full or half siblings, they were assigned a genetic coefficient of $r_g = .375$, intermediate between full ($r_g = .5$) and half siblings ($r_g = .25$). There were too few identical twin pairs ($n = 11$) to include them in these analyses, but the 23 fraternal twin pairs were included with other full siblings.

Measures of Child Problem Behavior

In each assessment across 4–13 years of age, mothers rated their children’s behavior using the Behavior Problem Index (BPI) (Peterson & Zill, 1986). The BPI was created for the CNLSY by selecting items from the *Child Behavior Checklist* (Achenbach, 1978) that were strongly correlated with factor scores. The mean of 3 BPI items rated on a 3-point scale were used to define *inattentive-hyperactive behavior*: has difficulty concentrating, cannot pay attention for long; impulsive, or acts without thinking; and restless or overly active, cannot sit still. The mean of 3 BPI items define *oppositional behavior*: argues too much; stubborn, sullen, or irritable; and has a strong temper and loses it easily. The mean of 7 BPI items rated on a 3-point scale define child *conduct problems* at each age: cheats or tells lies; has trouble getting along with teachers; disobedient at home; disobedient at school; bullies or is cruel or mean to others; breaks things on purpose or deliberately destroys things; and does not seem to feel sorry after misbehaving. Confirmatory factor analysis of these 15 items supported the distinction of

these 3 dimensions of problem behaviors (D'Onofrio et al., 2008). Each mean problem behavior score was multiplied by the number of items in the scale. The median alpha coefficients over ages 4–13 years for inattentive-hyperactive behavior, oppositional behavior, and conduct problems was .70, .73, and .74, respectively. These definitions of problem behaviors overlap substantially with those used in previous longitudinal studies of population-based samples (Fergusson & Horwood, 2002; Lahey et al., 2000; Moffitt, Caspi, Dickson, Silva, & Stanton, 1996). The CNLSY measure of child conduct problems is valid in the sense of robustly predicting convictions for nontrivial offenses during adolescence in the CNLSY (Lahey et al., 2006). Similarly, the inattentive-hyperactive score predicts tested academic achievement controlling for oppositional behavior and conduct problems (D'Onofrio et al., 2007).

Demographic Covariates

Mothers classified their offspring using mutually exclusive categories as African American, Hispanic, or all other race-ethnic groups. Total family income was assessed in the wave when the mothers were closest to 30 years old to index earning capacity, expressed as the logarithm of 1986 inflation-adjusted dollars. This includes all income received by the woman and her spouse (if married), including food stamps and other forms of governmental support. Income from any cohabiting partner was not included, but there are few substantive differences when this source of income is included in economic analyses (Avellar & Smock, 2005).

Statistical Analysis

Predictive Regression Analyses—We first tested the roles of inattentive-hyperactive and oppositional behavior at 4–7 years as predictors of conduct problems at 8–13 years in the same children using predictive regression analyses, controlling for conduct problems at 4–7 years. Because the response variable of conduct problems at 8–13 years was highly skewed and kurtotic with a mode of zero, log-linear regression was conducted specifying Poisson working distributions in SAS PROC GENMOD. In order to take the clustering of children within families into account, these regressions were conducted in generalized estimating equations using robust standard errors (Zeger & Liang, 1986).

Longitudinal Multivariate Behavior Genetic Analyses—To determine if genetic and environmental influences on conduct problems at 8–13 years were shared with those on inattentive-hyperactive and oppositional behavior at 4–7 years, after genetic and environmental influences on conduct problems at 8–13 years shared with conduct problems at 4–7 years were taken into account, two separate series of multivariate behavior genetic analyses were conducted. Total observed variance in each of the four measures of child problem behavior based on the correlations among siblings of difference genetic relatedness (Table 1) was partitioned into estimates of variance attributable to latent additive genetic influences, shared environmental influences (i.e., shared experiences such as neighborhood factors that make siblings more similar), and nonshared environmental influences (i.e., either only one sibling experiencing an event such as bullying, or events experienced by both siblings that influence them in different ways, plus measurement error) based on differences in correlations between sibling pairs of differing genetic relatedness. Separate multivariate Cholesky decompositions (Neale & Cardon, 1992) were fit for the two putative precursors, one for inattentive-hyperactive behavior and one for oppositional behavior. Cholesky decomposition is a saturated model with as many potential underlying additive genetic and environmental influences as measured variables (Loehlin, 1996).

In the first series of Cholesky decompositions, the order of the three phenotypes was conduct problems at 4–7 years, oppositional behavior at 4–7 years, and conduct problems at 8–13 years. As illustrated in Figure 1, the first genetic factor specified (A_1) influenced conduct problems at 4–7 years (path coefficient a_{11}), oppositional behavior at 4–7 years (a_{21}), and conduct

problems at 8–13 years (a_{31}). The second genetic factor (A_2) influenced oppositional behavior at 4–7 years (a_{22}) and potentially conduct problems at 8–13 years (a_{32}). If path a_{32} from A_2 to conduct problems at 8–13 years can not be dropped from the model without a significant loss of fit, that would indicate that there are genetic influences on oppositional behavior at 4–7 years that influence conduct problems at 8–13 years that are distinct from the genetic influences on conduct problems at 4–7 years. The shared environmental factors (C_1 and C_2) and nonshared environmental factors (E_1 and E_2) have same interpretations as A_1 and A_2 , but for these environmental influences. The same tests for pathways from these shared and nonshared environmental factors (i.e., paths c_{32} and e_{32} in Figure 1) were conducted separately for each factor to determine if early oppositional behavior shares environmental influences with conduct problems at 8–13 years that are distinct from the environmental influences on conduct problems at 4–7 years. A second series of models conducted the same tests for the relevant genetic and environmental parameters for inattentive-hyperactive behavior at 4–7 years.

In previous analyses of this sample, we found that variances for conduct problems at all ages were greater for children from blended families (those with at least one half sibling) than non-blended families (all full siblings) and the term for shared environmental influences in non-blended families was nonsignificant, even though a shared environmental influence term specific to blended families was significant (Waldman et al., 2008). Therefore, a term for shared environmental influences was specified only for blended families in the present biometric models.

In each of the two series of Cholesky decompositions, omnibus tests were conducted to determine if there were sex differences in the structure of genetic and environmental influences on developing antisocial behavior. Model 1 (Sex Equal) constrained all parameters to be equal for females and males. Model 2 (Scalar Sex Differences) constrained all parameters to be equal between sexes, but allowed the total variation/covariation in males to be a scalar multiple of the total variation/covariation in females (Neale & Cardon, 1992). If Model 2 were the best fitting model, that would imply that the total influence of genetic and environmental influences on conduct problems is proportionally less in one sex than the other. Model 3 (Quantitative Sex Differences) allowed all parameters to vary between sexes, but the underlying factors were the same for both sexes.

Because the distributions of the three phenotypes were non-normal at all ages, models were fitted in Mplus 4.2 using robust maximum likelihood estimation, taking the clustering of children within families into account (Muthén & Muthén, 2007). The significance of model parameters was determined by comparing the fit of alternative models. The overall fit of each model was expressed as a means-adjusted Satorra-Bentler χ^2 and differences between models were tested using the Satorra-Bentler scaled-difference χ^2 (Satorra & Bentler, 2001) and χ^2 tests. A non-significant change in χ^2 would indicate that the model with fewer parameters fits the data equally well and would be preferred based on parsimony. Because concerns have been raised about fitting Cholesky models (Carey, 2005; Dominicus, Skrondal, Gjessing, Pedersen, & Palmgren, 2006), we present fit statistics in addition to the Satorra-Bentler χ^2 . Lower values of the Bayesian information criterion (BIC) reflect better fit and/or greater parsimony. The root mean square error of approximation (RMSEA) reflects the discrepancy between observed and predicted covariances per degree of freedom. Values of RMSEA < .05 are indicative of a close fit and values < .08 indicate a good fit to the data (MacCallum, Browne, & Sugawara, 1996).

Tests of Sample Bias—The 6,994 offspring used in the present analyses are a subset of the total 11,428 identified CNLSY offspring. Some offspring were excluded from the present analyses because they were too old when assessments of offspring began in 1986 to be assessed at 4–7 years of age. Most excluded youth were too young in 2004 to have been assessed at 8–13 years of age, and some had missing data in one age range due to non-participation. Compared

to the 4,437 CNLSY offspring whose data were not used, the offspring who participated in the present analyses were not significantly more likely at $p < .05$ to be boys and did not differ in log family income when the mother was 30 years old, but were more likely to be Hispanic, odds ratio = 1.46 (95% confidence interval: 1.32–1.61) and more likely to be African American, odds ratio = 1.52 (95% confidence interval: 1.40–1.66). These demographic variables were used as covariates in the predictive regression analyses to estimate population parameters more accurately.

RESULTS

The means and standard deviations of each measure of child problem behavior at 4–7 years and conduct problems at 8–13 years are presented in Table 2. As is typical for maternal ratings of problem behaviors across these ages (Leve, Kim, & Pears, 2005), there was a slight decrease in conduct problems at the older ages. The Spearman rank correlation between inattentive-hyperactive behavior and oppositional behavior at 4–7 years was $\rho = .55$ ($n = 3559$, $p < .0001$) for boys and $\rho = .52$ ($n = 3443$, $p < .0001$) for girls. The correlation between inattentive-hyperactive behavior and conduct problems at 4–7 years was $\rho = .58$ ($n = 3559$, $p < .0001$) for boys and $\rho = .52$ ($n = 3433$, $p < .0001$) for girls. The correlation between oppositional behavior and conduct problems at 4–7 years was $\rho = .62$ ($n = 3560$, $p < .0001$) for boys and $\rho = .58$ ($n = 3433$, $p < .0001$) for girls. The only significant sex difference in correlations among the measures of disruptive behavior at 4–7 years was between oppositional behavior and conduct problems, $p < .01$.

Prospective Prediction Analyses

Formal Predictive Tests—The first set of analyses formally tested the extent to which child conduct problems, inattentive-hyperactive behavior, and oppositional behavior at 4–7 years predict conduct problems at 8–13 years using log-linear regression. As shown in Table 3, when each early putative precursor and early conduct problems were tested in separate regression models, controlling the child's sex, ethnicity (two contrasts for African American versus other and Hispanic versus other), and log family income, each early putative predictor accounted for significant variance in conduct problems at 8–13 years. When early inattentive-hyperactive behavior, oppositional behavior, and conduct problems were all entered simultaneously with the demographic covariates, each behavioral predictor still accounted for significant variance in conduct problems at 8–13 years, although the magnitudes of the beta coefficients for inattentive-hyperactive and oppositional behavior dropped by more than 67%. When interactions between child sex and each of the three early behavioral predictors was tested in this joint model, only the sex-by-early conduct problems interaction was significant, $\beta = -.05$, $z = -2.07$, $p < .04$, reflecting a slightly stronger prediction in girls.

Explained Variance—In order to provide easily interpreted estimates of effect sizes, correlations between each predictor at 4–7 years and conduct problems at 8–13 years are presented in Table 4. Because the distributions of the behavior problem scores were not normal, Spearman rank correlations were calculated. Sex differences in the magnitudes of all of the predictive zero-order correlations between earlier and later problem behaviors in Table 4 were small, but statistically significant in each case in this large sample.

To reveal the extent to which inattentive-hyperactive and oppositional behaviors independently predicted conduct problems at 8–13 years after the correlation with conduct problems at 4–7 years was taken into account, two sets of partial correlations also were calculated to estimate predictive effect sizes for each predictor at 4–7 years (1) controlling the demographic covariates only, and (2) controlling the demographic covariates and both of the other two behavioral predictors at 4–7 years.

The zero-order Spearman correlations show that conduct problems at 4–7 years explained 28–37% of the variance in conduct problems at 8–13 years. Similarly, inattentive-hyperactive and oppositional behaviors at 4–7 years each explain 17–23% of the variance in conduct problems at 8–13 years. When the demographic covariates were partialled, there was little reduction in the variance in conduct problems at 8–13 years explained by the three predictors. In contrast, when inattentive-hyperactive and oppositional behaviors were tested partialling both the demographic covariates and conduct problems at 4–7 years, oppositional behavior explained only 2% of the variance in conduct problems at 8–13 years in both sexes and inattentive-hyperactive behavior explained 2–4% of the variance in conduct problems at 8–13 years. Thus, inattentive-hyperactive and oppositional behavior each explained little variance in future conduct problems after early conduct problems were taken into account.

Genetic and Environmental Influences Shared With Early Oppositional Behavior

Tests of omnibus sex differences—The first series of Cholesky decompositions tested for omnibus sex differences in the sharing of genetic and environmental influences among oppositional behavior and conduct problems. As reported in Table 5, the scalar sex-differences model fit significantly better than the model allowing no sex differences, but the quantitative sex-differences model did not fit significantly better than the scalar sex-differences model. Therefore, the scalar sex-differences model was used as the basis for the tests of specific genetic and environmental parameters to provide a standard of reference.

Tests of hypotheses regarding specific genetic and environmental parameters

—As reported in Table 6, a second series of Cholesky decompositions was conducted to test hypotheses regarding specific genetic and environmental pathways from oppositional behavior at 4–7 years to conduct problems at 8–13 years. The base model for this series was Model 2 (scalar sex differences) in Table 5. Because pathway a_{33} from genetic factor A_3 (see Figure 1) was estimated to be 0 for both sexes in Model 2, we first determined if a_{33} (unique genetic influences on conduct problems at 8–13 years) could be dropped by eliminating pathway a_{33} from Model 2. The lack of a significant Wald χ^2 between Model 3 and Model 2 indicated that a_{33} could be dropped without significant loss in fit, indicating that Model 2 was preferred on the basis of parsimony. Therefore, the key hypotheses regarding the specific pathways relevant to the sharing of genetic and environmental influences between the putative developmental precursor of oppositional behavior at 4–7 years and conduct problems at 8–13 years (a_{32} , c_{32} , and e_{32}) were tested by comparing the fits of Models 5, 6, and 7 (in which specific pathways were dropped) to Model 2.

In Model 5, parameter a_{32} (genetic influences shared by oppositional behavior and later conduct problems, separate from genetic factor A_1) was dropped from Model 2 (along with a_{33}).

Comparison of Model 5 to Model 2 revealed that dropping genetic parameter a_{32} did not result in statistically significant decrease in fit, indicating that Model 5 was preferred. BIC for Model 5 was slightly lower than for Model 2, reflecting its greater parsimony.

In Model 6, parameter c_{32} (shared environmental influences shared by oppositional behavior and later conduct problems, separate from shared environmental factor C_1) was dropped from Model 2 (along with a_{33}). Comparison of Model 6 with Model 2 revealed that it was possible to drop c_{32} without significant loss in fit, indicating that Model 6 was preferred. BIC for Model 6 was slightly lower than for Model 2, reflecting its greater parsimony.

In Model 7, parameter e_{32} (nonshared environmental influences shared by oppositional behavior and later conduct problems, separate from nonshared environmental factor E_1) was dropped from Model 2 (along with a_{33}). Comparison with Model 4 revealed that was possible to drop the nonshared environmental parameter e_{32} without significant loss in fit, indicating

that Model 7 was preferred. Again, BIC for Model 7 was lower than for Model 2, reflecting its greater parsimony.

Squared standardized coefficients for each path in the more complete Model 2 are presented in Figure 2. Based on the comparative tests of the alternative models reported in Table 6, paths a_{32} , c_{32} , and e_{32} were omitted from Figure 2. Path e_{33} was not omitted even though its 95% confidence interval included 0 because it includes the estimate of error in conduct problems at 8–13 years. Overall, Figure 2 shows that genetic factor A_1 robustly influences conduct problems at both ages (a_{11} and a_{31}) and early oppositional behavior (a_{21}). Indeed, there were no genetic influences on conduct problems at 8–13 years that were not shared with earlier conduct problems. In contrast, genetic factor A_2 , which also influences early oppositional behavior, did influence later conduct problems. There also appeared to be very limited common environmental influence on early and later problem behaviors.

Genetic and Environmental Influences Shared With Early Inattentive-Hyperactive Behavior

Tests of omnibus sex differences—As shown in Table 7, the scalar sex-differences model fit significantly better than the model allowing no sex differences, but the quantitative sex-differences model fit significantly better than the scalar sex-differences model. Therefore, the quantitative sex-differences model was used as the basis for the tests of specific genetic and environmental parameters. As a result, the alternative models had to be tested separately for the two sexes, but the results were almost always the same.

Tests of hypotheses regarding specific genetic and environmental parameters

—The base model for the series of Cholesky decompositions to test hypotheses regarding specific genetic and environmental pathways from inattentive-hyperactive behavior at 4–7 years to conduct problems at 8–13 years was Model 3 in Table 7. Because pathway a_{33} from genetic factor A_3 (see Figure 1) was estimated to be essentially 0 for both sexes, we first determined if a_{33} (unique genetic influences on conduct problems at 8–13 years) could be dropped from Model 3 for inattentive-hyperactive behavior. The non-significant Wald χ^2 (Satorra & Bentler, 2001) between Model 3 and Model 4 indicated that a_{33} could be dropped without significant loss in fit. Therefore, Model 4 was preferred on the basis of parsimony, which is reflected in the slightly lower BIC for Model 4 than for Model 3. Thus, the hypotheses regarding specific pathways relevant to the sharing of genetic and environmental influences between the putative developmental precursor of inattentive-hyperactive behavior at 4–7 years and conduct problems at 8–13 years (a_{32} , c_{32} , and e_{32}) were tested by comparing models with each dropped pathway to Model 4 (from which a_{33} was dropped).

In Model 5, parameter a_{32} (genetic influences shared by inattentive-hyperactive behavior and later conduct problems, separate from genetic factor A_1) was dropped (along with a_{33}). As shown in Table 7, the comparison of Model 5 to Model 4 showed that it was possible to drop genetic parameter a_{32} without a significant decrease in fit. Correspondingly, the BIC for the more parsimonious Model 5 was slightly smaller than for Model 4. In Model 6, parameter c_{32} (shared environmental influences shared by inattentive-hyperactive behavior and later conduct problems, separate from shared environmental factor C_1) was dropped (along with a_{33}). Comparison of Model 6 with Model 4 revealed that it was not possible to drop parameter c_{32} (shared environmental influences in blended families that are common to both inattentive-hyperactive behavior at 4–7 years and to conduct problems at 8–13 years) without a significant loss in fit. This reflects the sex difference in paths from C_2 . As shown in the two panels of Figure 3, either c_{22} or c_{32} is negligible for one of the sexes. Either way, the implication is that there are no appreciable shared environmental influences common to both inattentive-hyperactive behavior at 4–7 years and conduct problems at 8–13 years independent of conduct problems at 4–7 years.

In Model 7, parameter e_{32} (nonshared environmental influences shared by inattentive-hyperactive behavior and later conduct problems, separate from nonshared environmental factor E_1) was dropped (along with a_{33}). Comparison of Model 7 with Model 4 revealed that it was possible to drop e_{32} without significant loss in fit. BIC for Model 7 was slightly lower than for Model 4, reflecting its greater parsimony.

Squared standardized coefficients for each path in Model 4 for each sex are presented in Figure 3, except that parameters were omitted for each sex when their 95% confidence intervals included 0 for that sex. Coefficient a_{32} was not presented in Figure 3 for either sex because it could be dropped without significant loss in fit and its 95% confidence intervals included 0 for both sexes. Path e_{33} was not omitted from Figure 3 even though its 95% confidence interval included 0 because it includes the estimate of error in conduct problems at 8–13 years.

DISCUSSION

As in many previous studies (Lahey et al., 2000; Lilienfeld & Waldman, 1990), conduct problems at 8–13 years were significantly predicted by both inattentive-hyperactive and oppositional behaviors at 4–7 years in the present sample. This suggests that the measures of these putative precursors used in the CNLSY are prospectively related to conduct problems at 8–13 years in the much same way as the various measures of these constructs used in other studies. When conduct problems at 4–7 years were added as a simultaneous predictor to the prediction models, however, inattentive-hyperactive and oppositional behaviors at 4–7 years each accounted for very little independent variance in later conduct problems. This suggests that knowing that young children exhibit inattentive-hyperactive or oppositional behaviors adds very little to our ability to predict conduct problems in middle and late childhood after early conduct problems are considered.

In addition, the longitudinal multivariate behavior genetic analyses revealed that the same genetic and environmental factors that influence conduct problems at both 4–7 and 8–13 years also influence the putative early precursors at 4–7 years. After the genetic and environmental influences that the putative precursors share with early conduct problems (genetic factor A_1) were taken into account, however, inattentive-hyperactive and oppositional behavior each shared little or no unique genetic influences with later conduct problems (genetic factor A_2). This suggests that, after early conduct problems are considered, early inattentive-hyperactive and oppositional behavior do not function as useful early manifestations of additional genetic and environmental influences that later give rise to conduct problems in middle and late childhood.

These findings argue for a reconceptualization of the early childhood manifestations of later conduct problems. Prospective longitudinal studies beginning in early childhood have revealed that young children engage in serious conduct problems to a much greater extent than previously suspected (Keenan et al., 2007; Tremblay, 2004). This is important, as the present findings reveal that the best indicator that a child will exhibit conduct problems at 8–13 years is his or her conduct problems at 4–7 years. Indeed, early inattentive-hyperactive and oppositional behaviors are almost entirely redundant predictors of later conduct problems after early conduct problems are controlled. Thus, inconsistent with their hypothesized role as developmental precursors, early inattentive-hyperactive and oppositional behaviors do not function as informative early behavioral manifestations of future conduct problems after early conduct problems are considered. Similarly, the longitudinal Cholesky models revealed that early inattentive-hyperactive and oppositional behaviors only minimally reflect the same genetic and environmental influences that will give rise to later conduct problems after the genetic and environmental influences on early conduct problems are considered. Thus, little is gained by viewing inattentive-hyperactive and oppositional behaviors as developmental

precursors to later conduct problems. Rather than focusing on early inattentive-hyperactive or oppositional behavior, early interventions designed to prevent conduct problems in late childhood should focus on children who already exhibit conduct problems in early childhood.

This is not to say that inattentive-hyperactive and oppositional behaviors are unimportant, however. Each is a significant dimension of psychopathology in its own right that is associated with different forms of functional impairment (August & Garfinkel, 1993; Lahey, Applegate et al., 2004; Lahey, Pelham et al., 2004). The present findings are consistent with previous cross-sectional studies of shared genetic and environmental influences on ADHD, ODD, and conduct problems (Burt et al., 2005; Dick et al., 2005; Nadder et al., 2002; Waldman et al., 2001) in revealing that inattentive-hyperactive behaviors, oppositional behaviors, and conduct problems at 4–7 years are all influenced to a substantial extent by the same genetic and environmental influences. Nonetheless, Figures 2 and 3 indicate that inattentive-hyperactive and oppositional behaviors each have some unique genetic (a_{22}) and environmental influences (c_{22} for males and a_{22} for both sexes).

Limitations and Future Research

Like most previous longitudinal studies of the predictive associations of inattentive-hyperactive and oppositional behaviors with later conduct problems (Hechtman et al., 1984; Lambert, 1988; Loney et al., 1981; Magnusson, 1984; Mannuzza et al., 1993; Mannuzza et al., 1991), the present analyses were not based on DSM measures of ADHD or ODD. The present findings were quite consistent with the smaller set of previous studies that did use DSM-based measures in terms of the predictive associations found between inattentive-hyperactive and oppositional behaviors and later conduct problems (Biederman et al., 1996; Lahey et al., 2000; Loeber et al., 1995; Rowe et al., 2002), however. This suggests that the present findings are relevant to DSM conceptualizations of these disorders. Nonetheless, it is important to note that different results might have been found had DSM measures been used. For this reason, longitudinal studies of large representative samples using DSM-based measures would be very useful.

It also will be very important for future studies to examine links between early inattentive-hyperactive and oppositional behaviors and delinquency during adolescence, as relations between adolescent delinquency and early problem behaviors could be quite different. In particular, the present findings do not rule out the possibility that inattentive-hyperactive and oppositional behaviors could play roles as developmental precursors to childhood-onset delinquency (Moffitt, 1993, 2006). Because childhood-onset delinquency is characterized by high levels of early child conduct problems, however, it is possible that early inattentive-hyperactive and oppositional behaviors do not independently predict childhood-onset delinquency trajectories after early conduct problems are controlled. Fortunately, it will be possible to test this hypothesis when enough children in the present sample have completed assessments from early childhood through adolescence in the future.

References

- Achenbach TM. The Child Behavior Profile: I. Boys aged 6–11. *Journal of Consulting and Clinical Psychology* 1978;46:478–488. [PubMed: 670491]
- Applegate B, Lahey BB, Hart EL, Biederman J, Hynd GW, Barkley RA, et al. Validity of the age of onset criterion for attention-deficit/hyperactivity disorder: A report from the DSM-IV field trials. *Journal of the American Academy of Child and Adolescent Psychiatry* 1997;36:1211–1221. [PubMed: 9291722]
- August GJ, Garfinkel BD. The nosology of attention-deficit hyperactivity disorder. *Journal of the American Academy of Child and Adolescent Psychiatry* 1993;32:155–165. [PubMed: 8428868]
- Avellar S, Smock PJ. The economic consequences of the dissolution of cohabiting unions. *Journal of Marriage and Family* 2005;67:315–327.

- Biederman J, Faraone SV, Milberger S, Jetton JG, Chen L, Mick E, et al. Is childhood oppositional defiant disorder a precursor to adolescent conduct disorder? Findings from a four-year follow-up of children with ADHD. *Journal of the American Academy of Child and Adolescent Psychiatry* 1996;35:1193–1204. [PubMed: 8824063]
- Burt SA, McGue M, Krueger RF, Iacono WG. Sources of covariation among the child-externalizing disorders: Informant effects and the shared environment. *Psychological Medicine* 2005;35:1133–1144. [PubMed: 16116939]
- Carey G. Cholesky problems. *Behavior Genetics* 2005;35:653–665. [PubMed: 16184491]
- Chase-Lansdale PL, Mott FL, Brooks-Gunn J, Phillips DA. Children of the National Longitudinal Survey of Youth: A unique research opportunity. *Developmental Psychology* 1991;27:918–931.
- D’Onofrio BM, Van Hulle CA, Waldman ID, Rodgers JL, Harden KP, Rathouz PJ, et al. Smoking during pregnancy and offspring externalizing problems: An exploration of genetic and environmental confounds. *Development and Psychopathology* 2008;20:139–164. [PubMed: 18211732]
- D’Onofrio BM, Van Hulle CA, Waldman ID, Rodgers JL, Rathouz PJ, Lahey BB. Causal inferences regarding exposure to prenatal alcohol and childhood conduct problems. *Archives of General Psychiatry* 2007;64:1296–1304. [PubMed: 17984398]
- Dick DM, Viken RJ, Kaprio J, Pulkkinen L, Rose RJ. Understanding the covariation among childhood externalizing symptoms: Genetic and environmental influences on Conduct Disorder, Attention Deficit Hyperactivity Disorder, and Oppositional Defiant Disorder Symptoms. *Journal of Abnormal Child Psychology* 2005;33:219–229. [PubMed: 15839499]
- Dominicus A, Skrondal A, Gjessing HK, Pedersen NL, Palmgren J. Likelihood ratio tests in behavioral genetics: Problems and solutions. *Behavior Genetics* 2006;36:331–340. [PubMed: 16474914]
- Fergusson DM, Horwood LJ. Male and female offending trajectories. *Development and Psychopathology* 2002;14:159–177. [PubMed: 11893091]
- Hechtman L, Weiss G, Perlman T. Hyperactives as young adults: Past and current substance abuse and antisocial behavior. *American Journal of Orthopsychiatry* 1984;54:415–425. [PubMed: 6331769]
- Keenan K, Wakschlag LS, Danis B, Hill C, Humphries M, Duax J, et al. Further evidence of the reliability and validity of DSM-IV ODD and CD in preschool children. *Journal of the American Academy of Child and Adolescent Psychiatry* 2007;46:457–468. [PubMed: 17420680]
- Lahey BB, Applegate B, Waldman ID, Loft JD, Hankin BL, Rick J. The structure of child and adolescent psychopathology: Generating new hypotheses. *Journal of Abnormal Psychology* 2004;113:358–385. [PubMed: 15311983]
- Lahey BB, Loeber R, Quay HC, Applegate B, Shaffer D, Waldman I, et al. Validity of DSM-IV subtypes of conduct disorder based on age of onset. *Journal of the American Academy of Child and Adolescent Psychiatry* 1998;37:435–442. [PubMed: 9549965]
- Lahey, BB.; McBurnett, K.; Loeber, R. Are attention-deficit hyperactivity disorder and oppositional defiant disorder developmental precursors to conduct disorder?. In: Sameroff, A.; Lewis, M.; Miller, S., editors. *Handbook of developmental psychopathology*. 2. New York: Plenum; 2000. p. 431–446.
- Lahey BB, Pelham WE, Loney J, Kipp H, Ehrhardt A, Lee SS, et al. Three-year predictive validity of DSM-IV attention deficit/hyperactivity disorder in children diagnosed at 4–6 years of age. *American Journal of Psychiatry* 2004;161:2014–2020. [PubMed: 15514401]
- Lahey BB, Van Hulle CA, Waldman ID, Rodgers JL, D’Onofrio BM, Pedlow S, et al. Testing descriptive hypotheses regarding sex differences in the development of conduct problems and delinquency. *Journal of Abnormal Child Psychology* 2006;34:737–755. [PubMed: 17033935]
- Lambert NM. Adolescent outcomes for hyperactive children: Perspectives on general and specific patterns of childhood risk for adolescent educational, social, and mental health problems. *American Psychologist* 1988;43:786–799. [PubMed: 3195797]
- Leve LD, Kim HK, Pears KC. Childhood temperament and family environment as predictors of internalizing and externalizing trajectories from ages 5 to 17. *Journal of Abnormal Child Psychology* 2005;33:505–520. [PubMed: 16195947]
- Lilienfeld SO, Waldman ID. The relation between childhood attention-deficit hyperactivity disorder and adult antisocial behavior reexamined: The problem of heterogeneity. *Clinical Psychology Review* 1990;10:699–725.

- Loeber, R. Behavioral precursors and accelerators of delinquency. In: Buikhuisen, W.; Mednick, SA., editors. *Explaining criminal behavior*. Leiden, NE: Brill; 1988. p. 51-67.
- Loeber R, Green SM, Keenan K, Lahey BB. Which boys will fare worse? Early predictors of the onset of conduct disorder in a six-year longitudinal study. *Journal of the American Academy of Child and Adolescent Psychiatry* 1995;34:499–509. [PubMed: 7751264]
- Loeber R, Green SM, Lahey BB, Christ MA, Frick PJ. Developmental sequences in the age of onset of disruptive child behaviors. *Journal of Child and Family Studies* 1992;1:21–41.
- Loehlin JC. The Cholesky approach: A cautionary note. *Behavior Genetics* 1996;26:65–69.
- Loney, J.; Kramer, J.; Milich, RS. The hyperactive child grows up: Predictors of symptoms, delinquency, and achievement at follow-up. In: Gadow, KD.; Loney, J., editors. *Psychosocial aspects of drug treatment for hyperactivity*. Boulder, CO: Westview; 1981. p. 381-415.
- MacCallum RC, Browne MW, Sugawara HM. Power analysis and determination of sample size for covariance structure modelling. *Psychological Methods* 1996;1:130–149.
- Magnusson, D. Henry Tajfel Memorial Lecture. Oxford University; 1984. Early conduct and biological factors in the developmental background of adult delinquency. Cited in Loeber, R. Behavioral precursors and accelerators of delinquency. In: Buikhuisen, W.; Mednick, SA., editors. *Explaining criminal behavior*. Leiden: Brill; 1988. p. 51-67.
- Mannuzza S, Klein RG, Abikoff H, Moulton JL. Significance of childhood conduct problems to later development of conduct disorder among children with ADHD: A prospective follow-up study. *Journal of Abnormal Child Psychology* 2004;32:565–573. [PubMed: 15500034]
- Mannuzza S, Klein RG, Bessler A, Malloy P, LaPadula M. Adult outcome of hyperactive boys: Educational achievement, occupational rank, and psychiatric status. *Archives of General Psychiatry* 1993;50:565–576. [PubMed: 8317950]
- Mannuzza S, Klein RG, Bonagura N, Malloy P, Giampino TL, Addalli KA. Hyperactive boys almost grown up: Replication of psychiatric status. *Archives of General Psychiatry* 1991;48:77–83. [PubMed: 1984764]
- Moffitt TE. Adolescence-limited and life-course-persistent antisocial behavior: A developmental taxonomy. *Psychological Review* 1993;100:674–701. [PubMed: 8255953]
- Moffitt, TE. Life-course persistent versus adolescence-limited antisocial behavior. In: Cicchetti, D.; Cohen, D., editors. *Developmental Psychopathology*. 2. New York: Wiley; 2006. p. 570-598.
- Moffitt TE, Caspi A, Dickson N, Silva P, Stanton W. Childhood-onset versus adolescent-onset antisocial conduct problems in males: Natural history from ages 3 to 18 years. *Development and Psychopathology* 1996;8:399–424.
- Muthén, B.; Muthén, L. Mplus 4.2. Los Angeles: Muthén & Muthén; 2007.
- Nadder TS, Rutter M, Silberg JL, Maes HH, Eaves LJ. Genetic effects on the variation and covariation of attention deficit-hyperactivity disorder (ADHD) and oppositional-defiant disorder/conduct disorder (ODD/CD) symptomalogies across informant and occasion of measurement. *Psychological Medicine* 2002;32:39–53. [PubMed: 11883729]
- Neale, MC.; Cardon, LR. *Methodology for genetic studies of twins and families*. Boston: Kluwer Academic; 1992.
- Peterson JL, Zill N. Marital disruption, parent-child relationships, and behavior problems in children. *Journal of Marriage and the Family* 1986;48:295–307.
- Rodgers, JL.; Johnson, A.; Bard, DE. NLSY-Children/Young Adult (1986–2002) kinship linking algorithm. Norman, OK: University of Oklahoma; 2005.
- Rodgers JL, Rowe DC, Li C. Beyond nature versus nurture: DF analysis of nonshared influences on problem behaviors. *Developmental Psychology* 1994;30:374–384.
- Rowe R, Maughan B, Pickles A, Costello EJ, Angold A. The relationship between DSM-IV oppositional defiant disorder and conduct disorder: Findings from the Great Smoky Mountains Study. *Journal of Child Psychology and Psychiatry* 2002;43:365–373. [PubMed: 11944878]
- Satorra A, Bentler PM. A scaled difference chi-square test statistic for moment structure analysis. *Psychometrika* 2001;66:507–514.
- Satterfield JH, Schell AM. A prospective study of hyperactive boys with conduct problems and normal boys: Adolescent and adult criminality. *Journal of the American Academy of Child and Adolescent Psychiatry* 1997;36:1726–1735. [PubMed: 9401334]

- Tremblay RE. Decade of Behavior Distinguished Lecture: Development of Physical Aggression During Infancy. *Infant Mental Health Journal* 2004;25:399–407.
- Van Hulle CA, Rodgers JL, D’Onofrio BM, Waldman ID, Lahey BB. Sex differences in the causes of self-reported adolescent delinquency. *Journal of Abnormal Psychology* 2007;116:236–248. [PubMed: 17516757]
- Waldman, ID.; D’Onofrio, BM.; Van Hulle, CA.; Rodgers, JL.; Rathouz, PJ.; Lahey, BB. Developmental trends and heterogeneity in genetic and environmental influences on childhood conduct problems in blended and non-blended families. 2008. Manuscript under review
- Waldman, ID.; Rhee, SH.; Levy, F.; Hay, DA. Causes of the overlap among symptoms of ADHD, oppositional defiant disorder, and conduct disorder. In: Levy, F.; Hay, DA., editors. *Attention, genes, and ADHD*. New York: Brunner-Routledge; 2001. p. 115-138.
- Willoughby MT, Curran PJ, Costello EJ, Angold A. Implications of early versus late onset of attention-deficit/hyperactivity disorder symptoms. *Journal of the American Academy of Child and Adolescent Psychiatry* 2000;39:1512–1519. [PubMed: 11128328]
- Zeger SL, Liang KY. Longitudinal data analysis for discrete and continuous outcomes. *Biometrics* 1986;42:121–130. [PubMed: 3719049]

Figure 1.

Illustration of the full Cholesky decomposition model for oppositional behavior at 4–7 years and conduct problems at both 4–7 and 8–13 years. The same model was used for inattentive-hyperactive behavior at 4–7 years, substituting that phenotype for oppositional behavior. Please see text under *Longitudinal Multivariate Behavior Genetic Analyses* for definitions of A_1 – A_3 , C_1 – C_3 , and E_1 – E_3 .

Figure 2.

Squared standardized coefficients for each path in Model 4 for females and males together in a sex-scalar model. Note: Path a_{33} was set to 0 in this model; paths were omitted for each sex if their 95% confidence interval included 0 for that sex. The coefficient for path e_{33} is shown even though its 95% confidence interval included 0 because it includes the estimate of error in conduct problems at 8–13 years. Please see text under *Longitudinal Multivariate Behavior Genetic Analyses* for definitions of A_1 – A_3 , C_1 – C_3 , and E_1 – E_3 .

Figure 3.

Squared standardized coefficients for each path in Model 4 for females (top) and males (bottom). Note: Path a_{33} was set to 0 in this model; paths were omitted for each sex if their 95% confidence interval included 0 for that sex. The coefficient for path e_{33} is shown even though its 95% confidence interval included 0 because it includes the estimate of error in conduct problems at 8–13 years. Please see text under *Longitudinal Multivariate Behavior Genetic Analyses* for definitions of A_1 – A_3 , C_1 – C_3 , and E_1 – E_3 .

Table 1

Descriptive Data

Demographic Characteristics of the Full Sample			
Sex	<u>Females</u>	<u>Males</u>	
	3,194	3,272	
Race-Ethnicity	<u>African American</u>	<u>Hispanic</u>	<u>Non-Hispanic White and Other</u> ¹
	1,972	1,340	3,154
Total family income ²	<u>Mean</u>	<u>Standard Deviation</u>	
	31,789	72,469	

Number of Sibling Pairs of Each Degree of Genetic Relatedness in the Behavior Genetic Analyses				
	<u>r_g</u>	<u>Male Pairs</u>	<u>Female Pairs</u>	<u>Different-sex Pairs</u>
Half Siblings	.25	259	255	572
Ambiguous Siblings	.375	177	193	361
Full Siblings (blended)	.5	190	184	313
Full Siblings (non-blended)	.5	711	644	1422

Intra-Class Correlations Between Siblings on Each Measure of Child Problems Behavior Among Each Sibling Type				
	<u>Half</u>	<u>Ambiguous</u>	<u>Full Blended</u>	<u>Full Non-blended</u>
Hyperactive-Inattentive 4–7	.21	.24	.33	.30
Oppositional-Defiant 4–7	.32	.31	.38	.36
Conduct problems 4–7	.34	.30	.41	.43
Conduct problems 8–13	.36	.37	.42	.46

¹ Almost exclusively non-Hispanic European American, but includes some Asian and Native American children.

² In 1986 dollars.

Means and Standard Deviations of Each Measure of Child Problem Behavior at 4–7 Years and Conduct Problems at 8–13 Years Among Females and Males (N = 6,466)

Table 2

	Males		Females		Tests of sex differences in means ^d		Test of sex differences in variances ^b	
	M	SD	M	SD	M	SD	X ²	F
Conduct Problems (4–7 Years)	2.53	2.07	2.01	1.72			130.51	1.45
Oppositional Behavior (4–7 Years)	1.73	1.39	1.61	1.34			12.87	1.07
Inattentive Hyperactive Behavior (4–7 Years)	1.94	1.41	1.50	1.29			184.82	1.20
Conduct Problems (8–13 Years)	2.47	2.20	1.80	1.74			201.59	1.60

^aTested using Poisson regression; all $p < .001$

^bAll sex differences in variances $p < .001$

Table 3

Formal Tests of Mean Inattentive-Hyperactive Behavior, Oppositional Behavior, and Conduct Problems at 4–7 Years as Predictors of Mean Conduct Problems at 8–13 Years Using Log-Linear Regression in Generalized Estimating Equations

Predictor at 4–7 Years	β	z	$p <$
<u>Separate Models: In Which Each Single Predictor¹ and the Demographic Covariates Were Entered Simultaneously in Three Separate Models</u>			
Inattentive-hyperactive behavior	.34	33.82	.0001
Oppositional behavior	.36	35.74	.0001
Conduct problems	.39	36.74	.0001
<u>Single Model: In Which All Three Early Predictors and Demographic Covariates Were Entered Simultaneously in One Model</u>			
Inattentive-hyperactive behavior	.11	8.84	.0001
Oppositional behavior	.10	7.75	.0001
Conduct problems	.27	20.36	.0001

¹ Each predictor was converted to a standardized z score to facilitate comparisons.

Table 4

Spearman Correlations and Partial Spearman Correlations Between Each Problem Behavior at 4–7 Years and Conduct Problems at 8–13 Years Among Females and Males in the Full Sample

	Spearman correlations (and r^2)	Partial Spearman correlations (and r^2): Demographic covariates	Partial Spearman correlations (and r^2): Demographic covariates and conduct problems at 4–7)
Conduct Problems (4–7 Years)			
Males ^a	.61 ^b (.37)	.58 (.34)	-
Females	.53 (.28)	.52 (.27)	
Oppositional Behavior (4–7 Years)			
Males	.45 (.20)	.45 (.20)	.14 (.02)
Females	.41 (.17)	.41 (.17)	.16 (.02)
Inattentive-Hyperactive Behavior (4–7 Years)			
Males	.48 (.23)	.46 (.21)	.19 (.04)
Females	.41 (.17)	.38 (.14)	.16 (.02)

^a Sex differences in correlations between each measure of early problem behavior were tested by testing the sex-by-early predictor in Poisson regression

^b All correlations and partial correlations significantly different from zero, $p < .0001$

Results of Multivariate Cholesky Decompositions Testing for Sex Differences in the Sharing of Genetic and Environmental Influences Among Oppositional Behavior and Conduct Problems at 4–7 Years and Conduct Problems 8–13 Years

Table 5

	SB χ^2	df	BIC	RMSEA	SB $\Delta\chi^2$	df	P
Model 1 (No Sex Differences)	858.66	282	113544	.068			
Model 2 (Scalar Sex Differences)	574.71	279	113115	.049	100.73	3	<.0001
Model 3 (Quantitative Sex Differences)	567.84	264	113189	.051	18.9	15	n.s.

Notes: SB χ^2 = Satorra-Bentler chi-square; SB $\Delta\chi^2$ = Satorra-Bentler scaled-difference χ^2 ; df = degrees of freedom; BIC = Bayesian information criterion; RMSEA = root mean square error of approximation. Model 1 (No Sex Differences) constrained all parameters to be equal between sexes. Model 2 (Scalar Sex Differences) constrained all parameters to be equal between sexes, but allowed the total variation/covariation in males to be a scalar multiple of the total variation/covariation in females. Model 3 (Quantitative Sex Differences) allows quantitative sex differences in the structure of genetic and environmental influences. The Satorra-Bentler scaled-difference χ^2 is based on a comparison of each model with the model above it.

Results of Multivariate Cholesky Decompositions Testing Specific Parameters in the Sharing of Genetic and Environmental Influences Among Oppositional Behavior and Conduct Problems at 4–7 Years and Conduct Problems 8–13 Years

Table 6

	SB X ²	Df	BIC	RMSEA	Wald Test	df	P
Model 2 (Full Model)	574.71	279	113115	.049			
Model 4 (Drop a ₃₃)	576.77	280	113106	.049	2.63	2	.27
Model 5 (Drop a ₃₃ and a ₃₂)	577.33	281	113098	.049	0.18	2	.92
Model 6 (Drop a ₃₃ and c ₃₂)	576.39	281	113100	.049	0.65	2	.72
Model 7 (Drop a ₃₃ and e ₃₂)	576.71	281	113101	.049	0.76	2	.68

Notes: SB X² = Satorra-Bentler chi-square; SB ΔX² = Satorra-Bentler scaled-difference X²; df = degrees of freedom; BIC = Bayesian information criterion; RMSEA = root mean square error of approximation. All models allow for scalar sex differences. Pathway labels are shown in Figure 1. Model 2 is the full model illustrated in Figure 1. Model 4 is the same as the full model but dropping a₃₃, which was estimated to be 0. The fits of models 5–7 are compared to Model 4. In Model 5, genetic pathways a₃₂ and a₃₃ were dropped. In Model 6, genetic pathway a₃₃ and environmental pathway c₃₂ were dropped. In Model 7, genetic pathway a₃₃ and environmental pathway e₃₂ were dropped. The Satorra-Bentler scaled-difference X² is based on a comparison of each model with Model 4.

Results of Multivariate Cholesky Decompositions Testing for Sex Differences in the Sharing of Genetic and Environmental Influences Among Inattentive-Hyperactive Behavior and Conduct Problems at 4–7 Years and Conduct Problems 8–13 Years

Table 7

	SB χ^2	df	BIC	RMSEA	SB $\Delta\chi^2$	df	p
Model 1 (No Sex Differences)	910.40	282	113938	.083			
Model 2 (Scalar Sex Differences)	631.91	279	113517	.064	101.11	3	<.001
Model 3 (Quantitative Sex Differences)	585.50	264	113550	.061	42.00	15	<.001

Notes: SB χ^2 = Satorra-Bentler chi-square; SB $\Delta\chi^2$ = Satorra-Bentler scaled-difference χ^2 ; df = degrees of freedom; BIC = Bayesian information criterion; RMSEA = root mean square error of approximation. Model 1 (No Sex Differences) constrained all parameters to be equal between sexes. Model 2 (Scalar Sex Differences) constrained all parameters to be equal between sexes, but allowed the total variation/covariation in males to be a scalar multiple of the total variation/covariation in females. Model 3 (Quantitative Sex Differences) allows quantitative sex differences in the structure of genetic and environmental influences. The Satorra-Bentler scaled-difference χ^2 is based on a comparison of each model with the model above it.

Table 8
Results of Multivariate Cholesky Decompositions Testing Specific Parameters in the Sharing of Genetic and Environmental Influences Among Inattentive-Hyperactive Behavior and Conduct Problems at 4–7 Years and Conduct Problems 8–13 Years

	SB χ^2	df	BIC	RMSEA	Wald Test	df	p
Model 3 (Full Model)	585.50	264	113550	.053			
Model 4 (Drop a_{33})	589.75	266	113533	.053	.83	2	.66
Model 5 (Drop a_{32} and a_{33})	593.31	268	113521	.053	3.38	2	.18
Model 6 (Drop a_{33} and c_{32})*	644.43	268	113563	.057	64.0	2	<.001
Model 7 (Drop a_{33} and e_{32})	596.91	268	113530	.053	2.9	2	.22

Notes: SB χ^2 = Satorra-Bentler chi-square; SB $\Delta\chi^2$ = Satorra-Bentler scaled-difference chi-square; df = degrees of freedom; BIC = Bayesian information criterion; RMSEA = root mean square error of approximation. All models allow for quantitative sex differences. Pathway labels are shown in Figure 1. Model 3 is the full model illustrated in Figure 1. Model 4 is the same full model but dropping a_{33} , which was estimated to be 0. The fits of models 5–7 are compared to Model 4. In Model 5, genetic pathways a_{32} and a_{33} were dropped. In Model 6, genetic pathway a_{33} and environmental pathway c_{32} were dropped. In Model 7, genetic pathway a_{33} and environmental pathway e_{32} were dropped.

* A Wald test indicates that dropping c_{32} from the model results in a significant loss in fit.