Published in final edited form as:

Biochem Pharmacol. 2010 October 15; 80(8): 1221–1229. doi:10.1016/j.bcp.2010.06.030.

Identification and Characterization of a Novel Estrogenic Ligand Actinopolymorphol A

Emily Powell¹, Sheng-Xiong Huang², Yong Xu⁴, Scott R. Rajske², Yidan Wang¹, Noel Peters³, Song Guo³, H. Eric Xu⁴, F. Michael Hoffmann^{1,3}, Ben Shen², and Wei Xu¹

¹McArdle Laboratory for Cancer Research, University of Wisconsin-Madison, Madison, WI, USA, 53706

²Division of Pharmaceutical Science, University of Wisconsin-Madison, Madison, WI, USA, 53706

³UWCCC Small Molecule Screening Facility, University of Wisconsin-Madison, Madison, WI, USA, 53706

⁴Laboratory of Structural Sciences, Van Andel Research Institute, 333 Bostwick Avenue, N.E., Grand Rapids, Michigan 49503, USA

Abstract

Xenoestrogenic compounds are abundant in the modern environment including phytoestrogens from plants, chemical by-products from industry, and secondary metabolites from microbes; all can profoundly affect human health. Consequently mechanism-based screens are urgently needed to improve the rate at which the xenoestrogens are discovered. Estrogen Receptor (ER) dimerization is required for target gene transcription. The three ER dimer pairs (ER α / α homodimers, ER β / β homodimers, and ER α / β heterodimers) exhibit diverse physiological responses in response to ligand-dependent activation with ER α / α homodimers being pro-proliferative and ER β / β homodimers being anti-proliferative. The biological role of the ER α / β heterodimer remains unclear. We previously developed a cell-based, bioluminescence resonance energy transfer (BRET) assay that can distinguish natural estrogenic compounds based on their abilities to activate the three diverse ER dimer pairs. Using BRET assays, we sought to identify novel xenoestrogens from soil bacteria that preferentially activate ER α / β heterodimer with hopes of shedding light on the biological function of this elusive dimer pair. Here we describe the application of BRET assays in high throughput screens of crude bacterial extracts not previously screened for ER modulatory function and originating from unique ecological niches. Here we report the discovery and biological evaluation of a new natural product, actinopolymorphol A (1), that preferentially induces ER α / β dimerization. Actinopolymorphol A represents the first representative of a new ER modulatory scaffold.

Keywords

Estrogen receptor α ; estrogen receptor β ; BRET; xenoestrogens; soil bacteria

Address correspondence to: Wei Xu, Ph.D., 1400 University Avenue, McArdle Laboratory Room 421A, Madison, WI 53706, Phone: 608-265-5540, wxu@oncology.wisc.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

The Estrogen Receptors (ERs) are hormone dependent transcription factors existing in two forms: ER α and ER β . The binding of both endogenous (i.e. 17 β -estradiol, also known as E2) and exogenous estrogenic ligands to these receptors induces conformational changes leading to dissociation from the Hsp90 molecular chaperone complex, subsequent receptor dimerization, interaction with coactivator proteins, and recognition of Estrogen Response Elements (EREs) in the promoter regions of target genes to activate target gene transcription. Transcriptional activity of ERs is strongly influenced by ligands at each step including (i) the binding of a given ligand for ER α vs. ER β , (ii) the conformational changes induced upon ligand binding which influence dimer partner preference (i.e. ER α / α and ER β / β homodimers, or ER α / β heterodimers), (iii) cofactor recruitment, and, (iv) interaction with chromatin. The differential regulation of ER α and ER β by endogenous and exogenous estrogenic compounds has extensive physiological implications, as transcriptional activation of ER α by these ligands is known to stimulate cellular proliferation, while transcriptional activation of ER β inhibits cell growth [1-6].

In addition to the genomic transcriptional activities of ER α and ER β in estrogenic signaling, ERs can also be regulated by growth factors such as epidermal growth factor (EGF) and insulin-like growth factor (IGF) [7-10]. The effects of many of these growth factor pathways are believed to reflect their abilities to change the phosphorylation state of ERs, as well as that of coregulators and other proteins with which ERs interact to modulate gene expression. Furthermore, in addition to the well-documented synergistic effects of estrogens and growth factors on gene transcription, estrogens also exert rapid membrane-initiated effects that are known to massively impact cell signaling and may also influence gene transcription in the nucleus. Membrane-bound ERs have been shown to mediate estrogenic effects in ER-negative cells via activation of the MAPK pathway [11]. These non-genomic mechanisms of estrogenic signaling should therefore be carefully considered as important mechanisms of global estrogen action.

The cellular functions of ER α and ER β homodimers are well-established. However, the biological role of the ER α / β heterodimer remains a topic for intense study and debate, due largely to the lack of tools to study ER α / β heterodimerization in a physiological context. The coexpression of ER α and ER β results in a heterogeneous pool of homodimers and heterodimers, and thus the activity of heterodimers cannot be deciphered from that of either homodimer, although evidence strongly suggests that ER β antagonizes the proliferative action of ER α via formation of growth-inhibitory heterodimers [1,3,4,12,13]. To elucidate the biological role of these heterodimers, we have developed novel Bioluminescent Resonance Energy Transfer (BRET) assays in order to study ER α / β heterodimerization in a cell-based, physiological environment in real time [14]. We have used these assays to study the basic intermolecular mechanism of ER α / β heterodimerization. Another important application of the BRET assay involves the identification of selective ER α / β heterodimer-inducing small molecules. Of particular significance here is the application of the BRET assay to identify new natural products able to activate ER α / β heterodimerization.

Exogenous estrogenic ligands are ubiquitous in the natural environment and include xenoestrogenic industrial by-products and phytoestrogens such as genistein, a principle constituent of soy. Metabolites of xenoestrogenic and phytoestrogenic compounds have been demonstrated to be produced by several bacterial strains including those present in the intestinal flora [15-20] and soil bacteria [21,22]. For example, bisphenol A (BPA) can be metabolized by many organisms ranging from microorganisms to animals, and these transformations represent an important pathway for its detoxification [23]. Other studies have shown that many natural products produced by bacteria serve as xenoestrogens [24,25].

Screening of such compounds for their ability to selectively activate ER homodimers and heterodimers is important in order to determine the physiological effects of these environmental ligands as they may act through pro-proliferative ER α / α homodimers or anti-proliferative ER β / β heterodimers. Moreover, the identification of such compounds represents an important undertaking as compounds displaying selective ER activation may serve as scaffolds which may be used for the development of novel therapeutics or biochemical tools. Inspired by the realization that the chemical structures of natural products remain either the source of, or the basis for, the majority of drug discovery and synthesis [26], this study sought to identify new natural products able to induce selective ER heterodimer formation leading to subsequent transcriptional activity with the rationale that these structures may be useful as a basis for chemical synthesis of therapeutically-useful ER dimer-selective ligands. Natural products of interest were produced by actinomycetes of terrestrial origin.

The application of our novel ER dimer-specific BRET assay [14,27] for high throughput screening (HTS) of a microbial library of crude extracts resulted in the identification of actinopolymorphol A from the actinomycete *Actinopolymorpha rutilus* whose structure has not previously been reported or characterized as an ER ligand. This discovery was enabled by the novelty of the BRET assay with its rapid in-cell format which circumvents the need for tissue culture grade crude extracts and serves as an excellent assay for activity-guided chemical fractionation of crude extracts containing an assortment of natural products.

2. Materials and Methods

2.1. High throughput Screening BRET of the UWCCC SMSF Discovery Library

HTS BRET was performed at the University of Wisconsin Small Molecule Screening Facility. ER α / α homodimerization was examined using ER α -RLuc and ER α -YFP, ER β / β homodimerization was examined using RLuc-ER β and YFP-ER β , and ER α / β heterodimerization was examined using ER α -RLuc and YFP-ER β using the optimized conditions described previously [14]. Cells were transfected with these fusion proteins (0.73 μ g RLuc fusion + 2.8 μ g YFP fusion) in batches on 10 cm plates to reduce well-to-well variation in Phenol Red Free DMEM + 5% SFS. Empty vector (pCMX-pL2) and RLuc fusions were also transfected alone in order to calculate the Correction Factor (CF) portion of the BRET ratio [14]. Twenty four hours after transfection, the cells were trypsinized from their 10 cm plates and resuspended to 10,000 cells per well of 384 well white bottom plates in PBS. On each plate, dimer pairs were plated by quadrant (i.e. ER α / α homodimers were plated in quadrant 1, ER β / β homodimers were plated in quadrant 2, and ER α / β heterodimers were plated in quadrants 3 and 4). Thus, all three dimer pairs were present within the same plate in order to avoid confounding plate-to-plate variation. Cells were treated with a final concentration of 5 μ M library compounds for 1 hour, and each condition was performed in triplicate for each compound. The RLuc substrate coelenterazine h was then added to a final concentration of 5 μ M. The RLuc and YFP emission signals were detected at 470 nm and 530 nm, respectively, on a Victor Wallac V plate reader (Perkin Elmer).

2.2. Cell Based assays

2.2.1 Cells and culture—MDA-MB-231 breast cancer cells were purchased from ATCC (cat. no. HTB-26) and were maintained in DMEM + 10% FBS. PC3 human prostate cancer cells were kindly provided by the laboratory of Dr. Douglas McNeel (Department of Medicine, UW Madison) and were maintained in DMEM + 10% FBS, and HC11 normal mouse mammary cells were kindly provided by the laboratory of Dr. Caroline Alexander (Department of Oncology, UW Madison) and were maintained in RPMI1640 + 10 ng/mL EGF, 5 μ g/mL insulin, and 10% FBS.

2.2.2. HEK293 ERE-luciferase reporter assays—HEK293 cells were transfected in batches in 48-well plates using 2.5 ng of each indicated ER and 50 ng tk-ERE-luc vector per well as described above. After allowing 48 hours for protein expression and incubating with the indicated ligands for 24 hours, cells were lysed, and firefly luciferase emission was detected upon addition of the firefly luciferase substrate (Promega) on a PerkinElmer Victor 3-V plate reader using a luminescence detection setting. β -gal was analyzed using the Tropix β -galactosidase detection kit (Tropix), and emission was detected on a PerkinElmer Victor 3-V plate reader using a luminescence detection setting. Luciferase counts were normalized to β -gal counts in each well.

2.2.3. Cell Growth and Viability Assays— 1×10^5 PC3 or HC11 cells were seeded onto 6 cm plates in phenol red free DMEM + 5% SFS and allowed to attach overnight. The next day, media was replaced with media containing the indicated concentration of ligands or 0.1% DMSO, and the total amount of DMSO per plate was kept constant at 0.1%. Time points were harvested at 24 hours, 48 hours, 72 hours, and 96 hours by trypsinizing cells, inactivating the trypsin with DMEM + 10% FBS and transferring to 2 mL eppendorf tubes, and centrifuging at 3000 rpm for 5 minutes. Media + trypsin was then removed from the cell pellets, and the pellet was resuspended in 200 μ L PBS + 200 μ L of a 1:1 dilution of trypan blue and PBS. Cells were then loaded onto cell counting chambers (18 μ L per chamber side) in duplicate and each side was read in triplicate using the Cellometer. The Cellometer protocol was “initial cell type” with a dilution of 2, and cell viability was measured on each read. The cell number per mL was then extrapolated to total cell number based on the total 400 μ L volume.

2.2.4 MTT assays for cellular metabolic activity—This assay measures mitochondrial activity when yellow MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) is reduced to its purple formazan metabolic product [28]. Thus, the ability of a cell to metabolize MTT to formazan is correlated to its metabolic activity. These assays were performed in a variety of cell types following the same protocol. Cells were seeded to a confluency of ~10% on Day 1 in phenol red free DMEM supplemented with 5% FBS stripped 6 times with charcoal and dextran (SFS) in 48-well plates. Cells were allowed to attach overnight, and on Day 2 the appropriate dilution of DMSO vehicle, 10 nM E2, or actinopolymorphol A was added with a final DMSO concentration of 0.1%. Four consecutive time points were then harvested on Days 3, 4, 5, and 6 (24 hours, 48 hours, 72 hours, and 96 hours post-treatment), and ligands were refreshed every 48 hours by replacing the media. Each time point was harvested by adding MTT to a final concentration of 500 μ g/mL and incubating for 30 minutes at 37°C, 5% CO₂ in humidified air. The media + MTT solution was removed from each well with suction and 50 μ L of DMSO was added to each well, incubated at room temperature with shaking to solubilize the purple formazan crystals, and transferred to a clear flat-bottom 96-well plate. The plate was then read at 595 nm for formazan absorbance and 650 nm for background absorbance, and these values were normalized by subtraction. Each condition was performed in triplicate.

2.3. Statistical analysis

T-Tests were employed to statistically analyze data. Comparisons were made between each ligand treatment condition concentration and treatment with the vehicle DMSO.

2.4. Quantitation of proteins and mRNA

2.4.1. Western Blots—Cells were lysed with Triton X-100 lysis buffer with added protease inhibitors and 50 μ g total protein was loaded onto an 8% SDS-PAGE gel. Proteins were transferred to nitrocellulose membranes for 1 hour (for ER α) or 2 hours (for ER β) at

0.35 A. Membranes were blocked overnight in 5% nonfat milk (for ER α) or 10% nonfat milk (for ER β) in PBS Tween. ER α was detected with a 1:10,000 dilution of Santa Cruz HC-20, and ER β was detected with a 1:5000 dilution of Santa Cruz H-150. β -actin was detected with a 1:5000 dilution of Sigma A5441.

2.4.2. RT-PCR—Cellular RNA was isolated using the Trizol extraction method. DNA was digested with DNase I, and RNA was quantitated using the Nanodrop (Thermo Scientific, Wilmington, DE). cDNA was reverse transcribed from RNA using the Superscript II Reverse Transcription kit according to the manufacturer's instructions, and PCR amplification was performed using the following primers: ER α Forward 5' TTATGGAGTCTGGTCCTGTG 3', Reverse 5' CATCATTCCTTCGTAGC 3'; ER β Forward 5' TTTGGGTGATTGCCAAGAGC 3', Reverse 5' AGCACGTGGGCATTTCAGC 3'; β -actin Forward 5'AGGCACCAGGGCGTGATGGT 3', Reverse 5' GGTCTCAAACATGATCTGGG 3'.

2.5. Fluorescence Polarization Assays for measuring ligand binding affinity to ER α and ER β

The binding affinity of ligands for ER α and ER β was measured using Estrogen Receptor Competitor Assays from Invitrogen (PanVera) (Part # P2614, P2698 for ER α ; Part # P2615, P2700 for ER β) according to the manufacturer's instructions. Using purified ER α or ER β provided in the kit, serial dilutions of test compounds were prepared ranging in concentration from 2 mM to 20 pM by preparing 1:10 dilutions in provided screening buffer. The concentration of DMSO was kept below 1%. Compounds were diluted 2-fold in the final reaction with a mixture of 30 nM purified ER α , 2 nM Fluormone, and screening buffer such that the final concentration of test ligand was serially diluted 1 mM to 10 pM, the final concentration of ER α was 15 nM, and the final concentration of Fluormone was 1 nM. Test ligands were prepared in a similar fashion for ER β using a final concentration of 10 nM ER β . This mixture was incubated in the dark at room temperature for 2 hours, and polarization values were read in individual glass tubes using a Beacon 2000 instrument.

2.6. Molecular Modeling

Crystal structure of ER α -estradiol (2OCF.pdb) and ER β -estradiol (2J7X.pdb) complex were retrieved from PDB databank (<http://www.rcsb.org>). The structure of actinopolymorphol A was built and minimized with OPLS 2005 force field in Maestro (Maestro, version 8.5, Schrodinger, LLC, New York, NY, 2008) interface. All the molecular files were prepared by Maestro in Schrodinger program. The grid-enclosing box was centered on the E2 present in the LBD with approximate dimension of 25 $\times$ 25 $\times$ 25 Å³. A scaling factor of 1.0 was set to van der Waals (VDW) radii of those receptor atoms with the partial atomic charge less than 0.25. The minimized actinopolymorphol A was docked into the ligand binding pocket of each ER using Glide (Glide, version 5.0, Schrödinger, LLC, New York, NY, 2008) software package using with standard parameters (SP). All structure figures were prepared using PyMOL (DeLano Scientific).

3. Results

3.1. Identification of ER α / β heterodimer inducing compound actinopolymorphol A by high throughput screening

3.1.1. High Throughput BRET Screening of the University of Wisconsin Discovery Library (WDL)—In order to identify ligands capable of differentially inducing ER α / α and ER β / β homodimerization and ER α / β heterodimerization, a BRET assay was developed and optimized [14]. Distinct from the existing ER reporter assay, the BRET assay is exquisitely sensitive and allows the formation of different dimer types to be detected

independently, including ER α / α , ER β / β , and ER α / β dimers. This is especially important in the case of the ER α / β heterodimer, as the co-expression of ER α and ER β allows the formation of all three dimer pairs, which prevents a clearly delineated understanding of heterodimer function *in vivo*. The BRET assay allows the visualization of ER α / β heterodimers and downstream function without interference from either homodimer. Moreover, because the BRET assay takes place in a physiological environment, it allows selection for small molecules that penetrate into the appropriate intracellular compartments. This assay involved the transfection of DNA encoding an ER α -Renilla Luciferase (RLuc) fusion protein and an ER β -YFP fusion protein into ER-negative HEK293 cells. The high transfectability (>90%), low doubling time, and ER-negative status of this cell line made it an attractive candidate for the high throughput ER BRET screening, as interference from endogenous ERs was not a concern. Cells were transfected with the fusion proteins described in the Methods section in batches in Phenol Red Free DMEM + 5% SFS on 10 cm plates to reduce well-to-well variation. Twenty four hours after transfection, the cells were trypsinized and resuspended to 10,000 cells per well in white 384-well plates. 1 μ L Library extract was then added and incubated with cells for 1 hour, at which point the RLuc substrate coelenterazine h was added to a final concentration of 5 μ M. Coelenterazine h induces RLuc emission at $\sim$ 470 nm; if ER dimerization has occurred, YFP is in close proximity to RLuc, which results in resonance energy transfer to YFP and its emission at 530 nm. Thus, YFP emission is indicative of dimerization (Figure 1). After the addition of coelenterazine h, RLuc and YFP signals were detected at 470 nm and 530 nm, respectively, on a Victor Wallac V plate reader (Perkin Elmer). These values were used to calculate the BRET ratio described previously [14,29]. A schematic of the BRET assay format is shown in Figure 1b. E2 was used as a positive control. Because the ER antagonist ICI 182,780 also induces dimerization [14], vehicle (DMSO) served as the sole negative control. Internal positive and negative controls were included on each plate. HEK293 cells transfected with the ER-RLuc fusion alone (in the absence of YFP) were included on each plate and treated with the vehicle DMSO in order to calculate the Correction Factor portion of the BRET ratio [14]. Each compound was tested in an ER α / α homodimer BRET assay, an ER β / β homodimer BRET assay, and an ER α / β heterodimer BRET assay. On each plate, dimer pairs were plated by quadrant (i.e. ER α / α homodimers were plated in quadrant 1, ER β / β homodimers were plated in quadrant 2, and ER α / β heterodimers were plated in quadrants 3 and 4). Thus, all three dimer pairs were present within the same plate in order to prevent confounding plate-to-plate variation. Each condition was performed in triplicate for each compound. This assay setup allowed verification of the specificity of each primary hit compound based on its ability to activate ER α / β heterodimers in comparison with its ability to activate each respective homodimer.

3.1.2. Rationale for Choice of University of WDL (natural products and microbial extracts library)—The University of Wisconsin is intensely interested in the discovery of new small molecules possessing novel and useful bioactivities. Reflective of this interest, researchers at UW have established a library of small molecules and microbial extracts referred to as the Wisconsin Discovery Library (WDL). The WDL is composed of a wide assortment of privileged natural products and natural product-inspired synthetic compounds in addition to > 1000 natural product extracts from un- and underexplored microorganisms. The molecular diversity displayed by WDL members is extraordinary particularly among the natural product extracts which likely possess heretofore unknown structural scaffolds with, as yet, unidentified bioactivities. For this reason, we applied the BRET assay to a selection of natural product extracts available to us through the WDL. Beyond the molecular diversity presented by natural product extracts, a strong motivator for interest in the screening of natural product extracts is their ready availability via scaled up fermentation of the producing organism and our ability to produce analogs via the

application of combinatorial biosynthesis methods. Finally, it is well established that terrestrial microbes are a rich source for xenoestrogens [21,22,30-33]. This realization supported our hypothesis that crude extracts from the WDL might be an excellent source of ER modulators.

3.1.3. Characterization of Hit Crude Extracts from the WDL—HTS ER α / β heterodimer BRET assays were performed on selected members of the WDL, and crude extracts capable of inducing ER α / β heterodimerization were re-tested alongside each homodimer pair. A total of 25% of the extracts induced ER α / β heterodimerization, and the strongest 1% (10 extracts) were re-tested alongside each homodimer pair. This high number of heterodimer-inducing compounds is likely due to the large size and flexibility of the ER α and ER β LBD, which is accommodating for a variety of chemical structures. Because heterodimerization only was initially tested, this number is not reflective of dimer selectivity. Because of the unrefined and unknown composition of crude extracts, the concentration of specific components could not be determined. Therefore extracts were tested at 1 μ L per well to avoid any toxicity from DMSO. Natural product extract SB83e appeared to exhibit a preference for inducing ER α / β heterodimerization relative to either homodimer pair (Figure 2), and was therefore chosen for subsequent investigation. While a low level of ER α / α homodimerization was also induced by this extract, the crude mixture of multiple compounds did not preclude the possibility of a heterodimer-specific compound coexistent with a dimer non-selective compound.

Bioactivity-guided fractionation of the crude extract SB83e led to the identification of one discrete compound, actinopolymorphol A, responsible for ER α / β heterodimerization observed in BRET assays (Huang et al., submitted for publication). Actinopolymorphol A (Figure 3a, inset) was found to activate the transcriptional activity of ER β alone and ER α + ER β , but was not able to activate the transcriptional activity of ER α alone. The dimer selectivity was subsequently confirmed using the BRET assay in a dose response from 1 μ M to 100 μ M (Figure 3a). These BRET assays showed an optimal 5-fold increase in ER α / β heterodimerization at 100 μ M purified actinopolymorphol A, and this concentration minimally induced either homodimer (less than 1.5 fold). 10 μ M purified actinopolymorphol A likewise induced 3-fold induction for ER α / β heterodimers; at this concentration neither ER α / α or ER β / β homodimerization was induced (Figure 3a). Similarly, both 10 μ M and 100 μ M exhibited a preference for the transcriptional induction of both ER β alone and ER α + ER β , but not ER α alone (Figure 3b). The fold-induction of both the BRET ratio and transcriptional activity of ER α + ER β was greater than that obtained with ER β alone, indicating that this compound retains some level of heterodimer selectivity. Furthermore, this transcriptional activity on an ERE-luciferase reporter was shown to be ER-specific in the presence of the ER pure antagonist ICI 182,780 (Figure 3b). ICI 182,780 completely abrogated actinopolymorphol A-dependent reporter activity in the presence of ER α + ER β (Figure 3b). This is in contrast to the failure of ICI 182,780 to reduce the basal transcriptional activity of ER β homodimer alone (Figure 3b). ER β was previously found to display a high level of ligand-independent dimerization [14] and transcriptional activity [34]. The apparent discrepancy between BRET and reporter assays for ER β / β homodimers at 10 μ M purified actinopolymorphol A is likely due to the lower sensitivity of the ER β / β homodimer BRET assay relative to the reporter assay because the BRET assay only captures receptor dimerization transiently whereas the reporter assay detects accumulated product formation over time. Synthetic actinopolymorphol A was confirmed to retain its ability to selectively induce ER α / β dimerization (Figure 4a) and to enhance transcriptional activity (Figure 4b) of ERs in a fashion comparable to the natural product from *A. rutilus*. Thus, because this compound selectively induced the activity of anti-proliferative ER β / β homodimers and ER α / β heterodimers while not having a pronounced effect on the activity of proliferative ER α / α homodimers, we hypothesized that this compound may be able to

inhibit cell growth by enhancing ER β dimerization leading to a dampening of the proliferative effects of ER α .

3.2. Molecular characterization of actinopolymorphol A

3.2.1. The novel ER heterodimer inducing actinopolymorphol A inhibits cellular proliferation—Synthetic actinopolymorphol A was used in cell proliferation and viability assays in ER α and ER β positive cell lines to determine its effect on cell growth. Figure 5a shows confirmation of ER α and ER β expression in these cell lines by RT-PCR (left panel) and western blotting (right panel). MDA-MB-231 was used as a negative control. Consistent with previous reports, HC11 mouse mammary epithelial cells and PC3 human prostate cancer cells have both ER α and ER β co-expressed [2, 35]. Figure 5b shows the effect of the novel compound in HC11 cells (top panels) and PC3 cells (bottom panels) on cell number (left panels) and viability (right panels). In HC11 cells, a statistically non-significant decrease in cell number was observed in the presence of 10 μ M compound, but these decreases are statistically significant at 100 μ M (p = 0.06 and 0.02, respectively). Neither concentration was generally cytotoxic compared to the vehicle DMSO in this cell line, as observed by trypan blue staining. In PC3 cells, statistically significant decreases in cell number are observed in the presence of both 10 μ M and 100 μ M actinopolymorphol A (p = 0.03 and p =0.002, respectively). Similarly to HC11 cells, neither concentration was generally cytotoxic to PC3 cells. We hypothesize that the difference in the compound's ability to influence cell growth in these two cell lines depends on the relative expression level of ER α and ER β as well as the expression levels of coactivator and corepressor proteins. The effect of actinopolymorphol A on growth and viability cannot be completely reversed by the antagonist ICI 182,780 in ER-positive HC11 and PC3 cells (Figure S1), suggesting that other pathways, including non-genomic estrogenic signaling pathways, could be partially responsible for this compound's effects on cell growth. Thus, actinopolymorphol A is a weak estrogenic ligand which may exert cellular effects through both genomic and non-genomic pathways.

3.2.2. Actinopolymorphol A binds differentially to both ER α and ER β —In order to determine the binding affinity of actinopolymorphol A to ER α and ER β , we employed *in vitro* Fluorescence Polarization (FP) Competition Binding Assays. The basis of this assay lies in the capturing of polarized light in horizontal and vertical planes if a fluorescent ligand remains stationary by binding to a protein during plane-polarized light excitation. A non-fluorescently labeled compound is then titrated to a saturating concentration of fluorescently-labeled ligand, and the degree of competition is measured as the fluorescently-labeled ligand is competed off and freely tumbled during the period of excitation, the emitted light will be random or depolarized. Measurements of the binding affinity of actinopolymorphol A for recombinant ER proteins using this assay were validated commercially [36,37] and in our experiment using the characterized compound genistein (Figure 6a). Genistein has been reported to have a ~10 fold greater binding affinity for ER β over ER α [38], and our results showed a ~15 fold greater binding affinity for ER β (Figure 6a). As shown in Figure 6b, actinopolymorphol A was able to effectively compete with fluorescently labeled E2 for binding to both ER α and ER β with a ~ 2-fold higher affinity for binding to ER β . This data strongly suggests the ability of this compound to bind within the ligand binding domain (LBD) of ER α and ER β , and furthermore, the differential binding affinity likely contributes to the compound's ability to selectively activate ER β / β homodimers relative to ER α / α homodimers. The IC₅₀ values for actinopolymorphol A binding to ER α and ER β were 29 μ M and 15 μ M, respectively, correlating to K_i values of 6.2 μ M and 4.3 μ M, respectively.

3.3. Actinopolymorphol A induces the agonist conformation of both ER α and ER β

Molecular modeling indicated that actinopolymorphol A is well accommodated in the agonist conformation in the LBD of both ER α and ER β (Figure 7). The C-3 hydroxyl group of A-ring of 17 β -estradiol forms strong hydrogen bonds with conserved residues Glu353 and Arg394 from one side of the ligand binding pocket of ER α (Glu260 and Arg301 in ER β) (Figure 7b and 7d). The 17 β -hydroxyl of estradiol makes direct hydrogen bond with the residue His524 on the other side of the binding pocket of ER α (His430 in ER β). The phenolic hydroxyl group of actinopolymorphol A mimics the C-3 hydroxyl group of 17 β -estradiol and participates in the same kind of hydrogen bond interactions observed with residues Glu and Arg in both ERs' binding pockets. However, by virtue of its limited size and simple architecture, actinopolymorphol A cannot extend to the right side of the LBD pocket to form tight interactions with residue His514 in ER α or His430 in ER β , which is likely responsible for the lower docking scores compared to E2. The relatively similar docking scores of actinopolymorphol A to ER α (-8.13) and ER β (-8.02) agree well with our findings of similar K_i values from Fluorescence Polarization competition binding assays. These high docking scores indicate that actinopolymorphol A takes an acceptable low-energy conformation and is accommodated by the physico-chemical environment of the pocket. Thus, based on the binding mode predicted by this molecular modeling, more potent ER dimer-selective estrogenic compounds could be designed based on the structure of this scaffold.

4. Discussion

It is well-established that phytochemicals serve as recruitment signals resulting in the symbiotic activation of plant growth signals by soil bacteria, and that the presence of endocrine disrupting compounds (EDCs) contained in pesticides and other industrial by-products can disrupt this process [38-40]. The direct production of xenoestrogenic compounds by soil bacteria is not as well established and therefore represents an opportunity for discovery of new chemical scaffolds with possible utility as, or the potential for optimization of, ER modulators. Secondary metabolites have been identified through drug discovery methods and include valuable compounds such as antitumor and antibacterial agents. The University of Wisconsin's WDL contains crude bacterial extracts consisting of various natural products and was therefore screened using three highly optimized BRET assays to identify novel inducers of ER α/α and ER β/β homodimerization as well as ER α/β heterodimerization. The application of these BRET assays to screen identify new ER modulators from crude extracts obtained from actinomycetes originating from unique ecological niches resulted in the identification of a novel, previously uncharacterized, dimer selective ER agonist named actinopolymorphol A.

High throughput, mechanism-based assays are on call to advance the discovery of xenoestrogens and drug leads at a rapid pace. Two types of high throughput assays are popularly used for large-scale screening of ER structural scaffolds and agonists or antagonist ligands. A fluorescence polarization (FP) method that measures the capacity of a competitor chemical to displace a high affinity fluorescent E2 from purified, recombinant ER α or ER β have been adapted for testing environmental chemicals for ER binding interactions [36]. However, this method requires pure preparation of receptor, and the fluorescence from the test compounds could interfere with fluorescence readout [36]. Thus far this method has been restricted to use with pure compounds and has not been applicable to whole cell extracts or bioassay-guided fractionation efforts.

Transcriptional reporter assays can be applied to library extracts or compounds but require 18-24 h incubations. Thus, these extracts or compounds must be sterile and of high-quality tissue culture grade to avoid contamination and concomitant ablation of the transcriptional

output signal. The BRET assays described herein circumvent this issue because the library extract or compound in question needs only to be incubated with cells expressing ER fusion proteins for 1 hour to induce dimerization. Furthermore, the three possible ER dimer pairs (ER α / α homodimers, ER β / β homodimers, and ER α / β heterodimers) may be directly examined in parallel yet in isolation from each other, thus providing an added layer of sensitivity and complexity to the library extract or compound's ability to act as an agonist or antagonist ligand. Thus, the utility of this cell-based assay for high throughput crude extract screening lies in its rapid assay time frame. In addition, this method does not require sterile tissue-culture grade extracts. Using this BRET screening method, the crude extract from *A. rutilus* was found to selectively induce formation and transcriptional activity of ER β / β homodimers and ER α / β heterodimers. This screening method allowed assay-guided fractionation of the extract, and the pure compound responsible for induction of dimerization and subsequent transcriptional activity was identified as actinopolymorphol A, a previously unknown natural product. However, estrogenic compounds identified by BRET assays may activate both genomic and non-genomic signaling pathways. Different from the classical genomic signaling through EREs, these non-genomic signaling pathways initiated at the cell membrane may also require receptor dimerization and couples with a variety of other signaling partners that can eventually culminate in the phosphorylation of transcription factors and their partners, ultimately influencing transcriptional outcome, and thus physiological effects such as cell division and apoptosis. Thus, the physiological effects of BRET identified compound await further characterization.

It is worth noting that BRET assays measure the ligand's ability to induce receptor dimerization and the FP method measures ligand replacement of E2 in ER pocket; neither of these assays can distinguish agonists from antagonists. Transcriptional reporter assays measure the ability of the lead compound to induce or inhibit (in the presence of E2) transcription of ER subtypes, allowing determination of agonist or antagonist activity of a ligand. For example, while a low level of ER α / α homodimerization was induced by this compound and substantiated via the BRET assay, these proliferative dimers were not transcriptionally active in ERE-luciferase reporter assays. Because of the limitation of each assay, all three were employed in the present study leading to the discovery of the estrogenic natural product.

The structure of this novel ER dimer-selective natural product from *A. rutilus* was determined by NMR and mass spectroscopic analysis and its absolute stereochemistry was established by total synthesis using an optically pure starting material ((*S*)-2-hydroxy-3-(4-hydroxyphenyl)-propionic acid). (Huang et al., submitted for publication). Combined, the results of structural characterization and coordinated BRET assays reveal that actinopolymorphol A represents a novel scaffold for estrogenic small molecule design. Molecular modeling suggests that, although the phenolic hydroxyl group of actinopolymorphol A mimics the C-3 hydroxyl group of 17 β -estradiol and makes the same hydrogen bond interactions with residues Glu and Arg in both ERs' binding pockets (Figure 7), other structural elements of the natural product do not strictly adhere to predictions likely to be made on the basis of other ER ligands such as tamoxifen or raloxifene. The modeled structure explains how actinopolymorphol A may compete with E2 in binding to the same LBD in FP assay while also displaying a lower binding affinity due to the absence of functionalities needed to H-bond with histidine distal to the Glu and Arg end of the ER LBD. This modeling indicates that the agonist or partial agonist conformation is adopted by ER α and ER β and that their ligand binding cavities are shaped into a low-energy conformation by actinopolymorphol A. Perhaps most significantly, actinopolymorphol A is, to the best of our knowledge, the first ER dimerization modulator identified from actinomycetes. As such, discovery of this natural product and subsequent association with ER modulatory function unveils a new molecular scaffold with a novel and potentially

useful bioactivity. This structure may serve as a molecular scaffold upon which chemical modifications may be made in order to increase the selectivity and efficacy of this novel compound.

The application of ER BRET assays for high throughput screening of crude natural product extracts and subsequent bioassay-guided fractionation leading to the identification of actinopolymorphol A showcases a new molecular scaffold but also highlights the utility of BRET assays in discovering new, otherwise difficult to detect, natural products with ER modulatory activity. These ER modulatory compounds may function through genomic or non-genomic signaling pathways. Follow-up assays showed that actinopolymorphol A is able to act as an agonist on ERs and can decrease the growth of ER α and ER β positive cell lines while not adversely affecting their viability. Despite its low activity on ERs, this novel structure is able to compete with endogenous E2 for LBD binding in both ER α and ER β . Competitive LBD binding by actinopolymorphol A is rationalized on the basis of molecular modeling which suggests the natural product can induce an agonist conformation upon binding to both ERs. Combined, these data reveal the unique application of BRET assays to find new ER modulators and reveal actinomycetes as a potentially rich source of such bioactive natural products, the apparent first example of which, highlights a unique molecular scaffold which may serve as a lead for drug discovery and therapeutic intervention in ER dependent diseases such as breast and prostate cancers.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Erin Shanle for proofreading the manuscript and the Analytical Instrumentation Center of the School of Pharmacy, UW-Madison for support in obtaining MS and NMR data. This work is supported by NIH grants R01CA125387 and R03MH089442 to W.X., T32 CA009135 to E.P. and CA113297 to B.S.

References

1. Chang EC, Frasor J, Komm B, Katzenellenbogen BS. Impact of estrogen receptor beta on gene networks regulated by estrogen receptor alpha in breast cancer cells. *Endocrinology*. 2006; 147:4831–42. [PubMed: 16809442]
2. Helguero LA, Faulds MH, Gustafsson JA, Haldosen LA. Estrogen receptors alpha (ERalpha) and beta (ERbeta) differentially regulate proliferation and apoptosis of the normal murine mammary epithelial cell line HC11. *Oncogene*. 2005; 24:6605–16. [PubMed: 16007178]
3. Liu MM, Albanese C, Anderson CM, Hilty K, Webb P, Uht RM, et al. Opposing action of estrogen receptors alpha and beta on cyclin D1 gene expression. *J Biol Chem*. 2002; 277:24353–60. [PubMed: 11986316]
4. Pettersson K, Delaunay F, Gustafsson JA. Estrogen receptor beta acts as a dominant regulator of estrogen signaling. *Oncogene*. 2000; 19:4970–8. [PubMed: 11042684]
5. Shoker BS, Jarvis C, Sibson DR, Walker C, Sloane JP. Oestrogen receptor expression in the normal and pre-cancerous breast. *J Pathol*. 1999; 188:237–44. [PubMed: 10419589]
6. Tilli MT, Frech MS, Steed ME, Hruska KS, Johnson MD, Flaws JA, et al. Introduction of estrogen receptor-alpha into the tTA/Tag conditional mouse model precipitates the development of estrogen-responsive mammary adenocarcinoma. *Am J Pathol*. 2003; 163:1713–9. [PubMed: 14578170]
7. Kato S, Endoh H, Masuhiro Y, Kitamoto T, Uchiyama S, Sasaki H, et al. Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase. *Science*. 1995; 270:1491–4. [PubMed: 7491495]
8. Aronica SM, Katzenellenbogen BS. Stimulation of estrogen receptor-mediated transcription and alteration in the phosphorylation state of the rat uterine estrogen receptor by estrogen, cyclic

- adenosine monophosphate, and insulin-like growth factor-I. *Mol Endocrinol*. 1993; 7:743–52. [PubMed: 7689695]
9. Ignar-Trowbridge DM, Nelson KG, Bidwell MC, Curtis SW, Washburn TF, McLachlan JA, et al. Coupling of dual signaling pathways: epidermal growth factor action involves the estrogen receptor. *Proc Natl Acad Sci U S A*. 1992; 89:4658–62. [PubMed: 1584801]
 10. Bunone G, Briand PA, Miksicek RJ, Picard D. Activation of the unliganded estrogen receptor by EGF involves the MAP kinase pathway and direct phosphorylation. *Embo J*. 1996; 15:2174–83. [PubMed: 8641283]
 11. Filardo EJ, Quinn JA, Bland KI, Frackelton AR Jr. Estrogen-induced activation of Erk-1 and Erk-2 requires the G protein-coupled receptor homolog, GPR30, and occurs via trans-activation of the epidermal growth factor receptor through release of HB-EGF. *Mol Endocrinol*. 2000; 14:1649–60. [PubMed: 11043579]
 12. Monroe DG, Secreto FJ, Subramaniam M, Getz BJ, Khosla S, Spelsberg TC. Estrogen receptor alpha and beta heterodimers exert unique effects on estrogen- and tamoxifen-dependent gene expression in human U2OS osteosarcoma cells. *Mol Endocrinol*. 2005; 19:1555–68. [PubMed: 15802376]
 13. Stossi F, Barnett DH, Frasor J, Komm B, Lyttle CR, Katzenellenbogen BS. Transcriptional profiling of estrogen-regulated gene expression via estrogen receptor (ER) alpha or ERbeta in human osteosarcoma cells: distinct and common target genes for these receptors. *Endocrinology*. 2004; 145:3473–86. [PubMed: 15033914]
 14. Powell E, Xu W. Intermolecular interactions identify ligand-selective activity of estrogen receptor alpha/beta dimers. *Proc Natl Acad Sci U S A*. 2008; 105:19012–7. [PubMed: 19022902]
 15. Liu X, Tani A, Kimbara K, Kawai F. Metabolic pathway of xenoestrogenic short ethoxy chain-nonylphenol to nonylphenol by aerobic bacteria, *Ensifer* sp. strain AS08 and *Pseudomonas* sp. strain AS90. *Appl Microbiol Biotechnol*. 2006; 72:552–9. [PubMed: 16528514]
 16. Gorbach SL. Estrogens, breast cancer, and intestinal flora. *Rev Infect Dis*. 1984; 6(Suppl 1):S85–90. [PubMed: 6326245]
 17. Xie LH, Ahn EM, Akao T, Abdel-Hafez AA, Nakamura N, Hattori M. Transformation of arctiin to estrogenic and antiestrogenic substances by human intestinal bacteria. *Chem Pharm Bull (Tokyo)*. 2003; 51:378–84. [PubMed: 12672988]
 18. Hori Y, Abe Y, Ezaki M, Goto T, Okuhara M, Kohsaka M. R1128 substances, novel non-steroidal estrogen-receptor antagonists produced by a *Streptomyces*. I Taxonomy, fermentation, isolation and biological properties. *J Antibiot (Tokyo)*. 1993; 46:1055–62. [PubMed: 8360099]
 19. Hori Y, Abe Y, Nishimura M, Goto T, Okuhara M, Kohsaka M. R1128 substances, novel non-steroidal estrogen-receptor antagonists produced by a *Streptomyces*. III Pharmacological properties and antitumor activities. *J Antibiot (Tokyo)*. 1993; 46:1069–75. [PubMed: 8360101]
 20. Hori Y, Takase S, Shigematsu N, Goto T, Okuhara M, Kohsaka M. R1128 substances, novel non-steroidal estrogen-receptor antagonists produced by a *Streptomyces*. II Physico-chemical properties and structure determination. *J Antibiot (Tokyo)*. 1993; 46:1063–8. [PubMed: 8360100]
 21. Kondo H, Nakajima S, Yamamoto N, Okura A, Satoh F, Suda H, et al. BE-14348 substances, new specific estrogen-receptor binding inhibitors. Production, isolation, structure determination and biological properties. *J Antibiot (Tokyo)*. 1990; 43:1533–42. [PubMed: 2276973]
 22. Matseliukh BP, Polishchuk LV, Lutchenko VV, Bambura OI, Kopyiko OP. Synthesis of daidzein and genistein by streptomycetes and their effect on production of antibiotics. *Mikrobiol Z*. 2005; 67:12–21. [PubMed: 16018212]
 23. Kang JH, Katayama Y, Kondo F. Biodegradation or metabolism of bisphenol A: from microorganisms to mammals. *Toxicology*. 2006; 217:81–90. [PubMed: 16288945]
 24. John DM, White GF. Mechanism for biotransformation of nonylphenol polyethoxylates to Xenoestrogens in *Pseudomonas putida*. *J Bacteriol*. 1998; 180:4332–8. [PubMed: 9721266]
 25. Clavel T, Henderson G, Alpert CA, Philippe C, Rigottier-Gois L, Dore J, et al. Intestinal bacterial communities that produce active estrogen-like compounds enterodiol and enterolactone in humans. *Appl Environ Microbiol*. 2005; 71:6077–85. [PubMed: 16204524]
 26. Newman DJ, Cragg GM, Snader KM. Natural products as sources of new drugs over the period 1981-2002. *J Nat Prod*. 2003; 66:1022–37. [PubMed: 12880330]

27. Powell E, Wang Y, Shapiro DJ, Xu W. Differential requirements of Hsp90 and DNA for the formation of estrogen receptor homodimers and heterodimers. *J Biol Chem.* 285:16125–34. [PubMed: 20353944]
28. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods.* 1983; 65:55–63. [PubMed: 6606682]
29. Pfleger KD, Eidne KA. Illuminating insights into protein-protein interactions using bioluminescence resonance energy transfer (BRET). *Nat Methods.* 2006; 3:165–74. [PubMed: 16489332]
30. Wang CL, Takenaka S, Murakami S, Aoki K. Isolation of a benzoate-utilizing *Pseudomonas* strain from soil and production of catechol from benzoate by transpositional mutants. *Microbiol Res.* 2001; 156:151–8. [PubMed: 11572455]
31. Tsai SC, Tsai LD, Li YK. An isolated *Candida albicans* TL3 capable of degrading phenol at large concentration. *Biosci Biotechnol Biochem.* 2005; 69:2358–67. [PubMed: 16377894]
32. Sweetman AJ, Valle MD, Prevedouros K, Jones KC. The role of soil organic carbon in the global cycling of persistent organic pollutants (POPs): interpreting and modelling field data. *Chemosphere.* 2005; 60:959–72. [PubMed: 15992603]
33. Du Y, Zhou M, Lei L. Role of the intermediates in the degradation of phenolic compounds by Fenton-like process. *J Hazard Mater.* 2006; 136:859–65. [PubMed: 16513260]
34. Tremblay A, Tremblay GB, Labrie F, Giguere V. Ligand-independent recruitment of SRC-1 to estrogen receptor beta through phosphorylation of activation function AF-1. *Mol Cell.* 1999; 3:513–9. [PubMed: 10230404]
35. Maruyama S, Fujimoto N, Asano K, Ito A, Usui T. Expression of estrogen receptor alpha and beta mRNAs in prostate cancers treated with leuporelin acetate. *Eur Urol.* 2000; 38:635–9. [PubMed: 11096250]
36. Bolger R, Wiese TE, Ervin K, Nestich S, Checovich W. Rapid screening of environmental chemicals for estrogen receptor binding capacity. *Environ Health Perspect.* 1998; 106:551–7. [PubMed: 9721254]
37. Parker GJ, Law TL, Lench FJ, Bolger RE. Development of high throughput screening assays using fluorescence polarization: nuclear receptor-ligand-binding and kinase/phosphatase assays. *J Biomol Screen.* 2000; 5:77–88. [PubMed: 10803607]
38. Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH, van der Saag PT, et al. Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology.* 1998; 139:4252–63. [PubMed: 9751507]
39. Baker ME. Flavonoids as hormones. A perspective from an analysis of molecular fossils. *Adv Exp Med Biol.* 1998; 439:249–67. [PubMed: 9781308]
40. Peters NK, Frost JW, Long SR. A plant flavone, luteolin, induces expression of *Rhizobium meliloti* nodulation genes. *Science.* 1986; 233:977–80. [PubMed: 3738520]

The abbreviations used are

BRET	bioluminescence resonance transfer
ERα	estrogen receptor α
ERβ	estrogen receptor β
LBD	ligand binding domain
FP	fluorescence polarization
HTS	high throughput screening
NPE	natural product extract
EDC	endocrine disrupting compound
MS	mass spectrum

HPLC high performance liquid chromatography
NMR nuclear magnetic resonance

Figure 1. BRET assay methodology. **(a)** Schematic representing ligand-dependent dimerization and resonance energy transfer between RLuc and YFP fusions via BRET. **(b)** Schematic representing the BRET assay 96-well format protocol.

Figure 2.

BRET HTS identifies one Natural product extract SB83e capable of preferentially inducing anti-proliferative ERβ/β and ERα/β dimers while minimally inducing pro-proliferative ERα/α homodimers. Error bars represent standard deviation from the mean. SB83e 1:10 diluted or non-diluted was incubated with HEK293 cells transfected with different pairs of ER fusion plasmids for one-hour before reading of Rluc and YFP signals.

Figure 3.

Dimer selective natural product extract SB83e was fractionated, and the constituent compound responsible for dimer selectivity was identified and purified. **(a)** BRET assays showing the dimer selectivity of actinopolymorphol A for ER β /ER β and ER α /ER β . **(b)** ERE-luciferase assays in HEK293 cells showing the ability of actinopolymorphol A to induce the transcription of ER α alone (ER α homodimers), ER β alone (ER β homodimers), or ER α + ER β (all three dimer pairs). The transcriptional response is ablated by the ER antagonist ICI 182,780. Error bars represent standard deviations from the mean.

Figure 4. The dimer selectivity for synthetic actinopolymorphol A was confirmed via BRET assays (a) and ERE-luciferase assays in HEK293 cells (b). Error bars represent standard deviations from the mean.

Figure 5.

The synthetic actinopolymorphol A inhibits cell growth. **(a)** ER α and ER β expression in PC3 and HC11 cells was confirmed via semi-quantitative RT-PCR (left panel) and western blotting (right panel). β -actin served as a loading control. **(b)** Cell growth and viability were decreased by treatment with the synthetic actinopolymorphol A in both cell lines as determined by cell counting and trypan blue staining. Error bars represent standard deviations from the mean.

Figure 6. Fluorescence Polarization Competition Binding assays for ER α and ER β . **(a)** Genistein binds to ER α and ER β with different affinities consistent with previous reports and served as a positive control for the Fluorescence Polarization Competition Binding Assay. **(b)** The synthetic actinopolymorphol A binds to ER β with 2-fold higher affinity than to ER α .

Figure 7.

Comparison of predicted binding mode of actinopolymorphol A and binding mode of 17 β -estradiol from crystal structure. A and D, Superimposition of actinopolymorphol A and 17 β -estradiol in the ligand binding domain of ER α (A) and ER β (D). B and E, important interaction between estradiol and residues in the ligand binding pocket of ER α (B) and ER β (E). C and F, Important interactions between actinopolymorphol A and residues in the ligand binding pocket of ER α (C) and ER β (F). The hydrogen bonds were shown as yellow dash line.