Invited reply

Sexual size dimorphism is the most consistent explanation for the body size spectrum of *Confuciusornis sanctus*

Recently, we suggested that the size spectrum of *Confuciusornis sanctus* is best explained by sexual size dimorphism (SSD) and is compatible with a bird-like life history (Peters & Peters 2009). We find the criticism of Chiappe *et al.* (2010) interesting, but irrelevant to our main argument.

Let us follow this argument from the beginning. Chiappe *et al.* (2008) reported a ‘bimodal size distribution’ in *C. sanctus* and hypothesized that it reflected ‘a mid-development phase of exponential growth that separates earlier and later phases of slower growth’ (figure 1*a*). Avian allometry suggests a body mass of approximately 700 g for the larger animals (size class sIII; Peters & Peters 2009) which, according to the hypothesis, were the most mature ones. Mature female birds of this weight produce hatchlings of approximately 30 g, which is an order of magnitude below the estimated body mass of the smaller cohort (sII) in the ‘bimodal distribution’. This gap between hatchlings and sII (figure 1*a*) was unaccounted for by Chiappe *et al.* (2008), who did not consider the egg or neonate masses implied by their hypothesis.

The gap in the hypothesis has no equivalent in the fossil record. Some specimens fall right between the estimated hatchling mass and sII (Peters & Peters 2009), corroborating independently that hatchlings must have been much smaller than sII. To close this gap in the original hypothesis, one may postulate that *C. sanctus* had a biphasic growth curve (figure 1*b*). Alternatively, the size spectrum may be interpreted as resulting from two monophasic growth curves with different asymptotes, as in species exhibiting SSD (figure 1*c*). Both hypotheses explain the size spectrum, but the former (figure 1*b*) postulates a novel, unique type of growth curve, whereas the latter (figure 1*c*) works with growth curves commonly observed. Consequently, we accepted the SSD hypothesis (figure 1*c*), precisely because it does *not* rely on ‘questionable ideas about growth patterns’ (Chiappe *et al.* 2010).

The above is the logical backbone of our argumentation (Peters & Peters 2009). Chiappe *et al.* (2010) do not comment on it. Instead, they state that we interpreted the evidence ‘through the lens of considering the growth pattern of *C. sanctus* as comparable to that

Figure 1. Schematic representation of the size spectrum of *Confuciusornis sanctus* (right in *a–c*) with three alternative life histories (growth curves; left in *a–c*) proposed to explain it. See main text for details.

of modern birds’. This is incorrect; our argument (figure 1) requires no assumption of this type. The SSD hypothesis is the most parsimonious explanation for the observed size spectrum; it also happens to be compatible with bird-like growth. Therefore, as explained by Peters & Peters (2009), the SSD hypothesis resolves the apparent conflict between the size spectrum and bird-like growth that Chiappe *et al.* (2008) had discussed. This resolution, however, is a consequence, not a premise, of the SSD hypothesis.

Chiappe *et al.* (2010) reject our alleged ‘claim that the larger cluster (i.e. sIII) is composed of females’, thus criticizing us for the opposite of what we actually said (‘it remains unclear whether this dimorphism was male or female biased’; Peters & Peters 2009). This misunderstanding is surprising, given that it is the hypothesis of Chiappe *et al.* (2008), not ours, that puts all mature animals into size class sIII which, consequently, must include reproductive females. Obviously, we could not deviate from Chiappe *et al.* (2008) in this point when we evaluated the plausibility of their hypothesis (Peters & Peters 2009; compare second paragraph of this text above).

The accompanying comment can be viewed at <http://dx.doi.org/doi:10.1098/rsbl.2009.1057>.

According to the SSD hypothesis (Peters & Peters 2009), only the smallest birds (sI) were juveniles, while the two larger cohorts (sII, sIII) represented sexually dimorphic adults. Chiappe *et al.* (2010) argue that this conclusion ‘cannot be sustained’ because the deviations of juveniles from allometry (fig. 1 of Peters & Peters 2009) lack statistical significance. This is misleading; the SSD hypothesis does not require juvenile deviation from allometry (fig. 1). Such deviation is an interesting developmental feature, though, and therefore we mentioned it. In the same context, Chiappe *et al.* (2010) state that our ‘inadequate solution’ is ‘compounded by ... *ad hoc* statistics’ because ‘the confidence intervals computed by Peters & Peters (2009) included the smallest individuals ... which indicates that all values can be fitted to a single regression line’. This argument is puzzling since we neither showed confidence intervals in our figures nor mentioned any in our text.

Elongated tail feathers occur in all size classes of *C. sanctus*, including the smallest specimens (Peters & Ji 1999). It is worthwhile remembering that Chiappe *et al.* (2008) rejected SSD as an explanation for the observed size spectrum for this one reason alone, namely the lack of correlation between body size and ‘the most apparent sexual trait’, elongated tail feathers. The assumption that these feathers are a sexual trait, however, is arbitrary (Peters & Peters 2009). Likewise, none of the issues raised by Chiappe *et al.* (2010)

affects the argument on which the SSD hypothesis rests (fig. 1). Thus, the SSD hypothesis remains the most consistent and least speculative interpretation of the currently available evidence.

Winfried S. Peters^{1,*} and Dieter Stefan Peters²

¹*Department of Biology, Indiana/Purdue University Fort Wayne, 2101 East Coliseum Boulevard, Fort Wayne, IN 46805-1499, USA*

²*Forschungsinstitut Senckenberg, Sektion für Ornithologie, Senckenberganlage 25, 60325 Frankfurt am Main, Germany*

*petersw@ipfw.edu

Chiappe, L. M., Marugán-Lobón, J., Ji, S. & Zhou, Z. 2008 Life history of a basal bird: morphometrics of the early cretaceous *Confuciusornis*. *Biol. Lett.* **4**, 719–723. (doi:10.1098/rsbl.2008.0409)

Chiappe, L. M., Marugán-Lobón, J. & Chinsamy, A. 2010 Palaeobiology of the Cretaceous bird *Confuciusornis*: a comment on Peters & Peters (2009). *Biol. Lett.* **6**, 529–530. (doi:10.1098/rsbl.2009.1057)

Peters, D. S. & Ji, Q. 1999 Mußte *Confuciusornis* klettern? *J. Ornithol.* **140**, 41–50. (doi:10.1007/BF02462087)

Peters, W. S. & Peters, D. S. 2009 Life history, sexual dimorphism and ‘ornamental’ feathers in the mesozoic bird *Confuciusornis sanctus*. *Biol. Lett.* **5**, 817–820. (doi:10.1098/rsbl.2009.0574)