Anterior Cruciate Ligament Injuries in the Prepubescent and Adolescent Athlete: Clinical and Research Considerations

Edward M. Wojtys, MD*†; Ashley M. Brower, BS†

*MedSport, †Department of Orthopaedic Surgery, University of Michigan, Ann Arbor

Exercise across the lifespan is an important maintenance program for good health. Particularly in the American population, exercise is essential to prevent such conditions as obesity, diabetes, and cardiac problems, which often result in premature death. However, those who exercise by participating in sports from a young age can suffer debilitating injuries that may curtail their ability to exercise throughout their lives, thus hampering their efforts to stay healthy and avoid disease.

EPIDEMIOLOGY

Data presented at the 2006 American Orthopaedic Society for Sports Medicine seminar on allografts¹ indicated that nearly 350 000 anterior cruciate ligament (ACL) reconstructions are performed annually in the United States; 60 000 of those involve allografts. The annual cost of care is estimated to be more than \$2 billion.² Long-term costs will no doubt be much higher.

Specific to age distribution, the Scandinavian ACL registries³ of 2004–2007 reported a total of 17 632 injuries, with a median age of injury of 25 years. The skeletally immature population appears to be sustaining rising numbers of ACL injuries^{4–6} and higher rates of midsubstance tears.^{3,7–9} The general risk of ACL injury in those with open growth plates is still low, but participation in organized sports significantly increases the risk.¹⁰ In a US report by Shea et al⁶ of 5- to 18-year-old soccer players, 30.8% of all injuries involved the knee, and 6.7% of all injuries were ACL tears. (Statistics were garnered from insurance data documenting 6 million athlete-years.¹¹) In a Finnish population-based cohort study of 14-, 16-, and 18-year-olds (46 472 responding; 78% response rate), 265 cruciate ligament injuries (194 in adolescent boys, 71 in adolescent girls; 92% were ACL tears) reflected an incidence rate of 60.9 per 100 000 person-years.¹⁰ Most interesting were the hazard ratios: 8.5 for adolescent girls and 4.0 for adolescent boys among those participating in organized sports 4 or more times per week.

Why the distribution of ACL injuries in males and females should change with maturation is an interesting research question. Prince et al¹² showed that ACL injuries were more common in immature boys, but after maturation, the risk and incidence in females appeared to increase.^{13,14} During this time, numerous changing factors, including hormones, height, weight, long-bone and muscle

tendon length, and strength distribution, may play a role in injury susceptibility.

In order to implement risk-factor surveillance and prevention programs for ACL injuries, an understanding of the mechanisms involved in ACL failure is essential. Because sports result in so many ACL injuries in adolescents, the mechanisms involved are probably found in common jump-landing and pivoting maneuvers. Although physical contact with another player does play a role in some injuries, most injuries, especially in females, still appear to be noncontact in nature,^{10,15} with up to 70% of ACL injuries in elite females occurring in the absence of contact.¹⁵

ASSOCIATED INJURIES

By itself, injury to the ACL increases the risk of osteoarthritis later in life, but the risk compounds when other structures are injured. Roos¹⁶ reported that if the menisci can be preserved after the ACL injury, the risk of osteoarthritis is reduced. In a retrospective review of 39 patients 14 years of age or younger (30 girls, 9 boys; mean age = 13.6 years; range, 10 to 14 years), Millet et al¹⁷ found that medial meniscus tears were 4 times more common in those whose ACL injuries were treated 6 weeks after injury, suggesting that the time course from injury to treatment may be an important factor in preserving the menisci. Magnetic resonance imaging scans of skeletally immature patients with ACL tears showed that 36% had meniscal tears when the growth plates were wide open, whereas 52% had medial meniscal tears and 17% had lateral meniscal tears when the growth plates were partially closed. These findings suggest a trend toward fewer meniscal injuries in the skeletally immature population.¹²

OUTCOME OF ACL INJURY

Injured athletes retire sooner from sports than their uninjured counterparts.^{18,19} For those with ACL injuries, this may reflect the fact that such injury increases the risk of radiographic osteoarthritis later in life by 105 times.¹⁰ This increased risk of osteoarthritis after ACL injury is an important consideration when deciding whether a nonoperative or operative course of treatment should be undertaken for the skeletally immature patient. Mohtadi and Grant⁵ performed a very complete systematic review on the management of ACL injuries in the skeletally

immature; they reviewed 615 studies but found that only 7 provided adequate comparisons between nonoperative and operative groups. The best evidence available, unfortunately, was no better than level 3. The authors concluded that data were inadequate to determine which route of treatment was better for children and adolescents.⁵

ACL SURGERY

A major concern with early ACL reconstruction in the skeletally immature patient is growth-plate injury from tunnel placement or graft fixation.^{4,20–25} Although the concern is justified, very few reports of growth-plate arrest have been published,^{4,8,26} and growth-plate injury is undoubtedly avoidable with meticulous surgery.^{27–31} Physal anatomy is well known, and experience has shown us that the risks of surgery can be minimized with proper technique.³² The response to ACL surgery, however, is not always predictable: Femoral or tibial overgrowth is possible with stimulation of an active physis secondary to the operative procedure. More problematic can be the occurrence of a growth spurt in close proximity to an ACL reconstruction. This potential for asymmetric long-bone growth may increase, possibly justifying the postponement of ACL reconstruction in some cases.³³

Recent reports by Henry et al²⁹ and Marx et al³⁰ in 2009 were equally optimistic. When comparing early reconstruction in children with delayed reconstruction after skeletal maturity, Henry et al²⁹ noted fewer medial meniscus tears (16% versus 41%) in those children reconstructed initially and no growth-plate injuries after a mean follow-up time of 27 months. Similarly, Marx et al³⁰ monitored 55 children (age range, 8 to 16 years; mean, 13 years) for a mean of 3.2 years after ACL reconstruction (range, 1 to 7.5 years), finding no growth arrests and 88% normal or almost-normal Tegner scores (90 of 100 possible points).

REHABILITATION

Although numerous authors have reported on the success or failure of ACL rehabilitation in adults,^{34–39} few authors have focused on children. Wells et al⁴⁰ reviewed the course of 55 teenagers (40 adolescent girls, 15 adolescent boys; mean, 15.9 ± 1.65 years), showing that the average time from surgery to recovery of 85% of normal quadriceps strength was 5.42 ± 2.27 months. Of these patients, 59% achieved 85% or better strength in fewer than 6 months, with 50% returning to full athletic activity in 6 months. Despite the numerous challenges in the rehabilitation of children and adolescents, ranging from their potentially limited focus and attention span to the adult size of most exercise equipment, successful rehabilitation is possible when managed by those who are familiar with the physical, mental, and emotional challenges of these patient groups. Maintaining motivation over time can be important in enabling them to maximize their great physical potential.

FUTURE RESEARCH CONSIDERATIONS

Clinical

- The most urgent need is to resolve the question of the best clinical treatment for ACL injuries in the skeletally

immature patient. Although surgical reports (especially from the United States) appear promising, surgery is not without risk. Mohtadi and Grant⁵ identified the shortcomings in the current literature and emphasized the need for a randomized trial in which operative and nonoperative groups are matched in terms of age, activity, and other risk factors.

- It is well established that an ACL injury, regardless of treatment, significantly increases the risk of osteoarthritis in adults.¹⁶ It is easy to conclude that the same or perhaps even a worse prognosis would exist for children and adolescents. However, we need to establish whether, in fact, this is true or whether the healing and rehabilitation potential of young people overrides these risks and yields a better prognosis.
- In youngsters who can be successfully treated with a conservative program, is meniscal salvage the key to protecting articular cartilage in the face of a dynamically stable, well-rehabilitated knee? A thorough identification of risk factors and prognostic indicators would help direct parents and physicians in these difficult clinical situations.

Basic Science

- One of the most intriguing questions about the skeletally immature with regard to ligament reconstruction is “Does a ligament grow in a child after implantation and revascularization?” Understanding these mechanisms may prove helpful in providing optimal care at all ages.
- If the prognosis for the skeletally immature with ACL injuries is as poor as in adults with ACL injuries,¹⁶ is there a role for the tissue-engineered repair procedure or a ligament scaffold to augment a repair? Because children appear to have better healing potential than adults, procedures that may not be advisable in adults may be more successful in children.
- Preventing ACL injuries in the young is obviously a high priority to avoid degenerative consequences later in life. An important question remains: Can we successfully conduct risk-factor surveillance in these at-risk groups to identify those who are most vulnerable? Along these same lines, if those with significant risk factors can be identified, should they be advised not to participate in high-risk sports such as soccer and basketball? If the susceptible individuals can be identified, are the injury mechanisms sufficiently well characterized to justify efforts at focused prevention strategies?
- No doubt, the onset of puberty in the skeletally immature patient presents challenges for athletes and appears to influence the injury rate in both boys and girls. Both groups experience a spike in the number of injuries in the early teenage years. Height, weight, muscle performance, and hormones all change during puberty; these alterations in body shape, physiology, and function probably play a role in the increased injury susceptibility. Monitoring the physical development and performance

of children and adolescents may yield many answers for the ACL injury-susceptibility question.

SUMMARY

An ACL injury in a child or adolescent is a major risk factor for the development of early-onset osteoarthritis. Although definitive evidence for recommending surgical management of these injuries is lacking, aggressive surgical treatment appears to provide the best outcome due to meniscal-salvage opportunities, especially in those who remain physically active.

REFERENCES

- Allografts in Sports Medicine: What Do We Know, Need to Know, and Need to Do? Round table discussion: American Orthopaedic Society for Sports Medicine; February 13, 2006; Park City, UT.
- Gottlob CA, Baker CL Jr, Pellissier JM, Colvin L. Cost effectiveness of anterior cruciate ligament reconstruction in young adults. *Clin Orthop Relat Res*. 1999;367(10):272–282.
- Granan LP, Forssblad M, Lind M, Engebretsen L. The Scandinavian ACL registries 2004–2007: baseline epidemiology. *Acta Orthop*. 2009;80(5):563–567.
- Kocher MS, Saxon HS, Hovis WD, Hawkins RJ. Management and complications of anterior cruciate ligament injuries in skeletally immature patients: survey of the Herodius Society and The ACL Study Group. *J Pediatr Orthop*. 2002;22(4):452–457.
- Mohtadi N, Grant J. Managing anterior cruciate ligament deficiency in the skeletally immature individual: a systematic review of the literature. *Clin J Sport Med*. 2006;16(6):457–464.
- Shea KG, Apel PJ, Pfeiffer RP. Anterior cruciate ligament injury in paediatric and adolescent patients: a review of basic science and clinical research. *Sports Med*. 2003;33(6):455–471.
- Kannus P, Jarvinen M. Knee ligament injuries in adolescents: eight-year follow-up of conservative management. *J Bone Joint Surg Br*. 1988;70(5):772–776.
- Lipscomb AB, Anderson AF. Tears of the anterior cruciate ligament in adolescents. *J Bone Joint Surg Am*. 1986;68(1):19–28.
- McCarroll JR, Rettig A, Shelbourne KD. Anterior cruciate ligament injuries in the young athlete with open physes. *Am J Sports Med*. 1988;16(1):44–47.
- Parkkari J, Pasanen K, Mattila VM, Kannus P, Rimpela A. The risk for a cruciate ligament injury of the knee in adolescents and young adults: a population-based cohort study of 46,500 people with a 9-year follow-up. *Br J Sports Med*. 2008;42(6):422–426.
- Shea KG, Pfeiffer R, Wang JH, Curtin M, Apel RJ. Anterior cruciate ligament injury in pediatric and adolescent soccer players: an analysis of insurance data. *J Pediatr Orthop*. 2004;24(6):623–628.
- Prince JS, Laor T, Bean JA. MRI of anterior cruciate ligament injuries and associated findings in the pediatric knee: changes with skeletal maturity. *AJR Am J Roentgenol*. 2005;185(3):756–762.
- Fayad LM, Parellada JA, Parker L, Schweitzer ME. MR imaging of anterior cruciate ligament tears: is there a gender gap? *Skeletal Radiol*. 2003;32(11):639–646.
- Huston LJ, Greenfield ML, Wojtyk EM. Anterior cruciate ligament injuries in the female athlete: potential risk factors. *Clin Orthop Relat Res*. 2000;372:50–63.
- Pasanen K, Parkkari J, Kannus P, et al. Injury risk in female floorball: a prospective one-season follow-up. *Scand J Med Sci Sports*. 2007;18(1):49–54.
- Roos EMR. Joint injury causes knee osteoarthritis in young adults. *Curr Opin Rheumatol*. 2005;17(2):195–200.
- Millet PJ, Willis A, Warren RF. Associated injuries in pediatric and adolescent anterior cruciate ligament tears: does a delay in treatment increase the risk of meniscal tear? *Arthroscopy*. 2002;18(9):955–959.
- Myklebust G, Holm I, Maehleum S, Engebretsen L, Bahr R. Clinical, functional, and radiographic outcome in team handball players 6 to 11 years after anterior cruciate ligament injury: a follow-up study. *Am J Sports Med*. 2003;31(6):981–989.
- Thelin N, Holmberg S, Thelin A. Knee injuries account for the sports-related increased risk of knee osteoarthritis. *Scand J Med Sci Sports*. 2006;16(5):329–333.
- Fehnel DJ, Johnson R. Anterior cruciate ligament injuries in the skeletally immature athlete: a review of treatment outcomes. *Sports Med*. 2000;29(1):51–63.
- Guzzanti V, Falciglia F, Stanitski CL. Preoperative evaluation and anterior cruciate ligament reconstruction for skeletally immature patients in Tanner stages 2 and 3. *Am J Sports Med*. 2003;31(6):949–953.
- Paletta GA Jr. Special considerations: anterior cruciate ligament reconstruction in the skeletally immature. *Orthop Clin North Am*. 2003;34(1):65–77.
- Mylle J, Renders P, Broos P. Transepiphyseal fixation of anterior cruciate avulsion in a child: report of a complication and review of the literature. *Arch Orthop Trauma Surg*. 1993;112(2):101–103.
- Pavlovich R, Goldberg SH, Bach BR Jr. Adolescent ACL injury: treatment considerations. *J Knee Surg*. 2004;17(2):79–93.
- Wester W, Canale ST, Dutkowsky JP, Warner WC, Beaty JH. Prediction of angular deformity and leg-length discrepancy after anterior cruciate ligament reconstruction in skeletally immature patients. *J Pediatr Orthop*. 1994;14(4):516–521.
- Koman JD, Sanders JO. Valgus deformity after reconstruction of the anterior cruciate ligament in a skeletally immature patient: a case report. *J Bone Joint Surg Am*. 1999;81(5):711–715.
- Bisson LJ, Wickiewicz T, Levinson M, Warren R. ACL reconstruction in children with open physes. *Orthopedics*. 1998;21(6):659–663.
- Edwards PH, Grana WA. Anterior cruciate ligament reconstruction in the immature athlete: long-term results of intra-articular reconstruction. *Am J Knee Surg*. 2001;14(4):232–237.
- Henry J, Chotel F, Chouteau J, Fessy MH, Berard J, Moya B. Rupture of the anterior cruciate ligament in children: early reconstruction with open physes or delayed reconstruction to skeletal maturity? *Knee Surg Sports Traumatol Arthrosc*. 2009;17(7):748–755.
- Marx A, Siebold R, Sobau C, Saxler G, Ellermann A. ACL reconstruction in skeletally immature patients [in German]. *Sportverletz Sportschaden*. 2009;23(1):47–51.
- Schneider FJ, Kraus T, Linhart WE. Anterior cruciate ligament reconstruction with semitendinosus tendon in children [in German]. *Oper Orthop Traumatol*. 2008;20(4–5):409–422.
- Anderson AF. Transepiphyseal replacement of the anterior cruciate ligament using quadruple hamstring grafts in skeletally immature patients. *J Bone Joint Surg Am*. 2004;86(suppl 1, pt 2):201–209.
- Stricker SJ, Hunt T. Evaluation of leg length discrepancy in children. *Int J Pediatr*. 2004;19(3):134–142.
- Beynon BD, Uh BS, Johnson RJ, et al. Rehabilitation after anterior cruciate ligament reconstruction: a prospective, randomized, double-blind comparison of programs administered over 2 different time intervals. *Am J Sports Med*. 2005;33(3):347–359.
- Harilainen A, Sandelin J. Post-operative use of knee brace in bone-tendon-bone patellar tendon anterior cruciate ligament reconstruction: 5-year follow-up results of a randomized prospective study. *Scand J Med Sci Sports*. 2006;16(1):14–18.
- Hooper DM, Morrissey MC, Drechsler W, Morrissey D, King J. Open and closed kinetic chain exercises in the early period after anterior cruciate ligament reconstruction: improvements in level walking, stair ascent, and stair descent. *Am J Sports Med*. 2001;29(2):167–174.
- Risberg MA, Holm I, Myklebust G, Engebretsen L. Neuromuscular training versus strength training during first 6 months after anterior cruciate ligament reconstruction: a randomized clinical trial. *Phys Ther*. 2007;87(6):737–750.
- Shaw T, Williams MT, Chipchase LS. Do early quadriceps exercises affect the outcome of ACL reconstruction? A randomised controlled trial. *Aust J Physiother*. 2005;51(1):9–17.
- Tagesson S, Öberg B, Good L, Kvist J. A comprehensive rehabilitation program with quadriceps strengthening in closed versus

open kinetic chain exercise in patients with anterior cruciate ligament deficiency: a randomized clinical trial evaluating dynamic tibial translation and muscle function. *Am J Sports Med.* 2008;36(2):298–307.

40. Wells L, Dyke JA, Albaugh J, Ganley T. Adolescent anterior cruciate ligament reconstruction: a retrospective analysis of quadriceps strength recovery and return to full activity after surgery. *J Pediatr Orthop.* 2009;2(5):486–489.

Editor's note: Edward M. Wojtys, MD, is a professor in the Department of Orthopaedic Surgery; Chief of the Sports Medicine Service; Medical Director of the MedSport Sports Medicine Program; and Associate Director, Bone & Joint Injury Prevention & Rehabilitation Center, University of Michigan, Ann Arbor. Ashley M. Brower, BS, is a research associate at the University of Michigan Health System. Address correspondence to Edward M. Wojtys, MD, The University of Michigan, 24 Frank Lloyd Wright Drive, Ann Arbor, MI 48106-0391. Address e-mail to edwojtys@umich.edu.