Published in final edited form as:

Angew Chem Int Ed Engl. 2010 September 3; 49(37): 6589–6591. doi:10.1002/anie.201002972.

Chiral Phosphoric Acid Catalyzed Peroxidation of Imines**

Wenhua Zheng,

Department of Chemistry, University of South Florida 4202 E. Fowler Avenue, CHE 205A, Tampa, FL 33620 (USA)

Lukasz Wojtas, and

Department of Chemistry X-Ray Facility University of South Florida, Tampa, FL 33620 (USA)

Jon C. Antilla*

Department of Chemistry, University of South Florida 4202 E. Fowler Avenue, CHE 205A, Tampa, FL 33620 (USA)

Keywords

asymmetric catalysis; imines; organocatalysis; peroxides; phosphoric acids

Peroxide-containing compounds are a class of biologically important targets, many of which possess antitumor, anti-bacterial, and antimalarial activities.[1] For example, the peroxide-containing natural product artemisinin (qinghaosu) has been shown to be a highly effective antimalarial drug.[2] The chiral peroxide moiety in this natural product was proven to be essential for the high antimalarial potency.[2] For this reason, the development of stereoselective methods and strategies for the preparation of chiral peroxides has become considerably important to the practitioners of chemical and pharmaceutical synthesis. However, catalytic syntheses of chiral peroxides is still a challenging goal to synthetic chemists.[3–5] Despite these challenges, the groups of Deng[6] and List[7] independently reported the use of a chiral base-catalyzed enantioselective peroxidation of α,β -unsaturated ketones in 2008.

Chiral α -amino peroxide moieties, though somewhat rare, can be found in natural products. For example, verruculogen (Figure 1) contains an α -amino peroxide, and has been found to alter GABA receptor binding and inhibit the mammalian cell cycle.[1a] In addition, dioxetanone is a high-energy natural product containing a chiral α -amino peroxide found in the Japanese firefly.[8] To the best of our knowledge, there is no catalytic asymmetric method that can give direct access to chiral α -amino peroxides.[9]

Chiral phosphoric acids have proven to be efficient catalysts for a large number of important enantioselective transformations.[10–11] So far most of the phosphoric acid catalyzed reactions have utilized carbon nucleophiles. Recently, our group and others reported on the addition of heteroatoms such as amines and alcohols, to imines catalyzed by chiral phosphoric acids.[12] Herein, we report the first example of a chiral phosphoric acid catalyzed addition

[**]We thank the National Institutes of Health (NIH GM-082935) and the National Science Foundation (NSF-0847108) for financial support.

© 2010 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

[*]Fax: (+ 1) 813-974-1733 jantilla@usf.edu.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201002972>.

of hydroperoxides to imines, thereby allowing the direct access to chiral α -amino peroxides in high yield and enantiomeric excess (Scheme 1).

Initial studies were conducted by examining the addition of *tert*-butyl hydroperoxide to imine **2a** in the presence of a chiral phosphoric acid. Catalyst screening revealed that phosphoric acid **1** was the most effective catalyst for the peroxidation of imine **2a** in terms of enantioselectivity. [13] With toluene as the solvent, the product was obtained in 88% yield and 75% *ee* (Table 1, entry 1). The reaction could also be conducted in CH₂Cl₂, THF, and EtOAc to give the desired product in high yields (Table 1, entries 2–4). Isopropyl acetate, a solvent having a variety of uses in manufacturing, was found to be the best with respect to asymmetric induction (Table 1, entry 5). To additionally optimize the reaction conditions, variation of protecting group on the nitrogen atom of the imine was investigated. As shown in Table 1, the enantiomeric excess increased with introduction of a methoxy group in the *para* position of phenyl ring (Table 1, entry 6). Placement of the same substituent in the *meta* position gave an even higher *ee* value (Table 1, entries 7–8). Installation of electron-rich groups in the *meta* position of the phenyl ring enhanced the *ee* value to 97% (Table 1, entry 9) and 96% (Table 1, entry 10). Two equivalents of *tert*-butyl hydroperoxide were used to ensure reaction completion and did not have a detrimental effect on the enantioselectivity or reaction efficiency. Notably, prolonged reaction times did not lead to a decrease in optical purity. This suggests that retro-addition leading to subsequent product racemization through an iminium ion was not observed.

With the optimized conditions in hand, we started to investigate the substrate scope for the asymmetric peroxidation of imines. The representative results are summarized in Scheme 2. To our delight, catalyst **1** gave excellent results for the reaction of a wide range of different imines with hydroperoxides. With 3,5-dimethoxybenzoyl as a protecting group, electron-withdrawing substituents (F, Cl, and Br) in the *para* position (**4g**, **4h**, and **4i**) were all shown to be excellent substrates for the peroxidation reaction. Likewise, electron-donating substituents in the *para* and *ortho* positions (**4j** and **4k**) also provided excellent asymmetric induction. With 3-methoxybenzoyl as the protecting group, substrates having electron-withdrawing and electron-donating groups led to highly enantioselective products (**4m** and **4n**). When cumene hydroperoxide was used as a nucleophile, products **4p**, **4q**, and **4r** were obtained with high enantioselectivity.

The absolute configuration of the peroxide product was determined to be (*R*)-**4i** by single-crystal X-ray diffraction analysis of compound **4i** (Figure 2). [14] Configurations of other products were assigned by analogy.

We believe the bifunctional nature of the chiral phosphoric acid is responsible for concurrent activation of both the nucleophile and the electrophile through hydrogen-bonding interactions (Figure 3). These hydrogen-bonding interactions presumably serve to create a chiral environment, which allows for high selectivity during the addition of the peroxide to the activated imine.

In summary, we have successfully discovered a novel catalytic enantioselective peroxidation of imines using a chiral phosphoric acid. This approach provides direct access to optically active α -amino peroxides with high enantioselectivities. Additional investigation into the expansion of the reaction scope and synthetic utility, with the aim of developing more enantioselective transformations (novel chiral oxidizing reagents), reaction methodologies, and applications to total synthesis, are currently underway in our laboratory and will be reported in due course.

Experimental Section

General procedure: The imine **2** (0.1 mmol) and catalyst (*R*)-**1** (5 mol %) were placed in a flame-dried reaction tube. Dry isopropyl acetate (0.6 mL) was added to the mixture. The reaction mixture was stirred for 5 min, and then hydroperoxide **3** (0.2 mmol) was added and the mixture was stirred for an additional 24 h at RT. The reaction mixture was directly subjected to silica gel column chromatography (hexanes/ethyl acetate 10:1) to give the pure product **4**.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

- [1]. For reviews, see: a Casteel DA. *Nat. Prod. Rep* 1999;16:55–73.; b Casteel DA. *Nat. Prod. Rep* 1992;9:289–312. [PubMed: 1522977] ; c Rappoport, Z. *The Chemistry of Peroxide*. Wiley; New York: 2006. p. 915–1000.
- [2] a). Wu Y. *Acc. Chem. Res* 2002;35:255–259. [PubMed: 12020162] b Posner GH, O'Neil PM. *Acc. Chem. Res* 2004;37:397–404. [PubMed: 15196049]
- [3]. For selected examples of chiral epoxide syntheses, see: a O'Neil PM, Searle NL, Raynes KJ, Maggs JL, Ward SA, Storr RC, Park BK, Posner GH. *Tetrahedron Lett* 1998;39:6065–6068.; b Dussault PH, Eary CT, Woller KR. *J. Org. Chem* 1999;64:1789–1797. [PubMed: 11674266] ; c Dussault PH, Trullinger TK, Noor-e-Ain F. *Org. Lett* 2002;4:4591–4593. [PubMed: 12489937] ; d Kawanishi M, Kotoku N, Itagaki S, Horii T, Kobayashi M. *Bioorg. Med. Chem* 2004;12:5297–5307. [PubMed: 15388157]
- [4]. For a kinetic resolution of a racemic peroxide, see: Driver TG, Harris JR, Woerpel KA. *J. Am. Chem. Soc* 2007;129:3836–3837. [PubMed: 17358067]
- [5]. For copper-catalyzed allylic and benzylic peroxidation, see: Schulz M, Kluge R, Gelalcha FG. *Tetrahedron: Asymmetry* 1998;9:4341–4360.
- [6]. Lu X, Liu Y, Sun B, Cindric B, Deng L. *J. Am. Chem. Soc* 2008;130:8134–8135. [PubMed: 18529055]
- [7]. Reisinger CM, Wang X, List B. *Angew. Chem* 2008;120:8232–8235. *Angew. Chem. Int. Ed* 2008;47:8112–8115.
- [8]. Chung LW, Hayashi S, Lundberg M, Nakatsu T, Kato H, Morokuma K. *J. Am. Chem. Soc* 2008;130:12880–12881. [PubMed: 18767834]
- [9]. For syntheses of chiral α -acylaminoperoxides and hydroperoxides utilizing stoichiometric chiral reagents or mediators, see: a Rebek J, Mc Cready R. *J. Am. Chem. Soc* 1980;102:5602–5605.; b Rebek J. *Heterocycles* 1981;15:517–545.; c Schmidt U, Hausler J. *Angew. Chem* 1976;88:538–539.; *Angew. Chem. Int. Ed. Engl* 1976;15:497–498. [PubMed: 822758] ; d Machin PJ, Sammes PG. *J. Chem. Soc. Perkin Trans. 1* 1976:628–634.; e Markham JL, Sammes PG. *J. Chem. Soc. Perkin Trans. 1* 1979:1885–1888.; f Kienle M, Argyrakis W, Baro A, Laschat S. *Tetrahedron Lett* 2008;49:1971–1974.; g Blumenthal H, Liebscher J. *ARKIVOC* 2009;xi:204–220.
- [10]. For reviews of chiral phosphoric acids, see a Akiyama T. *Chem. Rev* 2007;107:5744–5758. [PubMed: 17983247] ; b Doyle AG, Jacobsen EN. *Chem. Rev* 2007;107:5714–5743.; c Terada M. *Chem. Commun* 2008:4097–4112.
- [11]. For selected examples of chiral phosphoric acid catalyzed organic transformations, see: a Uruguchi D, Terada M. *J. Am. Chem. Soc* 2004;126:5356–5357. [PubMed: 15113196] ; b Akiyama T, Itoh J, Yokota K, Fuchibe K. *Angew. Chem* 2004;116:1592–1594.; *Angew. Chem. Int. Ed* 2004;43:1566–1568.; c Kang Q, Zhao Z-A, You S-L. *J. Am. Chem. Soc* 2007;129:1484–1485. [PubMed: 17283980] ; d Wang X, List B. *Angew. Chem* 2008;120:1135–1138.; *Angew. Chem. Int. Ed* 2008;47:1119–1122.; e Cheng X, Goddard R, Buth G, List B. *Angew. Chem* 2008;120:5157–5159.; *Angew. Chem. Int. Ed* 2008;47:5079–5081.; f Cheon CH, Yamamoto H. *J. Am. Chem. Soc* 2008;130:9246–9247. [PubMed: 18582053] ; g Rueping M, Antonchick AP. *Angew. Chem* 2008;120:5920–5922.; *Angew. Chem. Int. Ed* 2008;47:5836–5838.; h Hu W, Xu X, Zhou J, Liu W-J, Huang H, Hu J, Yang L, Gong L-Z. *J. Am. Chem. Soc* 2008;130:7782–7783. [PubMed:

18512907] ; i Sickert M, Schneider C. *Angew. Chem* 2008;120:3687–3690.; *Angew. Chem. Int. Ed* 2008;47:3631–3634.; j Xu S, Wang Z, Zhang X, Zhang X, Ding K. *Angew. Chem* 2008;120:2882–2885.; *Angew. Chem. Int. Ed* 2008;47:2840–2843.; k Guo Q-S, Du D-M, Xu J. *Angew. Chem* 2008;120:771–774.; *Angew. Chem. Int. Ed* 2008;47:759–762.; l Liu H, Dagousset G, Masson G, Retailleau P, Zhu J. *J. Am. Chem. Soc* 2009;131:4598–4599. [PubMed: 19334771] ; m Terada M, Tanaka H, Sorimachi K. *J. Am. Chem. Soc* 2009;131:3430–3431. [PubMed: 19231855] ; n Terada M, Toda Y. *J. Am. Chem. Soc* 2009;131:6354–6355. [PubMed: 19374414] ; o Momiyama N, Tabuse H, Terada M. *J. Am. Chem. Soc* 2009;131:12882–12883. [PubMed: 19694430] ; p Lu M, Zhu D, Lu Y, Zeng X, Tan B, Xu Z, Zhong G. *J. Am. Chem. Soc* 2009;131:4562–4563. [PubMed: 19290662] ; q Han Z-Y, Xiao H, Chen X-H, Gong L-Z. *J. Am. Chem. Soc* 2009;131:9182–9183. [PubMed: 19518048] ; r Akiyama T, Katoh T, Mori K. *Angew. Chem* 2009;121:4290–4292.; *Angew. Chem. Int. Ed* 2009;48:4226–4228.; s Mori K, Katoh T, Suzuki T, Noji T, Yamanaka M, Akiyama T. *Angew. Chem* 2009;121:9832–9834.; *Angew. Chem. Int. Ed* 2009;48:9652–9654.; t Yue T, Wang M-X, Wang D-X, Masson G, Zhu J. *Angew. Chem* 2009;121:6845–6849.; *Angew. Chem. Int. Ed* 2009;48:6717–6721.; u Gu Q, Rong Z-Q, Zheng C, You S-L. *J. Am. Chem. Soc* 2010;132:4056–4057. [PubMed: 20205477] ; v Liao S, List B. *Angew. Chem* 2010;122:638–641.; *Angew. Chem. Int. Ed* 2010;49:628–631.

[12] a). Rowland GB, Zhang H, Rowland EB, Chennamadhavuni S, Wang Y, Antilla JC. *J. Am. Chem. Soc* 2005;127:15696–15697. [PubMed: 16277499] b Liang Y, Rowland EB, Rowland GB, Perman JA, Antilla JC. *Chem. Commun* 2007:4477–4479. c Li G, Fronczek FR, Antilla JC. *J. Am. Chem. Soc* 2008;130:12216–12217. [PubMed: 18722443] d Cheng X, Vellalath S, Goddard R, List B. *J. Am. Chem. Soc* 2008;130:15786–15787. [PubMed: 18975905] e Rueping M, Antonchick AP, Sugiono E, Grenader K. *Angew. Chem* 2009;121:925–927. *Angew. Chem. Int. Ed* 2009;48:908–910.

[13]. Please see the Supporting Information for a table of chiral phosphoric acids that were screened.

[14]. CCDC 775266 ((*R*)-**4i**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif

Figure 1.
Natural products possessing a chiral α -amino peroxide.

Figure 2. ORTEP representation of the X-ray structure of Compound (*R*)-**4i**. The thermal ellipsoids are shown at 50% probability.

Figure 3.
Proposed transition state for the chiral phosphoric acid catalyzed peroxidation of imines.

Scheme 1.
Catalytic synthesis of optically active α -amino peroxides.

Scheme 2.

Substrate scope for the chiral phosphoric acid catalyzed enantioselective peroxidation of imines. Reaction conditions: imine **2g-o** (0.1 mmol), hydroperoxide **3a-b** (0.2 mmol), **(R)-1** (5 mol%), *i*PrOAc (0.6 mL) at room temperature. The reported yields are for the isolated product, and the reported *ee* values were determined by HPLC analysis using a chiral stationary phase.

Table 1Optimization of reaction conditions.^[a]

Entry	Imine (R ¹)	Solvent	Yield [%] ^[b]	ee [%] ^[c]
1	2a (Ph)	toluene	4a : 88	75
2	2a	CH ₂ Cl ₂	4a : 88	79
3	2a	THF	4a : 86	47
4	2a	EtOAc	4a : 93	84
5	2a	<i>i</i> PrOAc	4a : 84	86
6	2b (4-MeOC ₆ H ₄)	<i>i</i> PrOAc	4b : 83	90
7	2c (2-MeC ₆ H ₄)	<i>i</i> PrOAc	4c : 92	84
8	2d (3-MeC ₆ H ₄)	<i>i</i> PrOAc	4d : 83	88
9	2e (3-MeOC ₆ H ₄)	<i>i</i> PrOAc	4e : 82	97
10	2f (3,5-(MeO) ₂ C ₆ H ₃)	<i>i</i> PrOAc	4f : 81	96

^[a] Reaction conditions: imine **2a–f** (0.1 mmol), hydroperoxide **3a** (0.2 mmol), (*R*)-**1** (5 mol%), solvent (0.6 mL) at room temperature.

^[b] Yield of isolated product.

^[c] Determined by HPLC analysis using a chiral stationary phase.