Published in final edited form as:

Cancer Prev Res (Phila). 2010 September ; 3(9): 1060–1065. doi:10.1158/1940-6207.CAPR-10-0175.

Metformin and Other Biguanides in Oncology: Advancing the Research Agenda

Michael Pollak

Department of Oncology, McGill University and Jewish General Hospital, Montreal, Quebec, Canada

Abstract

Retrospective studies suggest that diabetics treated with metformin have a substantially reduced cancer burden compared with other diabetics, studies that may be impractical to confirm prospectively. It is unclear if this reflects a chemopreventive effect, an effect on transformed cells, or both. It also remains to be established if these data have relevance to people without diabetes. Laboratory models, however, provide independent impressive evidence for activity of metformin and other biguanides in both cancer treatment and chemoprevention. Investigations of biguanide mechanisms have revealed considerable complexity and have identified important gaps in knowledge that should be addressed to ensure the optimal design of clinical trials of these agents. Such trials may define important new indications of biguanides for the prevention and/or treatment of many common cancers.

Introductory History

The fascinating story leading to current interest in the use of metformin for cancer prevention and treatment began in medieval Europe. Herbalists of that time recognized that polyuria (excessive urination) could occasionally be alleviated by *Galega officinalis* (French lilac). Nothing was known concerning the active ingredient of the plant. Even the disease responsible for polyuria that responded to *Galega officinalis*, diabetes mellitus, had not been defined. In the modern era, active fractions of the natural product were identified, leading to drug development efforts. After several false starts, the biguanides metformin, phenformin, and buformin were developed. Clinical experience showed that metformin has an excellent therapeutic index for diabetes, for which the drug was introduced in Europe in the 1970s and approved in the U.S. in 1995 (1).

Clinical trials demonstrated that metformin has advantages over other agents used in the management of type II diabetes (2). This finding, in conjunction with the global diabetes epidemic, has made metformin one of the most frequently prescribed drugs on the planet, with an estimated 40 million prescriptions filled in the U.S. alone in 2008. Its safety has allowed further clinical trials, which documented effectiveness in other conditions including weight gain induced by antipsychotic medications (3), polycystic ovary disease (4), and diabetes prevention for subjects at a high risk (5). Early laboratory studies of biguanides explored antimicrobial activity, and biguanide derivatives are now widely used in applications ranging from swimming pool decontamination to contact-lens sterilization and are under study for

Corresponding Author: Michael Pollak, M.D., 3755 Cote-Ste.-Catherine, Montreal, Quebec, H3T 1E2, Canada. Phone: 514-340-8222 ext 4139; Fax: 514-340-8600, michael.pollak@mcgill.ca.

Disclosure of Potential Conflicts of Interest The author has consulted for Merck, Novo-Nordisk, Lilly, Pfizer, and Sanofi-Aventis.

AIDS treatment (6). This review outlines the basis for current interest in potential applications of biguanides in oncology.

Mechanism of Biguanide Action for Current Indications

The precise mechanism of action of biguanides in therapy of type II diabetes remains an active research topic. As these compounds lower both the hyperglycemia and the hyperinsulinemia associated with this disease, early investigations explored the concept that biguanides enhance insulin-receptor activation and downstream signaling, thus reducing insulin resistance, which is a key aspect of type II diabetes pathophysiology. More-recent work showed that biguanides impair mitochondrial adenosine-5'-triphosphate (ATP) production (7,8), which results in activation of the liver kinase B1 (LKB1)-5' adenosine monophosphate-activated protein kinase (AMPK) signaling pathway. This pathway is central to the regulation of cellular energy homeostasis, and its activation under conditions of energy stress leads to physiologic downregulation of energy-consuming processes, such as protein synthesis and fatty-acid synthesis, in order to restore ATP levels (9). This homeostatic system is conserved in evolution from yeast to humans, and LKB1 had previously been identified as a tumor-suppressor gene (10).

In multicellular organisms, the LKB1-AMPK pathway has acquired further specialized roles in regulating energy metabolism at the level of the whole organism in addition to the cellular level. The system is involved in appetite control by the central nervous system (11), and in the special case of hepatocytes, activation of the LKB1-AMPK pathway downregulates gluconeogenesis, which represents export of energy from hepatocytes to the organism in the form of glucose (12). This effect in turn reduces blood-glucose concentration, which results in a secondary decrease in insulin level. Although other mechanisms may also contribute, the inhibition of hepatic gluconeogenesis (by LKB1-AMPK activation and/or other processes; ref. 13) is now felt to be a key process underlying the utility of biguanides in therapy of type II diabetes. Precise mechanisms of action in polycystic ovary disease and antipsychotic-medication-induced obesity are unclear, but a reduction of insulin level is likely involved.

Metformin and Neoplastic Disease: Early Clues

Important early evidence suggesting that metformin is relevant to oncology came from population surveys examining associations of diabetes and diabetes treatments with cancer. Several retrospective studies showed that diabetics had an increased cancer mortality compared with non-diabetics and that diabetics on metformin had a substantially (~40%) reduced cancer burden compared with diabetics on other treatments (reviewed in refs. 14,15). Although such associations are hypothesis-generating rather than definitive, the consistency of findings among different independent studies certainly contributed to the justification for further research.

Early laboratory studies also were supportive. Experiments demonstrated that, whereas metformin activates the LKB1-AMPK pathway to inhibit gluconeogenesis in hepatocytes (12), it activates this pathway with different consequences in mouse embryo fibroblasts and human prostate, colon, and breast cancer cell lines. In these cells, consequences of LKB1-AMPK activation include inhibition of mammalian target of rapamycin (mTOR) and protein synthesis (consistent with the need to reduce energy expenditure) and thereby a reduction in proliferation (16-20). Of note, despite the early view that metformin may act as an "insulin sensitizer" in classic insulin target tissues, the afore-mentioned experimental systems revealed the opposite effect in cancer cells: Metformin inhibited insulin-stimulated mTOR activation and proliferation in an AMPK-dependent manner. The mechanisms of metformin are discussed further by Engelman and Cantley elsewhere in this issue of the journal (21).

Recent Findings Suggesting Oncologic Applications of Biguanides

Many laboratory models (for example, those reported in refs. 16,17,22-35) have been used to demonstrate the antineoplastic activity of metformin, either as a single agent or in combinations. Not all of these studies rigorously defined mechanisms, but many have provided molecular correlates of activity. There is evidence indicating that growth of untransformed epithelial cells and a subset of cancers is stimulated by insulin (reviewed in refs. 36,37) and that the systemic insulin-lowering action of metformin may inhibit proliferation of these cancers (22-24). The site of the action of metformin that leads to insulin-lowering is the liver. Therefore, the insulin-lowering effect may operate *in vivo* even in the absence of detectable metformin in neoplastic tissue or, in the context of prevention, in at-risk tissue. This action of metformin shares mechanistic features with the manner by which caloric restriction inhibits the growth of certain cancers (38,39). Anti-neoplastic action of biguanides attributable to insulin reduction is most likely to occur if the two following conditions are met: A lack of activating mutations in the signaling pathways downstream of the insulin receptor in neoplastic cells and hyperinsulinemia at baseline in the host. This mechanism may be relevant to prevention as well as treatment; indeed, the first condition is more likely to be satisfied in a prevention context.

Efforts to confirm that the separate *direct* mechanism of action of metformin involving LKB1-AMPK pathway activation within tumor cells documented *in vitro* (16-20) also operates *in vivo* have yielded a surprise. This mechanism may indeed play a role *in vivo*, but evidence also suggests that metformin can inhibit growth of LKB1-deficient tumors by a separate mechanism involving induction of cellular-energy deficiency. In the context of LKB1 deficiency, the metformin-induced decrease in mitochondrial ATP production fails to trigger a fully compensatory decrease in energy expenditure, resulting in an energy crisis and necrosis (40). If this mechanism is confirmed, it implies an attractive therapeutic index of metformin for tumors with loss of *LKB1* tumor-suppressor gene function and in Peutz Jaghers syndrome; it also suggests that agents that reduce ATP production have advantages over direct activators of AMPK (41). This implication is conceptually similar to a prior finding of enhanced activity of metformin in cancers with loss of function of p53 (25), as p53 is involved in mediating the antiproliferative, energy-conserving response to AMPK activation, and its loss accentuates metformin-induced energy stress.

The laboratory demonstration that dose and route influence benefit in the context of a lung-cancer prevention model (reported in this issue of the journal; ref. 26) reminds us that pharmacokinetic considerations need to be addressed: One cannot assume that regimens optimized for diabetes control will be optimal for possible future oncology indications. Although this study provides evidence that indirect metformin effects are mechanistically important in prevention, another encouraging recent study, which involved a colon cancer-prevention model, favored a role for direct AMPK activation (27).

Additional data of interest are provided by recent population studies. A retrospective study showed that the rate of pathologic complete response to neoadjuvant chemotherapy in diabetic breast-cancer patients was significantly higher for patients taking metformin than for those taking other antidiabetic therapies (42), suggesting a role of metformin in sensitizing cancers to chemotherapy. Support for investigations of metformin for breast-cancer prevention is provided by a provocative study that found a 56% decrease in breast cancer among diabetics receiving metformin compared with diabetics on other therapies (43). This study is not definitive, however, as it involved a small sample size and was retrospective.

Gaps in Knowledge

The collective impact of the findings summarized here is to raise enthusiasm for further investigations of biguanides for cancer prevention and treatment, not only because the data suggest activity but also because they identify important gaps in knowledge. Several key questions based on these gaps (as follows) need to be addressed before clinical studies can be optimally designed.

What are the important mechanisms of action? What is the optimum compound and dosing regimen?

The question of mechanism is not merely academic, since clarification of mechanism can facilitate clinical development. Candidate mechanisms are complex and involve both systemic (for example, insulin-lowering) and direct effects. We know with certainty from diabetes experience that conventional metformin doses are sufficient to achieve concentrations in the liver that reduce glucose production and secondarily reduce insulin levels that are elevated at baseline. One strategy to evaluate the importance of the systemic insulin-lowering actions of metformin will be to compare its efficacy with that of specific small-molecule receptor tyrosine kinase inhibitors of the insulin/insulin-like growth factor 1 (IGF-I) receptor family (36). These agents block all the relevant receptors, whereas the systemic action of metformin merely reduces insulin levels, with little effect on IGF-I or IGF-II. Therefore, any *in vivo* evidence suggesting that the activity of metformin is superior to that of specific insulin-IGF-I inhibitors would imply mechanisms of action beyond reduction of the concentration of circulating insulin. A related point is that, particularly for indications involving short-term treatment, any forthcoming evidence defining settings where the insulin-lowering actions of metformin are mechanistically important will suggest applications for insulin/IGF-I-receptor-blocking drugs, which may reduce signal transduction more effectively than would a decrease in concentration of one of the relevant ligands.

If the direct mechanism of action is important, there is a clear need for human pharmacokinetic and pharmacodynamic studies to identify the best compounds and dosing regimens for effects in extrahepatic tissues. Presence of the organic cation transporter 1 (OCT1) membrane transporter (44) may be important for cellular uptake of metformin, but less so for other biguanides, which may enter the cell in the absence of active transport. There are data implying that phenformin may be superior to metformin in activity for oncologic indications (28), and this may be due to differences in potency and pharmacokinetics. Although phenformin was withdrawn from the market because its risk of toxicity was felt to be unacceptable for long-term treatment of diabetes, it is safer than many antineoplastic agents in common use today. For cancer therapy, the toxicity of phenformin would likely be justified if it offered benefits over metformin in terms of efficacy, particularly if duration of therapy was limited, as might be the case in combinations with chemotherapy or radiotherapy. On the other hand, the risk of toxicity in the prevention setting must be minimal, and here metformin has an obvious advantage, with large data bases showing its long-term safety and tolerability; metformin is considered by many to be the safest available pharmacologic treatment for diabetes and to be safe enough for prevention indications (2,5).

Last, the possibility of novel mechanisms of action of biguanides, including immune-system modulation (45), should be considered.

Are there rational combinations to investigate?

Early results and theoretical considerations suggest that biguanides may be more effective in combinations than as single agents for cancer treatment, combined not only with chemotherapy (29) and radiotherapy (30) but also with rapalogs (rapamycin-related compounds) or with other

agents targeting energy metabolism (31). Many combinations may be effective in relatively short-term treatment durations, whereas single-agent treatment might be practical for longer durations, similar to current hormonal therapies for prostate or breast cancer. The improved rate of pathologic complete response to neoadjuvant chemotherapy in the presence of metformin (42) is evidence that at least some benefits of this biguanide do not require long-term exposure. Metformin deserves investigation for prostate-cancer therapy in the special context of preventing the hyperinsulinemia that arises as a consequence of castration (46), which may contribute to progression to castration-resistant disease (47). Combinations in the prevention setting are unexplored.

What are the relevant predictive host or tumor biomarkers?

Identification of predictive host or tumor biomarkers represents a critical issue in the contexts of both prevention and treatment, but few data are available. Possibilities include phosphatidylinositol 3-kinase (PI3K), LKB1-AMPK status in neoplastic tissue, and/or baseline fasting or postprandial serum insulin levels. Since metformin has little effect on normal insulin levels, high baseline levels might predict an increased chance of any benefit mediated by a reduction in insulin concentration. Constitutive activation of PI3K is an example of a tumor characteristic that would be expected to confer resistance to metformin's insulin-lowering effects, but not necessarily to its direct actions involving mTOR inhibition. Indeed, metformin may have some advantages over rapamycin since its inhibition of mTOR does not appear to be associated with increased AKT activation (32).

The functional status of the LKB1-AMPK pathway in neoplastic tissue may also predict activity, although it is difficult to predict the direction. Tumors or at-risk tissue where this tumor-suppressor pathway is operational may respond via the AMPK activation mechanism. On the other hand, if drug concentration is adequate, tumors or at-risk tissue with loss of function of this pathway may be sensitive to the mechanism involving metformin-induced energy stress.

Do studies of cancer incidence and mortality among diabetics in relation to metformin have implications for non-diabetics? Do they imply a chemopreventive action?

The available population studies should be interpreted with caution as they are retrospective and, even more important, are confined to diabetics, in whom risk and prognostic factors that are relatively unimportant in the general population may have dominant roles. At present, we have no clinical evidence, even retrospective, for antineoplastic activity of biguanides in non-diabetics. Phase II clinical trials that address this issue will be of key importance in guiding subsequent clinical research.

Only a few *in vivo* models have treated host metabolic status as a variable in order to study its influence on the anti-neoplastic activity of biguanides. These models (22-24) suggest that metformin is most active in the context of a high-energy diet that leads to hyperinsulinemia. If confirmed, this finding would suggest that the insulin-lowering action of metformin is important and that hyperinsulinemic subjects would be of special interest for clinical trials.

Available data suggest that metformin-associated reductions in cancer mortality are similar in magnitude to reductions in cancer incidence, raising the possibility that the main effects may be preventive, notwithstanding the experimental evidence that biguanides have antiproliferative activity for established cancers.

What potential clinical oncologic applications of biguanides deserve study?

Available data suggest that biguanides have potential applications not only for adjuvant and palliative cancer treatment, but also in cancer prevention. Apart from this broad agenda, there

is the challenge (and opportunity) related to the apparent general rather than organ-specific effects of biguanides for cancer prevention and treatment. Preclinical data justify at a minimum further research in treating or preventing colon, prostate, breast, and lung cancer. Although many research efforts are organized on a strictly organ-specific basis, biguanides are an example of where an effort to coordinate research across disease sites may accelerate progress.

For all indications, one of the key challenges in clinical-trial design is deciding who is likely enough to benefit to be included in a trial. One possibility, using the available epidemiologic clues, would be to confine early studies to diabetics. In practice, however, this would be difficult for many reasons, including the possibility that some physicians or patients would not be at equipoise for a randomization that might exclude metformin. A subgroup where randomization might be acceptable, and where biguanides might have a higher chance of efficacy than in an unselected population, would be individuals with insulin resistance and hyperinsulinemia but no formal diabetes diagnosis. Of course, it is easiest to carry out trials in unselected populations, but such studies must be large if they are to have sufficient power to detect benefits that are confined to subgroups of special interest, such as hyperinsulinemic individuals or, in an extreme example, people with Peutz Jagers syndrome.

The Way Forward

A substantial body of evidence justifies further research regarding oncologic applications of biguanides, but considerable gaps remain in the knowledge base required to optimize designs of clinical trials of these agents. We need to ascertain the best biguanide or related compound, determine the appropriate dose, and determine biologically based eligibility criteria that define subpopulations likely to benefit before conducting a costly phase-III trial involving thousands of patients and prolonged follow-up. Therefore, early steps should be to make every effort to conduct small-scale trials involving pharmacodynamic endpoints and to accelerate ongoing laboratory and epidemiologic studies. These steps may yield data that not only will contribute to the rationale for major phase-III trials but also may help optimize their design. The role of the private sector in the exploration of new indications for metformin may be problematic because the drug is off-patent, but mechanisms have been proposed to justify investment in new indications for approved agents (48). On the other hand, there is a clear incentive from an intellectual-property perspective to screen libraries of biguanides and related compounds to identify patentable molecules that may have advantages over metformin for cancer treatment.

With respect to cancer prevention, it is noteworthy that metformin may have broad activity against carcinogenesis in general, unlike intensively studied compounds that have utility only for a specific site (e.g., tamoxifen or raloxifene for the breast and finasteride or dutasteride for the prostate). If such broad-spectrum activity is confirmed, metformin could become a far more attractive chemoprevention agent than are disease-site-specific compounds. Metformin also may have benefits in the context of aging-related diseases (33) in addition to known benefits in diabetes prevention and treatment. Therefore, if the next research steps provide sufficient justification for large-scale cancerprevention trials, endpoints should include overall health outcomes. Such trials would be most justifiable, however, if future research provides criteria to identify a specific high-risk population in which metformin is expected to provide benefit.

We now have both motivating data and the tools of epidemiology, laboratory studies, and translational research to pursue specific settings where biguanides may be useful for cancer control.

Acknowledgments

Dr. Pollak holds the Alexander-Goldfarb Research Chair at McGill University and his lab is supported by Prostate Cancer Canada, the Prostate Cancer Foundation (Santa Monica), National Institutes of Health, and the Canadian Cancer Society Research Institute.

References

1. Bailey CJ, Turner RC. Metformin. *N Engl J Med* 1996;334:574–9. [PubMed: 8569826]
2. Bolen S, Feldman L, Vassy J, et al. Systematic review: comparative effectiveness and safety of oral medications for type 2 diabetes mellitus. *Ann Intern Med* 2007;147:386–99. [PubMed: 17638715]
3. Wu RR, Zhao JP, Jin H, et al. Lifestyle intervention and metformin for treatment of antipsychotic-induced weight gain: a randomized controlled trial. *JAMA* 2008;299:185–93. [PubMed: 18182600]
4. Palomba S, Falbo A, Zullo F, Orio F Jr. Evidence-based and potential benefits of metformin in the polycystic ovary syndrome: a comprehensive review. *Endocr Rev* 2009;30:1–50. [PubMed: 19056992]
5. Knowler WC, Barrett-Connor E, Fowler SE, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403. [PubMed: 11832527]
6. Thakkar N, Pirrone V, Passic S, et al. Persistent interactions between biguanide-based compound NB325 and CXCR4 result in prolonged inhibition of human immunodeficiency virus type 1 infection. *Antimicrob Agents Chemother* 2010;54:1965–72. [PubMed: 20231400]
7. Hawley SA, Ross FA, Chevtzoff C, et al. Use of cells expressing gamma subunit variants to identify diverse mechanisms of AMPK activation. *Cell Metab* 2010;11:554–65. [PubMed: 20519126]
8. El Mir MY, Nogueira V, Fontaine E, Averet N, Rigoulet M, Leverve X. Dimethylbiguanide inhibits cell respiration via an indirect effect targeted on the respiratory chain complex I. *J Biol Chem* 2000;275:223–8. [PubMed: 10617608]
9. Kahn BB, Alquier T, Carling D, Hardie DG. AMP-activated protein kinase: ancient energy gauge provides clues to modern understanding of metabolism. *Cell Metab* 2005;1:15–25. [PubMed: 16054041]
10. Shackelford DB, Shaw RJ. The LKB1-AMPK pathway: metabolism and growth control in tumour suppression. *Nat Rev Cancer* 2009;9:563–75. [PubMed: 19629071]
11. Minokoshi Y, Alquier T, Furukawa N, et al. AMP-kinase regulates food intake by responding to hormonal and nutrient signals in the hypothalamus. *Nature* 2004;428:569–74. [PubMed: 15058305]
12. Shaw RJ, Lamia KA, Vasquez D, et al. The kinase LKB1 mediates glucose homeostasis in liver and therapeutic effects of metformin. *Science* 2005;310:1642–6. [PubMed: 16308421]
13. Foretz M, Hebrard S, Leclerc J, et al. Metformin inhibits hepatic gluconeogenesis in mice independently of the LKB1/AMPK pathway via a decrease in hepatic energy state. *J Clin Invest* 2010;120:2267–70. [PubMed: 20577046]
14. Giovannucci E, Harlan DM, Archer MC, et al. Diabetes and cancer: a consensus report. *CA Cancer J Clin* 2010;60:207–21. [PubMed: 20554718]
15. Johnson JA, Pollak M. Insulin, glucose and the increased risk of cancer in patients with type 2 diabetes. *Diabetologia*. 2010 In press.
16. Dowling RJ, Zakikhani M, Fantus IG, Pollak M, Sonenberg N. Metformin inhibits mammalian target of rapamycin-dependent translation initiation in breast cancer cells. *Cancer Res* 2007;67:10804–12. [PubMed: 18006825]
17. Zakikhani M, Dowling R, Fantus IG, Sonenberg N, Pollak M. Metformin is an AMP kinase-dependent growth inhibitor for breast cancer cells. *Cancer Res* 2006;66:10269–73. [PubMed: 17062558]
18. Shaw RJ, Kosmatka M, Bardeesy N, et al. The tumor suppressor LKB1 kinase directly activates AMP-activated kinase and regulates apoptosis in response to energy stress. *Proc Natl Acad Sci U S A* 2004;101:3329–35. [PubMed: 14985505]
19. Shaw RJ, Bardeesy N, Manning BD, et al. The LKB1 tumor suppressor negatively regulates mTOR signaling. *Cancer Cell* 2004;6:91–9. [PubMed: 15261145]
20. Gwinn DM, Shackelford DB, Egan DF, et al. AMPK phosphorylation of raptor mediates a metabolic checkpoint. *Mol Cell* 2008;30:214–26. [PubMed: 18439900]

21. Engelman JA, Cantley LC. Chemoprevention meets glucose control. *Cancer Prev Res* 2010;3 XXX-- [Ed: Please complete once issue is paginated].
22. Algire C, Zakikhani M, Blouin M-J, Shuai JH, Pollak M. Metformin attenuates the stimulatory effect of a high energy diet on in vivo H59 carcinoma growth. *Endocr Relat Cancer* 2008;15:833–9. [PubMed: 18469156]
23. Algire C, Amrein L, Zakikhani M, Panasci L, Pollak M. Metformin blocks the stimulative effect of a high energy diet on colon carcinoma growth in vivo and is associated with reduced expression of fatty acid synthase. *Endocr Relat Cancer* 2010;17:351–60. [PubMed: 20228137]
24. Phoenix KN, Vumbaca F, Fox MM, Evans R, Claffey KP. Dietary energy availability affects primary and metastatic breast cancer and metformin efficacy. *Breast Cancer Res Treat.* 2009 In press.
25. Buzzai M, Jones RG, Amaravadi RK, et al. Systemic treatment with the antidiabetic drug metformin selectively impairs p53-deficient tumor cell growth. *Cancer Res* 2007;67:6745–52. [PubMed: 17638885]
26. Memmott RM, Mercado JR, Maier CR, Kawabata S, Fox SD, Dennis PA. Metformin prevents tobacco carcinogen-induced lung tumorigenesis. *Cancer Prev Res* 2010;3 XXX-- [Ed: Please complete once issue is paginated].
27. Hosono K, Endo H, Takahashi H, et al. Metformin suppresses azoxymethane-induced colorectal aberrant crypt foci by activating AMP-activated protein kinase. *Mol Carcinog* 2010;49:662–71. [PubMed: 20564343]
28. Huang X, Wullschlegel S, Shpiro N, et al. Important role of the LKB1-AMPK pathway in suppressing tumorigenesis in PTEN deficient mice. *Biochem J* 2008;412:211–21. [PubMed: 18387000]
29. Hirsch HA, Iliopoulos D, Tsiachlis PN, Struhl K. Metformin selectively targets cancer stem cells, and acts together with chemotherapy to block tumor growth and prolong remission. *Cancer Res* 2009;69:7507–11. [PubMed: 19752085]
30. Sanli T, Rashid A, Liu C, et al. Ionizing Radiation Activates AMP-Activated Kinase (AMPK): A Target for Radiosensitization of Human Cancer Cells. *Int J Radiat Oncol Biol Phys.* 2010 In press.
31. Ben SI, Laurent K, Giuliano S, et al. Targeting cancer cell metabolism: the combination of metformin and 2-deoxyglucose induces p53-dependent apoptosis in prostate cancer cells. *Cancer Res* 2010;70:2465–75. [PubMed: 20215500]
32. Zakikhani M, Blouin M-J, Piura E, Pollak M. Metformin and rapamycin have distinct effects on the AKT pathway and proliferation in breast cancer cells. *Breast Cancer Res Treat* 2010;123:271–9. [PubMed: 20135346]
33. Anisimov VN, Egormin PA, Bershtein LM, et al. Metformin decelerates aging and development of mammary tumors in HER-2/neu transgenic mice. *Bull Exp Biol Med* 2005;139:721–3. [PubMed: 16224592]
34. Zakikhani M, Dowling RJ, Sonenberg N, Pollak MN. The effects of adiponectin and metformin on prostate and colon neoplasia involve activation of AMP-activated protein kinase. *Cancer Prev Res (Phila Pa)* 2008;1:369–75.
35. Vazquez-Martin A, Oliveras-Ferraro C, Menendez JA. The antidiabetic drug metformin suppresses HER2 (erbB-2) oncoprotein overexpression via inhibition of the mTOR effector p70S6K1 in human breast carcinoma cells. *Cell Cycle* 2009;8:88–96. [PubMed: 19106626]
36. Pollak M. Insulin and insulin-like growth factor signalling in neoplasia. *Nat Rev Cancer* 2008;8:915–28. [PubMed: 19029956]
37. Law JH, Habibi G, Hu K, et al. Phosphorylated insulin-like growth factor-1/insulin receptor is present in all breast cancer subtypes and is related to poor survival. *Cancer Res* 2008;68:10238–46. [PubMed: 19074892]
38. Pollak M. Do cancer cells care if their host is hungry? *Cell Metab* 2009;9:401–3. [PubMed: 19416708]
39. Kalaany NY, Sabatini DM. Tumours with PI3K activation are resistant to dietary restriction. *Nature* 2009;458:725–31. [PubMed: 19279572]
40. Pollak, M. Proceedings of International LKB1-AMPK Congress. Marseilles: Jul. 2010 Host metabolic factors and tumor molecular pathology (including LKB1 status) interact to determine the antineoplastic action of metformin and similar compounds. www.LKB1.com

41. Cool B, Zinker B, Chiou W, et al. Identification and characterization of a small molecule AMPK activator that treats key components of type 2 diabetes and the metabolic syndrome. *Cell Metab* 2006;3:403–16. [PubMed: 16753576]
42. Jiralerspong S, Palla SL, Giordano SH, et al. Metformin and Pathologic Complete Responses to Neoadjuvant Chemotherapy in Diabetic Patients With Breast Cancer. *J Clin Oncol* 2009;27:3297–302. [PubMed: 19487376]
43. Bodmer M, Meier C, Krahenbuhl S, Jick SS, Meier CR, Meier CR. Long-term metformin use is associated with decreased risk of breast cancer. *Diabetes Care* 2010;33:1304–8. [PubMed: 20299480]
44. Shu Y, Sheardown SA, Brown C, et al. Effect of genetic variation in the organic cation transporter 1 (OCT1) on metformin action. *J Clin Invest* 2007;117:1422–31. [PubMed: 17476361]
45. Pearce EL, Walsh MC, Cejas PJ, et al. Enhancing CD8 T-cell memory by modulating fatty acid metabolism. *Nature* 2009;460:103–7. [PubMed: 19494812]
46. Smith MR, Lee H, Nathan DM. Insulin sensitivity during combined androgen blockade for prostate cancer. *J Clin Endocrinol Metab* 2006;91:1305–8. [PubMed: 16434464]
47. Cox M, Gleave M, Zakikhani M, et al. Insulin receptor expression by human prostate cancers. *The Prostate* 2009;69:33–40. [PubMed: 18785179]
48. Morgan G. More patent protection for medicines with a new purpose. *Nature* 2010;465:1005. [PubMed: 20577188]