Published in final edited form as:

Org Lett. 2010 October 1; 12(19): 4271–4273. doi:10.1021/ol101646t.

A Stereocontrolled Synthesis of ($\pm$)-Xenovenine via a Scandium (III) Catalyzed Internal Aminodiene Bicyclization Terminated by a 2-(5-Ethyl-2-thienyl)ethenyl Group

Tao Jiang and Tom Livinghouse*

Department of Chemistry and Biochemistry, Montana State University, Bozeman, MT 59717

Abstract

A highly diastereoselective binary hydroamination of a 5-amino-1,8-diene containing a 2-(5-ethyl-2-thienyl)ethenyl terminator has been utilized in an efficient synthesis of ($\pm$)-xenovenine (**1**). A pronounced rate enhancement was observed for cyclization onto the 2-(heteroaromatic)ethenyl group in comparison to a simple 1,2-disubstituted alkene.

The catalytic hydroamination of alkenes constitutes a synthetic route to amines that is endowed with exceptional atom economy. Although main-group metal complexes have been used as catalysts for alkene hydroamination,^{1a} the most general cases for the synthesis of amines and their derivatives involve transition metal catalysis using complexes of rhodium, 1b–d ruthenium, 1e–f nickel 1g–i and palladium, 1j–p gold 1q–s as well as the group 32a–n and group 42o–y metals. In contrast to catalysis by complexes of the late transition metals, hydroaminations involving complexes of the group 3 metals proceed via syn-addition of the metal-amido (M–N) bond to the C=C double bond.^{2a,b} In addition, diastereoselectivities associated with intramolecular hydroaminations of chiral substrates by group 3 metals are correlated with both the covalent radius of the metal and its ligand environment.^{2a} We have previously shown that a non-metallocene complex of Sc(III) supported by a strongly electron donating chelating diamide ligand (e.g., **3**) provides superb trans/cis selectivities in the cyclization of 2-substituted-4-pentenamines.^{2c} In this communication, we describe an efficient synthesis of ($\pm$)-xenovenine (**1**) that relies on rate enhancement attributable to an unprecedented Sc(III) catalyzed internal hydroamination involving a 2-(5-ethyl-2-thienyl)ethenyl terminator, that serves as a surrogate for a conventional alkyl group.

livinghouse@chemistry.montana.edu.

Supporting Information Available: Experimental procedures and spectroscopic data for compounds **1**, **2b**, **2c**, **4bNTs**, **4bNTs**, **5c**, **6**, **7**, **9**, and their precursors (14 pages).

The toxic pyrrolizidine alkaloid (+)-xenovenine (**1**) was isolated from the skin of the frog *Solenopsis xenoveneum* in 1980.³ From a biological standpoint it is noteworthy that this defense alkaloid is not produced by the frog but instead originates in the venom of ants that constitute, in part, the diet of the amphibian. Xenovenine (**1**) has been the subject of six total syntheses, 4a–h one of which involved a sequential bicyclization of an amino(allene)ene in the presence of a constrained Sm(III) metallocene catalyst, where initial cyclization occurred at the more reactive allene.^{4c}

We have reported that 5-amino-1,8-nonadiene (**2a**) undergoes smooth sequential bicyclization to pyrrolizidine **5a** via pyrrolidine **4a** in the presence of complex **3** (5 mol%).^{2c} The most direct and obvious route to (±)-xenovenine (**1**) should therefore entail an analogous bicyclization of aminodiene **2b**. Unfortunately, common 1,2-disubstituted alkenes are known to be very reluctant participants in internal hydroaminations catalyzed by group 3 complexes.^{2d,e} This indeed proved to be the case as **2b**, although subject to facile monocyclization to provide **4b** (t/c = 49:1, characterized as the *p*-toluenesulfonamide), underwent the required subsequent cyclization only with great lethargy and incompletely at 120 °C (Scheme 1).

We have previously demonstrated that a representative primary amine bearing a 2-(phenyl)ethenyl moiety (e.g., 1-amino-2,2-dimethyl-5-phenyl-(*E*)-4-pentene) undergoes internal hydroamination at a significantly faster rate than simple 1,2-disubstituted aminoalkenes (i.e., 1-amino-2,2-dimethyl-(*E*)-4-hexene).^{2d,h} In light of this, it seemed reasonable that terminal substitution by heteroaromatic groups might also lead to rate acceleration of internal hydroamination. To test this hypothesis in the context of the synthesis of (±)-xenovenine (**1**), the aminodiene **2c** was prepared according to the following protocol.

Sequential lithiation of 5-hexen-2-one *N,N*-dimethylhydrazone **6** (LDA, THF) followed by alkylation with 1,3-dibromo-2-propene (THF, −78 °C → 0 °C) and subsequent hydrolysis (H₃O⁺) furnished ketone **7b** as a mixture of *Z*- and *E*- isomers in 94% overall yield. Coupling of 5-ethylthiophene-2-boronic acid (**8**)⁵ with **7b** [Pd(PPh)₄ (5 mol %), Na₂CO₃, LiCl, 1,2-DME] provided ketone **9** in 66% yield as a 4:1 mixture of *Z*- and *E*-geometrical isomers, from which the pure (*Z*)- isomer could be obtained in 48% isolated yield by chromatography on silica gel. Reductive amination of (*Z*)-**9** (NaBH₃CN, NH₄OAc, MeOH) then delivered the requisite aminodiene **2c** in 85% isolated yield. Exposure of **2c** to complex **3**^{2c} (10 mol%, toluene-D₈, 10 °C) resulted in an efficient and highly diastereoselective monocyclization to generate 2,5-disubstituted pyrrolidine **4c** (t/c = 49:1, characterized as the *p*-toluenesulfonamide) with >95% conversion. It is significant that simply increasing the reaction temperature to 60 °C led to comparatively rapid (18h) and stereospecific bicyclization to afford pyrrolizidine **5c** in 90% yield [(NMR), 86% after chromatography]. Reductive desulfurization of **5c** (Raney-Ni, EtOH, 23 °C) ultimately secured (±)-xenovenine (**1**)⁶ in 98% yield (44% overall in five steps from hydrazone **6**) after purification by bulb-to-bulb distillation (Scheme 2).

The foregoing application vividly illustrates a pronounced cyclization rate enhancement involving a 2-(heteroaromatic)ethenyl substituent as compared to the corresponding saturated 2-(alkyl)alkene in a Sc(III) catalyzed hydroamination. As the latter have proven substantial obstacles to the general utilization of internal hydroaminations catalyzed by early metal complexes in synthesis, the novel strategy described here is expected to enable new pathways to target molecules of interest.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Generous financial support for this research was provided by the National Institute of General Medical Science. Support of the MSU mass spectrometry facility has been provided in part by: The Murdoch Charitable Trust, INBRE P20 RR-16455-08, P20 RR-020185 and 1P20RR024237 from the NIH COBRE Program of the National Center For Research Resources.

References and Notes

- (a) Crimmin MR, Arrowsmith M, Barrett AGM, Casely IJ, Hill MS, Procopiou PA. *J Am Chem Soc* 2009;131:9670–9685. and references cited therein. [PubMed: 19552442] (b) Matsunaga S, Yuki G. *Kagaku Kyokaishi* 2006;64:778–779. (c) Takemiya A, Hartwig JF. *J Am Chem Soc* 2006;128:6042–6043. [PubMed: 16669666] (d) Bauer EB, Andavan GTS, Hollis TK, Rubio RJ, Cho J, Kuchenbeiser GR, Helgert TR, Letko CS, Tham FS. *Org Lett* 2008;10:1175–1178. [PubMed: 18293992] (e) Takaya J, Hartwig JF. *J Am Chem Soc* 2005;127:5756–5757. [PubMed: 15839651] (f) Knodo T, Okada T, Suzuki T, Mitsudo T. *J Organomet Chem* 2001;622:149–154. (g) Fadini L, Togni A. *Tetrahedron: Asymmetry* 2008;19:2555–2562. (h) Fadani L, Togni A. *Chem Commun* 2003;1:30–31. (i) Pawlas J, Nakao Y, Kawatsura M, Hartwig JF. *J Am Chem Soc* 2002;124:3669–3679. [PubMed: 11929257] (j) Cochran BM, Michael FE. *J Am Chem Soc* 2008;130:2786–2792. [PubMed: 18254623] (k) Michael FE, Cochran BM. *J Am Chem Soc* 2006;128:4246–4247. [PubMed: 16568997] (l) Nettekoven U, Hartwig JF. *J Am Chem Soc* 2002;124:1166–1167. [PubMed: 11841274] (m) Narsireddy M, Yamamoto Y. *J Org Chem* 2008;73:9698–9709. [PubMed: 18855453] (n) Shaffer AR, Schmidt JAR. *Organometallics* 2008;27:1259–1266. (o) Siriwardana AI, Kamada M, Nakamura I, Yamamoto Y. *J Org Chem* 2005;70:5932–5937. [PubMed: 16018688] (p) Shimada T, Yamamoto Y. *J Am Chem Soc* 2002;124:12670–12671. [PubMed: 12392408] (q) LaLond RL, Wang ZJ, Mba M, Lackner AD, Toste FD. *Angew Chem Int Ed Engl* 2010;49:598–601. [PubMed: 20014376] (r) Widenhofer RA, Han X. *Eur J Org Chem* 2006;4555–4563. and references cited therein. (s) Leseurre L, Toullec PV, Genet J, Michelet V. *Org Lett* 2007;9:4049–4052. [PubMed: 17824708]
- (a) Hong S, Marks TJ. *Acc Chem Res* 2004;37:673–686. and references cited therein. [PubMed: 15379583] (b) Jiang T, Livinghouse T. *Organometallics*. manuscript in preparation. (c) Kim JY, Livinghouse T. *Org Lett* 2005;7:4391–4393. [PubMed: 16178541] (d) Kim YK, Livinghouse T. *Angew Chem Int Ed Engl* 2002;41:3645–3647. [PubMed: 12370917] (e) Tian S, Arredondo VM, Stern CL, Marks TJ. *Organometallics* 1999;18:2568–2570. (f) Gribkov DV, Hultzsck KC, Hampel F. *J Am Chem Soc* 2006;128:3748–3759. and references cited therein. [PubMed: 16536549] (g) Kim YK, Livinghouse T, Horino Y. *J Am Chem Soc* 2003;125:9560–9561. [PubMed: 12904001] (h) Kim JY, Livinghouse T. *Org Lett* 2005;7:1737–1739. [PubMed: 15844894] (i) Kim YK, Livinghouse T, Bercaw JE. *Tetrahedron Lett* 2001;42:2933–2935. (j) Lauterwasser F, Hayes PG, Brase S, Piers WE, Schafer LL. *Organometallics* 2004;23:2234–2237. (k) Hannedouche J, Aillaud I, Collin J, Schulz E, Trifonov A. *Chem Commun* 2008;3552–3554. (l) Aillaud I, Collin J, Duhayon C, Guillot R, Lyubov D, Schulz E, Trifonov A. *Chem Eur J* 2008;14:2189–2200. (m) Yu X, Marks T. *Organometallics* 2007;26:365–376. (n) Marks TJ, Hong S. *J Am Chem Soc* 2002;124:7886–7887. [PubMed: 12095317] (o) Leitch DC, Payne PR, Dunbar CR, Schafer LL. *J Am Chem Soc* 2009;131:18246–18247. [PubMed: 19994887] (p) Reznichenko AL, Hultzsck KC. *Organometallics* 2010;29:24–27. (q) Wood MC, Leitch DC, Yeung CS, Kozak JA, Schafer LL. *Angew Chem Int Ed* 2007;46:354–358. (r) Kim H, Lee PH, Livinghouse T. *Chem Commun* 2005;41:5205–5207. (s) Kim H, Kim YK, Shim JH, Kim M, Han M, Livinghouse T, Lee PH. *Adv Synth Catal* 2006;348:2609–2618. (t) Thomson RK, Bexrud JA, Schafer LL. *Organometallics* 2006;25:4069–4071. (u) Kubiak R, Prochnow I, Doye S. *Angew Chem Int Ed* 2009;48:1153–1156. (v) Watson DA, Chiu M, Bergman RG. *Organometallics* 2006;25:4731–4733. [PubMed: 19079735] (w) Mueller C, Loos C, Schulenberg N, Doye S. *Eur J Org Chem* 2006;11:2499–2503. (x) Bexrud JA, Beard JD, Leitch DC, Schafer LL. *Org Lett* 2005;7:1959–1962. [PubMed: 15876029] (y) Mueller C, Koch R, Doye S. *Chem Eur J* 2008;14:10430–10436.
- Jones TH, Blum MS, Fales HM, Thompson CR. *J Org Chem* 1980;45:4778–4780.
- (a) Xiang Y, Wang X, Zheng X, Ruan Y, Huang P. *Chem Commun* 2009;45:7045–7047. (b) Uchiyama K, Hayashi Y, Narasaka K. *Tetrahedron* 1999;55:8915–8930. (c) Arredondo VM, Tain S, McDonald FE, Marks TJ. *J Am Chem Soc* 1999;121:3633–3639. (d) Dhiman H, Vanucci-Bacque C, Hamon L, Lhommet G. *Eur J Org Chem* 1998;9:1955–1963. (e) Oppolzer W. *Gazz Chim Ital* 1995;125:207–213. (f) Oppolzer W, Bochet CG, Merifield E. *Tetrahedron Lett* 1994;35:7015–7018. (g) Oppolzer W.

Pure Appl Chem 1994;66:2127–2130. (h) Takahata H, Bandoh H, Momose T. J Org Chem 1992;57:4401–4404.

5. Conveniently prepared in 72% overall yield by the sequential lithiation of 2-ethylthiophene (*n*-BuLi) followed by the addition of the resulting organolithium to B(OMe)₃ and hydrolysis.
6. (±)-Xenovenine prepared in this manner was spectroscopically identical to the natural and previously synthesized products.

Scheme 1.

Binary Hydroamination of Non-conjugated Aminodienes

Scheme 2.
Internal Hydroamination of a 2-(Heteroaryl)ethenyl Substituent