Paramyxovirus Disruption of Interferon Signal Transduction: STATus Report

Aparna Ramachandran and Curt M. Horvath

RNA viruses in the paramyxovirus family have evolved a number of strategies to escape host cell surveillance and antiviral responses. One mechanism exploited by a number of viruses in this family is direct targeting of cytokine-inducible transcription regulators in the STAT family. Diverse members of this large virus family effectively suppress STAT signaling by the actions of their V proteins, or the related proteins derived from alternate viral mRNAs. These viral proteins have distinct means of targeting STATs, resulting in a variety of negative effects on STATs and their signal transduction. Recent developments in understanding STAT targeting will be reviewed.

Interferon Signal Transduction and Antiviral Innate Immune Responses

IMMUNE DEFENSE AGAINST VIRUSES is initiated by the recognition of specific pathogen-associated molecular patterns (PAMPs) such as viral proteins and nucleic acids by cellular pathogen recognition receptor proteins (PRRs). The PRRs include a subset of Toll-like receptors (TLR3, 7 and 9) (Alexopoulou and others 2001; Diebold and others 2004; Mogensen and Paludan 2005) that are transmembrane proteins located on the cell surface of a variety of immune cells like macrophages and dendritic cells, as well as a family of DExD/H box RNA helicases (RIG-I and MDA5) (Kang and others 2002; Yoneyama and others 2004; Yoneyama and others 2005) localized in the cytoplasm. Upon recognition of the viral products produced during infection and replication, these receptors initiate an intracellular signaling cascade that leads to the phosphorylation and activation of latent transcription factors such as interferon regulatory factor 3 (IRF3) and nuclear factor- κ B (NF- κ B). The activation of these transcription factors is followed by their nuclear translocation and binding to the promoter region of the IFN- β gene, resulting in the production of type I IFN (including IFN- α and IFN- β). Once produced and secreted from the cell, the type I IFNs bind to the IFN- α/β receptor on the same cell (autocrine signaling) as well as adjacent cells (paracrine signaling) leading to downstream signaling to create a broadly effective antiviral state. The IFN receptor-associated Janus family tyrosine kinases, JAK1 and TYK2, phosphorylate the receptor on specific tyrosine residues, creating a docking

site for the latent STAT proteins that bind to the receptor via their Src homology 2 (SH2) domain (Stahl and others 1995). In turn, the receptor-associated STAT proteins get phosphorylated at their activating tyrosine residue (Y690 for STAT2 and Y701 for STAT1) and STAT-STAT interaction occurs immediately through reciprocal phosphotyrosine SH2 interaction (Improta and others 1994). The heterodimerized STAT1 and STAT2 join together with a third protein, the DNA-binding subunit IRF9, and form a heterotrimeric complex known as the interferon-stimulated gene factor 3 (ISGF3) (Fu and others 1990; Kessler and others 1990). ISGF3 rapidly translocates to the nucleus and binds to the conserved IFN-stimulated response element (ISRE) sequences on IFN- α/β -stimulated gene (ISG) promoters inducing their transcription (Reich and others 1987; Levy and others 1988). The accumulated effects of the induced interferon-stimulated gene products lead to an antiviral state in the IFN-stimulated cell that provides a broadly effective barrier protecting the cell against virus infections. The type II IFN responses (ie, IFN- γ) are induced analogously. Upon binding of IFN- γ to its receptor, STAT1 is phosphorylated and homodimerizes (Shuai and others 1994) to form the active IFN- γ -activated factor, GAF (Decker and others 1991) that accumulates in the nucleus and binds to the promoters of target genes bearing the IFN- γ -activated sequence, GAS.

The importance of the STAT-mediated IFN-signaling systems in mediating antiviral defense is highlighted by the fact that many viruses have evolved mechanisms to evade activation of this innate immune response. For some

viruses, including the paramyxoviruses, the early steps of virus detection and IFN induction are targeted by inhibitory mechanisms including dsRNA sequestration or signaling interference to antagonize IRF3 and NF- κ B pathways. In other cases, virus-encoded soluble IFN receptors or receptor antagonists block cytokine signaling. Individual viruses can also block specific antiviral effectors to preserve key cellular machinery needed for their replication. One host evasion mechanism common to many paramyxoviruses is evasion of IFN signaling by direct interference with the IFN-inducible STAT proteins. The molecular mechanisms of STAT-directed IFN evasion are known to be very diverse amongst individual viruses in this family, and most have been reviewed in detail elsewhere (Gotoh and others 2002; Horvath 2004a, 2004b). In this article, more recent developments in understanding STAT targeting by different paramyxoviruses are highlighted.

Paramyxoviruses

The *paramyxoviruses* are enveloped negative strand RNA viruses that are subdivided into several genera including *Respirovirus* (eg, Sendai virus), *Rubulavirus* (eg, PIV5, HPIV2, mumps), *Avulavirus* (eg, Newcastle disease virus), *Morbillivirus* (eg, measles virus), and *henipaviruses* (eg, Hendra and Nipah virus). All of these viruses share common structural, biochemical, and genetic elements including the single-stranded RNA genome that encodes a small number of

proteins, including surface glycoproteins and several subunits of an RNA-dependent RNA polymerase (reviewed in Lamb (2001)). A noteworthy property found in all the paramyxoviruses is the ability of its P gene to code for more than 1 protein (Thomas and others 1988). Sendai virus P gene, for example, directs the expression of at least 7 proteins, including P, V, W, C', C, Y1, and Y2. Other paramyxoviruses express fewer but more than 1 protein from their P/V/C gene, either from overlapping open-reading frames accessed by alternate translation initiation or by a unique mechanism referred to as RNA editing that produces alternate mRNAs by site-specific cotranscriptional insertion of non-templated nucleotides (Thomas and others 1988). As a result of the RNA editing phenomenon, the P locus produces proteins that share a common N terminus but differ in their C-terminal domain (CTD). Most commonly, this process produces 2 proteins called P and V, but in some viruses the third reading frame can produce a protein referred to as W (Harcourt and others 2000).

The V protein CTD is highly conserved among the paramyxoviruses (Fig. 1A). The CTD of the V protein is cysteine-rich and binds 2 zinc (Zn) molecules per V protein (Liston and Briedis 1994; Paterson and others 1995). A diverse range of host evasion activities, including IFN-signaling inhibition (Didcock and others 1999), prevention of apoptosis (He and others 2002; Wansley and Parks 2002), cell cycle alterations (Lin and Lamb 2000), inhibition of double-stranded RNA signaling (He and others 2002; Poole and others 2002), and prevention of IFN biosynthesis (He and others

FIG. 1. Distinct mechanism of STAT inhibition by different paramyxoviruses. **(A)** Sequence alignment of the highly conserved C-terminal domain (CTD) from various paramyxovirus V proteins mentioned in text. Histidine and 7 cysteine residues and zinc fingers 1 and 2 are indicated. **(B)** Schematic diagram of PIV5, measles, and Nipah V proteins drawn to scale showing the key features, including the CTD (pink box) and important regions and residues for STAT interaction. *(Top)* PIV5 V protein. A single region for STAT interaction has not been described, but residue N100 in the N-terminal domain shared between P and V has been shown to provide species-specific interaction with human STAT2. *(Middle)* Measles V protein. The measles V CTD has a unique extension (green) of unknown significance. The CTD alone is necessary and sufficient for STAT2 interaction. Within the CTD D248 has been found to be important in measles V protein STAT2 interaction. The STAT1 interaction in measles V protein is mediated by residues 110–130 (red box) that includes tyrosine 110 (Y110). *(Bottom)* Nipah V protein. The Henipavirus CTD is dispensable for STAT interaction. Residues 100–160 (red box) are necessary and sufficient for STAT1 binding, whereas STAT2 association requires STAT1 and a larger region involving residues 100–300. The STAT1-binding domain is also an interaction site for Polo-like kinase 1 (PLK1).

2002; Poole and others 2002; Wansley and Parks 2002), have been specifically ascribed to the paramyxovirus V proteins. The other proteins derived from this locus, including C and W, have also been implicated in evading IFN responses and antiviral signaling. In most cases, a fundamental activity associated with the host evasion proteins is in direct interference with STAT protein function by mechanisms that rely on protein interactions, but it is recognized that individual genera within the family exhibit remarkably distinct mechanisms of STAT inhibition (Fig. 1B).

Rubulaviruses

The *Rubulavirus* genus includes parainfluenza virus 5 (PIV5, formerly known as SV5), mumps virus, and the type 2 human parainfluenza virus (HPIV2). In this genus the V protein is encoded by the collinear transcript; it is the product of the unedited mRNA. Insertion of 2 non-templated guanine nucleotides produces an mRNA encoding the P protein (Thomas and others 1988). The PIV5 V protein was the first to be recognized as an IFN-signaling inhibitor, and sole expression of this protein in human cells results in rapid and specific loss of STAT1 by a proteasome-mediated degradation (Didcock and others 1999). In fact, all of the *Rubulavirus* V proteins have been shown to efficiently target STAT proteins for proteasome-mediated destruction with remarkable targeting specificity. The PIV5 V protein targets STAT1, the HPIV2 V protein targets STAT2 (although in some cases can also target STAT1), and the mumps virus V protein targets both STAT1 and STAT3 (Didcock and others 1999; Goodbourn and others 2000; Young and others 2000; Kubota and others 2001; Parisien and others 2001; Parisien and others 2002a, 2002b; Yokosawa and others 2002; Ulane and others 2003). While IFN signal transduction is not required for STAT targeting, somatic cell genetic analysis has demonstrated that both PIV5 V and mumps V protein require the cellular expression of STAT2 in order to target STAT1, while HPIV2 requires STAT1 to target STAT2. This requirement restricts the host range of PIV5, as the divergent murine STAT2 fails to associate with PIV5 V protein, and cannot support targeting of STAT1 (Parisien and others 2002a). A single amino acid N100 in the PIV5 V protein has been shown to provide species-specific interaction with human STAT2 (Young and others 2001). The requirement for human STAT2 in STAT1 destruction provides the first example of an intracellular signaling and transcription factor that functions as a host range determinant for any virus. This observation was further validated by the creation of a human STAT2 transgenic mouse model system in which human STAT2 was ubiquitously expressed in an immune-competent intact mouse (Kraus and others 2008). The presence of human STAT2 in this transgenic mouse enabled targeting of the murine STAT1 protein, and resulted in specific interference with IFN antiviral responses only in infected cells. This resulted in increased virus replication in the lungs of the humanized mouse, and created an enhanced cytokine and ISG induction profile in the infected animals.

Biochemical studies of STAT targeting by *Rubulavirus* V proteins indicates that they achieve STAT degradation by coordinating the assembly of a multicomponent E3 ubiquitin ligase complex that includes a number of cellular proteins including DDB1, the cullin family member Cul4A, and both STAT1 and STAT2 (Lin and others 1998; Andrejeva and others 2002; Ulane and Horvath 2002). This V-mediated targeting

complex is referred to as VDC, an acronym for V-dependent degradation complex, which fortuitously also denotes its central components, V, DDB1, and Cul4A (Fig. 2A). DDB1 is an essential component of the VDC, required for targeting STATs by all the *Rubulavirus* V proteins (Precious and others 2005; Ulane and others 2005). Data indicate that V proteins form separate interactions with their STAT substrates and the cellular ubiquitin ligase components. Furthermore, it has been observed that the V proteins can homo- or hetero-oligomerize through their highly conserved C-terminal domain (CTD) and form macromolecular spherical particles visible in the electron microscope. Based on these facts, it has been suggested that the V protein particle acts as an enzymatic scaffold for combining the hijacked E3 ubiquitin ligase activities with the V-associated STAT substrate, resulting in highly efficient targeting (Ulane and others 2005). The obligatory role of STAT2 is to act as a substrate adaptor, bridging STAT1 and the DDB1 complex (Precious and others 2005; Ulane and others 2005). Irrespective of the structural considerations, the *Rubulavirus* V protein acts as an efficient catalyst for the degradation of STAT1 to prevent the establishment of the IFN-induced antiviral state in the cells (Carlos and others 2005; Precious and others 2007).

Structural studies strongly support the molecular and biochemical conclusions regarding the Ub ligase complex assembled by *Rubulavirus* V proteins. Crystallographic visualization of PIV5 V protein in complex with DDB1 revealed molecular details about the V protein and its association with DDB1 (Li and others 2006) (Fig. 2A). This study showed that a unique zinc finger is formed that involves the first histidine of the V CTD, and 7 cysteine residues. The structure also demonstrated the importance of both N and C terminus in interacting with DDB1 confirming the previous work that STAT1 can be degraded by only full-length protein of PIV5 V (Andrejeva and others 2002) and that only V and not P protein can participate in targeting STAT1 (Didcock and others 1999). The data are also consistent with the findings that the C-terminal cysteine cluster is important for DDB1 binding (Lin and others 1998; Andrejeva and others 2002). In addition, a series of tryptophans with characteristic spacing (W-X3-W-X9-W) has also been shown to be important for STAT protein interactions with mumps and HPIV2 V proteins (Nishio and others 2002; Nishio and others 2005a).

Based on the structural and genetic features of the *rubulaviruses*, a new paramyxovirus, mapeura virus, has been recently isolated from bats. But unlike the *rubulaviruses*, this virus evades the IFN response by sequestering STAT proteins and preventing their nuclear translocation rather than degrading STATs (Hagmaier and others 2007). Another *Rubulavirus*, HPIV4, has been shown to be completely unable to evade the IFN response (Nishio and others 2005b). Sequence comparison reveals different amino acid composition in the N and C terminus of the virus V protein. However, as cell type and species-specific factors are already known to strongly influence the outcome of V-mediated STAT destruction, further work is needed to better understand the nature of these viruses, and their abilities to interface with the host IFN response and STAT proteins.

Henipaviruses

Hendra virus and Nipah virus are the 2 prototype species in the emerging *Henipavirus* genus that was recently

FIG. 2. Structural representations of paramyxovirus V proteins. **(A)** The PIV5 V-dependent degradation complex (VDC) complex (PDB accession no. 2B5L) crystal structure (Li and others 2006). The PIV5 V protein (in red) associates with the β -propeller cluster of DDB1 (in green), which is associated with Cullin 4A (in pink) and Roc 1 (in yellow). The crystal structure of PIV5 in complex with DDB1 shows that both the N-terminal domains (NTDs) and C-terminal domains (CTDs) are required to mediate interaction with the components of E3 ligase machinery. The PIV5 V protein is positioned at the substrate-binding cleft of the E3 complex. **(B)** Structural model of measles V protein CTD showing the surface residues unique to measles V protein that were demonstrated to be important for mediating interaction with STAT2. The measles virus V protein sequence was modeled on the SV5 structure (PDB accession no. 2B5L) by use of the Swiss PDB viewer. Coloration corresponds to the activity level of mutations for each position (Ramachandran and others 2008). Gray represents nonmutated residues, cyan indicates neutral substitutions, yellow represents mutations that retain 30% to 80% activity, and red represents mutations that are >70% defective (F246 and D248) (Ramachandran and others 2008).

identified and causes lethal diseases in humans. In the *henipaviruses*, the P protein is derived from the collinear transcript, and addition of non-templated guanine nucleotides produces mRNAs encoding 2 additional proteins, V and W with different carboxyl termini (Harcourt and others 2000). In addition, the C protein is produced from an alternate translational site within the P gene. Comparison of the mRNA editing frequency of the P gene of *henipaviruses* with other viruses has revealed that *henipaviruses* produce edited transcripts at twice the frequency of other viruses (Kulkarni and others 2009; Lo and others 2009); although the biological significance, if any, of this enhanced editing is not clear, it may lead to greater capacity for host immune evasion. Immunofluorescence analysis of Nipah virus-infected cells with specific antiserum has demonstrated that the P protein localizes throughout the cell, the C protein accumulates in the cytoplasm in a punctate pattern, and the V protein accumulates exclusively in cytoplasm while the W protein is predominantly in the nucleus. This steady state picture of protein accumulation does not give justice to the dynamics of nucleocytoplasmic movement, as it is established that the *Henipavirus* V protein shuttle between nucleus and cytoplasm with net nuclear export due to the presence of a potent export signal (Rodriguez and others 2004; Lo and others 2009). The C, V, and W proteins of Nipah virus inhibit virus transcription and RNA replication when expressed in a minigenome replicon assay (Sleeman and others 2008) suggesting that these proteins share similar functional properties.

Henipavirus P, V, and W proteins have been implicated in blocking the host IFN response by targeting cellular host proteins. Nipah virus P, V, and W proteins all form high-affinity

interactions with STAT1 through their common N-terminal domain (Rodriguez and others 2004; Shaw and others 2004), and this activity maps to residues 100–160, which is also present and functional in Hendra virus (Fig. 1B). Additional residues as well as the presence of STAT1 are required for these proteins to associate with STAT2 (Rodriguez and others 2004). The *Henipavirus* V proteins inhibit IFN responses by sequestering STAT1 and STAT2 in high-molecular-weight cytoplasmic complexes, thereby preventing their IFN-mediated tyrosine phosphorylation (Rodriguez and others 2002; Rodriguez and others 2003). The short CTD of the Nipah virus W protein possesses a potent nuclear localization signal (NLS) that interacts with host karyopherin alpha 3 and 4 (Shaw and others 2005), giving rise to its net nuclear accumulation. It has also been shown that W protein but not V protein can inhibit TLR3-mediated signaling due to loss of activated IRF3, but the mechanism of TLR3 signaling inhibition by W protein remains uncharacterized (Shaw and others 2005). Recently another cellular protein, Polo-like kinase 1 (PLK1), has been identified as a binding partner for *Henipavirus* P, V, and W proteins. The PLK1-binding site overlaps with the STAT1-binding site on the P, V, and W proteins (Ludlow and others 2008), as it maps between residues 100–160 of their shared N terminus (Rodriguez and others 2002). Mutagenic evaluation of PLK1 and STAT1 binding to *Henipavirus* V protein has revealed that Ser 130 and Ser 131 are both needed for binding to STAT1 (and therefore STAT2) and normal IFN-signaling interference (Ludlow and others 2008). The interaction of PLK1 with V and W are independent of the STAT interactions suggesting V:STAT and V:PLK exist as separate complexes, but the precise biological significance of PLK1 association with *Henipavirus* V proteins remains to

be determined. A therapeutically relevant outcome of this study was the demonstration that Nipah virus P proteins that no longer interact with STATs or PLK1 retain the ability to function in minigenome replication assays. As defective STAT1 interference is an obvious strategy for attenuation of *henipaviruses*, this result suggests that reverse genetics approaches will be successful in creation of vaccine strains. Indeed it was earlier found that a Nipah virus V protein with a single amino acid substitution fails to engage STAT1 (Hagmaier and others 2006).

Morbilliviruses

Measles virus is the prototype of the *Morbillivirus* genus that infects millions of people each year despite the availability of effective vaccines. For measles, the P mRNA is derived from the collinear transcript, and both V and C proteins are generated from RNA editing or translational mechanisms. The P/V/C locus of measles virus, like other paramyxoviruses, is associated with host immune evasion, and can disengage IFN signaling. The evasion activities are ascribed to the V protein, but specific cases of P- and C-mediated host evasion have been revealed (Devaux and others 2007; Fontana and others 2008), although it has been reported that V proteins from either attenuated or wild-type strains are more potent inhibitors of IFN-inducible reporter gene expression than C proteins (Fontana and others 2008). The C protein has also been implicated as a regulator of viral RNA synthesis (Nakatsu and others 2008) and shown to be important for virus spread in infected macaque lymphatic tissue (Takeuchi and others 2005). Other *morbilliviruses* such as rinderpest virus and canine distemper virus also block type I and type II IFN through their structural and non-structural proteins (Nanda and Baron 2006; von Messling and others 2006).

The mechanism of measles V protein-mediated inhibition of IFN signaling is distinct from other paramyxoviruses. The measles virus V protein does not degrade STATs, but effectively prevents IFN-induced STAT1 and STAT2 nuclear import (Palosaari and others 2003) by an unknown mechanism. Measles virus V protein can efficiently antagonize IFN- α/β signaling but the IFN- γ signaling inhibition has not always been confirmed (Palosaari and others 2003; Takeuchi and others 2003; Caignard and others 2007; Devaux and others 2007; Fontana and others 2008). Recently, distinct means for disengaging the IFN- γ and IFN- α/β signal transduction machinery through separate associations with STAT1 and STAT2 have been revealed (Fig. 1B). Results indicate that STAT2 is the primary target for measles V protein-mediated IFN- α/β -signaling evasion. Measles virus V protein engages STAT2 in the absence of STAT1 and in the absence of the V protein's STAT1-binding site (Ramachandran and others 2008). Unlike other paramyxovirus V proteins, measles V engages STAT2 specifically through the highly conserved CTD, and this domain is both necessary and sufficient for interaction with STAT2 and IFN- α/β -signaling evasion. Results demonstrate that both the zinc finger structure and specific amino acids present on the fingers are required for mediating interactions with STAT2 (Fig. 2B). Similar findings by other groups were also reported, not only verifying the importance of the CTD for STAT2 interaction, but also implicating conserved tryptophan residues W240 and W250 in mediating STAT2 interaction (Caignard and others 2009).

The direct and fundamental association with STAT2 clearly represents a target for design of a small molecule that could disengage the measles virus IFN-signaling evasion, perhaps creating an alternative therapy for measles outbreaks. This would be complicated however, as a distinct region of the molecule associates with STAT1. The STAT1-binding site on measles V has been mapped to residues 110–130 in the region shared between P and V proteins (Ramachandran and others 2008; Caignard and others 2009). This region is noted for high sequence conservation amongst *morbilliviruses* (Devaux and others 2007). This region originates with tyrosine 110 that has been shown to be important in the P protein-mediated block of STAT1 tyrosine phosphorylation (Devaux and others 2007). Interestingly, this residue was first implicated as an attenuating mutation in the original measles virus infectious clone, Ed-TAG, that was shown to be defective in IFN evasion (Radecke and others 1995; Combredet and others 2003; Ohno and others 2004). Subsequent reports of studies using the Edmonston ATCC strain, Moraten, Schwartz, or CAM70 strains confirmed the importance of Y110 in STAT1 interference (Caignard and others 2007; Devaux and others 2007; Fontana and others 2008; Ramachandran and others 2008). It is clear from our study and prior investigations (Takeuchi and others 2003) that measles virus V protein more effectively achieves IFN- α/β signal interference than IFN- γ interference. Both P and V have the intrinsic capacity to engage STAT1 via residues 110–130, consistent with the finding that infectious measles virus lacking V protein expression retains the ability to antagonize STAT1 signaling via the P protein in a Y110-dependent reaction (Devaux and others 2007). It is likely that the differential effects of measles virus on IFN- γ signaling may become more apparent during its normal life cycle in whole animals, and can only be revealed by the study of animal models or in the context of natural measles virus infections with wild-type strains.

Conclusions

The mechanisms of STAT-directed IFN-signaling evasion by the biomedically important paramyxovirus family has been a paradigm for the study of host innate immune evasion by viruses, and has revealed that fascinating diversity can be achieved by virus evolution in accomplishing the single task of overcoming host antiviral responses. These studies have provided further insight into the molecular underpinnings of virus replication, and provided targets for therapeutic intervention and support for novel attenuation strategies. While much has been learned in a short time, we are confident that the ongoing study of these small RNA viruses will yield more novel insights into the mechanics of innate immunity, and continue to provide amusement to the dedicated scientists who are passionately intrigued by their diversity.

Acknowledgments

The authors acknowledge that the large family of viruses reviewed here is matched by an equally large number of investigators who are committed to their study. Unfortunately limitations on space and time prevent us from mentioning them comprehensively, and we apologize for any who might feel left out of this general review. C.M.H. is particularly grateful for the long-term advice and

mentoring from Dr. Robert A. Lamb, who discovered the V proteins in his laboratory in 1988, and for the contributions of Dr. Richard Randall, who first noted the disappearance of STAT1 following infection of cells with PIV5. Work on paramyxovirus interactions with STAT signaling in the Horvath laboratory is supported by NIH grant AI050707 to C.M.H.

References

- Alexopoulou L, Holt AC, Medzhitov R, Flavell RA. 2001. Recognition of double-stranded RNA and activation of NF-kappaB by Toll-like receptor 3. *Nature* 413:732–738.
- Andrejeva J, Poole E, Young DF, Goodbourn S, Randall RE. 2002. The p127 subunit (DDB1) of the UV-DNA damage repair binding protein is essential for the targeted degradation of STAT1 by the V protein of the paramyxovirus simian virus 5. *J Virol* 76:11379–11386.
- Caignard G, Bourai M, Jacob Y, Tangy F, Vidalain PO. 2009. Inhibition of IFN-alpha/beta signaling by two discrete peptides within measles virus V protein that specifically bind STAT1 and STAT2. *Virology* 383:112–120.
- Caignard G, Guerbois M, Labernardiere JL, Jacob Y, Jones LM, Wild F, Tangy F, Vidalain PO. 2007. Measles virus V protein blocks Jak1-mediated phosphorylation of STAT1 to escape IFN-alpha/beta signaling. *Virology* 368:351–362.
- Carlos TS, Fearn R, Randall RE. 2005. Interferon-induced alterations in the pattern of parainfluenza virus 5 transcription and protein synthesis and the induction of virus inclusion bodies. *J Virol* 79:14112–14121.
- Combedret C, Labrousse V, Mollet L, Lorin C, Delebecque F, Hurtrel B, McClure H, Feinberg MB, Brahic M, Tangy F. 2003. A molecularly cloned Schwarz strain of measles virus vaccine induces strong immune responses in macaques and transgenic mice. *J Virol* 77:11546–11554.
- Decker T, Lew DJ, Mirkovitch J, Darnell JE, Jr. 1991. Cytoplasmic activation of GAF, an IFN-gamma-regulated DNA-binding factor. *EMBO J* 10:927–932.
- Devaux P, von Messling V, Songsunthong W, Springfield C, Cattaneo R. 2007. Tyrosine 110 in the measles virus phosphoprotein is required to block STAT1 phosphorylation. *Virology* 360:72–83.
- Didcock L, Young DF, Goodbourn S, Randall RE. 1999. The V protein of simian virus 5 inhibits interferon signalling by targeting STAT1 for proteasome-mediated degradation. *J Virol* 73:9928–9933.
- Diebold SS, Kaisho T, Hemmi H, Akira S, Reis e Sousa C. 2004. Innate antiviral responses by means of TLR7-mediated recognition of single-stranded RNA. *Science* 303:1529–1531.
- Fontana JM, Bankamp B, Bellini WJ, Rota PA. 2008. Regulation of interferon signaling by the C and V proteins from attenuated and wild-type strains of measles virus. *Virology* 374:71–81.
- Fu XY, Kessler DS, Veals SA, Levy DE, Darnell JE, Jr. 1990. ISGF3, the transcriptional activator induced by interferon alpha, consists of multiple interacting polypeptide chains. *Proc Natl Acad Sci USA* 87:8555–8559.
- Goodbourn S, Didcock L, Randall RE. 2000. Interferons: cell signaling, immune modulation, antiviral response and virus countermeasures. *J Gen Virol* 81:2341–2364.
- Gotoh B, Komatsu T, Takeuchi K, Yokoo J. 2002. Paramyxovirus strategies for evading the interferon response. *Rev Med Virol* 12:337–357.
- Hagmaier K, Stock N, Goodbourn S, Wang LF, Randall R. 2006. A single amino acid substitution in the V protein of Nipah virus alters its ability to block interferon signalling in cells from different species. *J Gen Virol* 87:3649–3653.
- Hagmaier K, Stock N, Precious B, Childs K, Wang LF, Goodbourn S, Randall RE. 2007. Mapuera virus, a rubulavirus that inhibits interferon signalling in a wide variety of mammalian cells without degrading STATs. *J Gen Virol* 88:956–966.
- Harcourt BH, Tamin A, Ksiazek TG, Rollin PE, Anderson LJ, Bellini WJ, Rota PA. 2000. Molecular characterization of Nipah virus, a newly emergent paramyxovirus. *Virology* 271:334–349.
- He B, Paterson RG, Stock N, Durbin JE, Durbin RK, Goodbourn S, Randall RE, Lamb RA. 2002. Recovery of paramyxovirus simian virus 5 with a V protein lacking the conserved cysteine-rich domain: the multifunctional V protein blocks both interferon-beta induction and interferon signaling. *Virology* 303:15–32.
- Horvath CM. 2004a. Silencing STATs: lessons from paramyxovirus interferon evasion. *Cytokine Growth Factor Rev* 15:117–127.
- Horvath CM. 2004b. Weapons of STAT destruction. Interferon evasion by paramyxovirus V protein. *Eur J Biochem* 271:4621–4628.
- Improta T, Schindler C, Horvath CM, Kerr IM, Stark GR, Darnell JE, Jr. 1994. Transcription factor ISGF-3 formation requires phosphorylated Stat91 protein, but Stat113 protein is phosphorylated independently of Stat91 protein. *Proc Natl Acad Sci USA* 91:4776–4780.
- Kang DC, Gopalkrishnan RV, Wu Q, Jankowsky E, Pyle AM, Fisher PB. 2002. mda-5: An interferon-inducible putative RNA helicase with double-stranded RNA-dependent ATPase activity and melanoma growth-suppressive properties. *Proc Natl Acad Sci USA* 99:637–642.
- Kessler DS, Veals SA, Fu XY, Levy DE. 1990. Interferon-alpha regulates nuclear translocation and DNA-binding affinity of ISGF3, a multimeric transcriptional activator. *Genes Dev* 4:1753–1765.
- Kraus TA, Garza L, Horvath CM. 2008. Enabled interferon signaling evasion in an immune-competent transgenic mouse model of parainfluenza virus 5 infection. *Virology* 371:196–205.
- Kubota T, Yokosawa N, Yokota S, Fujii N. 2001. C terminal CYS-RICH region of mumps virus structural V protein correlates with block of interferon alpha and gamma signal transduction pathway through decrease of STAT 1-alpha. *Biochem Biophys Res Commun* 283:255–259.
- Kulkarni S, Volchkova V, Basler CF, Palese P, Volchkov VE, Shaw ML. 2009. Nipah virus edits its P gene at high frequency to express the V and W proteins. *J Virol* 83:3982–3987.
- Lamb RA, Kolakofsky D. 2001. Paramyxoviridae: The viruses and their replication. Philadelphia: Lippincott-Raven Publishers.
- Levy DE, Kessler DS, Pine R, Reich N, Darnell JE, Jr. 1988. Interferon-induced nuclear factors that bind a shared promoter element correlate with positive and negative transcriptional control. *Genes Dev* 2:383–393.
- Li T, Chen X, Garbutt KC, Zhou P, Zheng N. 2006. Structure of DDB1 in complex with a paramyxovirus V protein: viral hijack of a propeller cluster in ubiquitin ligase. *Cell* 124:105–117.
- Lin GY, Lamb RA. 2000. The paramyxovirus simian virus 5 V protein slows progression of the cell cycle. *J Virol* 74:9152–9166.
- Lin GY, Paterson RG, Richardson CD, Lamb RA. 1998. The V protein of the paramyxovirus SV5 interacts with damage-specific DNA binding protein. *Virology* 249:189–200.
- Liston P, Briedis DJ. 1994. Measles virus V protein binds zinc. *Virology* 198:399–404.
- Lo MK, Harcourt BH, Mungall BA, Tamin A, Peeples ME, Bellini WJ, Rota PA. 2009. Determination of the henipavirus phosphoprotein gene mRNA editing frequencies and detection of the C, V and W proteins of Nipah virus in virus-infected cells. *J Gen Virol* 90:398–404.
- Ludlow LE, Lo MK, Rodriguez JJ, Rota PA, Horvath CM. 2008. Henipavirus V protein association with Polo-like kinase reveals functional overlap with STAT1 binding and interferon evasion. *J Virol* 82:6259–6271.
- Mogensen TH, Paludan SR. 2005. Reading the viral signature by Toll-like receptors and other pattern recognition receptors. *J Mol Med* 83:180–192.
- Nakatsu Y, Takeda M, Ohno S, Shirogane Y, Iwasaki M, Yanagi Y. 2008. Measles virus circumvents the host interferon response by different actions of the C and V proteins. *J Virol* 82:8296–8306.
- Nanda SK, Baron MD. 2006. Rinderpest virus blocks type I and type II interferon action: role of structural and nonstructural proteins. *J Virol* 80:7555–7568.

- Nishio M, Garcin D, Simonet V, Kolakofsky D. 2002. The carboxyl segment of the mumps virus V protein associates with Stat proteins in vitro via a tryptophan-rich motif. *Virology* 300:92–99.
- Nishio M, Tsurudome M, Ito M, Garcin D, Kolakofsky D, Ito Y. 2005a. Identification of paramyxovirus V protein residues essential for STAT protein degradation and promotion of virus replication. *J Virol* 79:8591–8601.
- Nishio M, Tsurudome M, Ito M, Ito Y. 2005b. Human parainfluenza virus type 4 is incapable of evading the interferon-induced antiviral effect. *J Virol* 79:14756–14768.
- Ohno S, Ono N, Takeda M, Takeuchi K, Yanagi Y. 2004. Dissection of measles virus V protein in relation to its ability to block alpha/beta interferon signal transduction. *J Gen Virol* 85:2991–2999.
- Palosaari H, Parisien JP, Rodriguez JJ, Ulane CM, Horvath CM. 2003. STAT protein interference and suppression of cytokine signal transduction by measles virus V protein. *J Virol* 77:7635–7644.
- Parisien JP, Lau JF, Horvath CM. 2002a. STAT2 acts as a host range determinant for species-specific paramyxovirus interferon antagonism and simian virus 5 replication. *J Virol* 76:6435–6441.
- Parisien JP, Lau JF, Rodriguez JJ, Sullivan BM, Moscona A, Parks GD, Lamb RA, Horvath CM. 2001. The V protein of human parainfluenza virus 2 antagonizes type I interferon responses by destabilizing signal transducer and activator of transcription 2. *Virology* 283:230–239.
- Parisien JP, Lau JF, Rodriguez JJ, Ulane CM, Horvath CM. 2002b. Selective STAT protein degradation induced by paramyxoviruses requires both STAT1 and STAT2 but is independent of alpha/beta interferon signal transduction. *J Virol* 76:4190–4198.
- Paterson RG, Leser GP, Shaughnessy MA, Lamb RA. 1995. The paramyxovirus SV5 V protein binds two atoms of zinc and is a structural component of virions. *Virology* 208:121–131.
- Poole E, He B, Lamb RA, Randall RE, Goodbourn S. 2002. The V proteins of simian virus 5 and other paramyxoviruses inhibit induction of interferon-beta. *Virology* 303:33–46.
- Precious B, Childs K, Fitzpatrick-Swallow V, Goodbourn S, Randall RE. 2005. Simian virus 5 V protein acts as an adaptor, linking DDB1 to STAT2, to facilitate the ubiquitination of STAT1. *J Virol* 79:13434–13441.
- Precious BL, Carlos TS, Goodbourn S, Randall RE. 2007. Catalytic turnover of STAT1 allows PIV5 to dismantle the interferon-induced anti-viral state of cells. *Virology* 368:114–121.
- Radecke F, Spielhofer P, Schneider H, Kaelin K, Huber M, Dotsch C, Christiansen G, Billeter MA. 1995. Rescue of measles viruses from cloned DNA. *EMBO J* 14:5773–5784.
- Ramachandran A, Parisien JP, Horvath CM. 2008. STAT2 is a primary target for measles virus V protein-mediated alpha/beta interferon signaling inhibition. *J Virol* 82:8330–8338.
- Reich N, Evans B, Levy D, Fahey D, Knight E, Jr, Darnell JE, Jr. 1987. Interferon-induced transcription of a gene encoding a 15-kDa protein depends on an upstream enhancer element. *Proc Natl Acad Sci USA* 84:6394–6398.
- Rodriguez JJ, Cruz CD, Horvath CM. 2004. Identification of the nuclear export signal and STAT-binding domains of the Nipah virus V protein reveals mechanisms underlying interferon evasion. *J Virol* 78:5358–5367.
- Rodriguez JJ, Parisien JP, Horvath CM. 2002. Nipah virus V protein evades alpha and gamma interferons by preventing STAT1 and STAT2 activation and nuclear accumulation. *J Virol* 76:11476–11483.
- Rodriguez JJ, Wang LF, Horvath CM. 2003. Hendra virus V protein inhibits interferon signaling by preventing STAT1 and STAT2 nuclear accumulation. *J Virol* 77:11842–11845.
- Shaw ML, Cardenas WB, Zamarin D, Palese P, Basler CF. 2005. Nuclear localization of the Nipah virus W protein allows for inhibition of both virus- and toll-like receptor 3-triggered signaling pathways. *J Virol* 79:6078–6088.
- Shaw ML, Garcia-Sastre A, Palese P, Basler CF. 2004. Nipah virus V and W proteins have a common STAT1-binding domain yet inhibit STAT1 activation from the cytoplasmic and nuclear compartments, respectively. *J Virol* 78:5633–5641.
- Shuai K, Horvath CM, Huang LH, Qureshi SA, Cowburn D, Darnell JE, Jr. 1994. Interferon activation of the transcription factor Stat91 involves dimerization through SH2-phosphotyrosyl peptide interactions. *Cell* 76:821–828.
- Sleeman K, Bankamp B, Hummel KB, Lo MK, Bellini WJ, Rota PA. 2008. The C, V and W proteins of Nipah virus inhibit minigenome replication. *J Gen Virol* 89:1300–1308.
- Stahl N, Farruggella TJ, Boulton TG, Zhong Z, Darnell JE, Jr, Yancopoulos GD. 1995. Choice of STATs and other substrates specified by modular tyrosine-based motifs in cytokine receptors. *Science* 267:1349–1353.
- Takeuchi K, Kadota SI, Takeda M, Miyajima N, Nagata K. 2003. Measles virus V protein blocks interferon (IFN)-alpha/beta but not IFN-gamma signaling by inhibiting STAT1 and STAT2 phosphorylation. *FEBS Lett* 545:177–182.
- Takeuchi K, Takeda M, Miyajima N, Ami Y, Nagata N, Suzuki Y, Shahnewaz J, Kadota S, Nagata K. 2005. Stringent requirement for the C protein of wild-type measles virus for growth both in vitro and in macaques. *J Virol* 79:7838–844.
- Thomas SM, Lamb RA, Paterson RG. 1988. Two mRNAs that differ by two nontemplated nucleotides encode the amino coterminal proteins P and V of the paramyxovirus SV5. *Cell* 54:891–902.
- Ulane CM, Horvath CM. 2002. Paramyxoviruses SV5 and HPIV2 assemble STAT protein ubiquitin ligase complexes from cellular components. *Virology* 304:160–166.
- Ulane CM, Kentsis A, Cruz CD, Parisien JP, Schneider KL, Horvath CM. 2005. Composition and assembly of STAT-targeting ubiquitin ligase complexes: paramyxovirus V protein carboxyl terminus is an oligomerization domain. *J Virol* 79:10180–10189.
- Ulane CM, Rodriguez JJ, Parisien JP, Horvath CM. 2003. STAT3 ubiquitylation and degradation by mumps virus suppress cytokine and oncogene signaling. *J Virol* 77:6385–6393.
- von Messling V, Svitek N, Cattaneo R. 2006. Receptor (SLAMF7) recognition and the V protein sustain swift lymphocyte-based invasion of mucosal tissue and lymphatic organs by a morbillivirus. *J Virol* 80:6084–6092.
- Wansley EK, Parks GD. 2002. Naturally occurring substitutions in the P/V gene convert the noncytopathic paramyxovirus simian virus 5 into a virus that induces alpha/beta interferon synthesis and cell death. *J Virol* 76:10109–10121.
- Yokosawa N, Yokota S, Kubota T, Fujii N. 2002. C-terminal region of STAT-1alpha is not necessary for its ubiquitination and degradation caused by mumps virus V protein. *J Virol* 76:12683–12690.
- Yoneyama M, Kikuchi M, Matsumoto K, Imaizumi T, Miyagishi M, Taira K, Foy E, Loo YM, Gale M, Jr, Akira S, Yonehara S, Kato A, Fujita T. 2005. Shared and unique functions of the DExD/H-box helicases RIG-I, MDA5, and LGP2 in antiviral innate immunity. *J Immunol* 175:2851–2858.
- Yoneyama M, Kikuchi M, Natsukawa T, Shinobu N, Imaizumi T, Miyagishi M, Taira K, Akira S, Fujita T. 2004. The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. *Nat Immunol* 5:730–737.
- Young DE, Chatziandreou N, He B, Goodbourn S, Lamb RA, Randall RE. 2001. Single amino acid substitution in the V protein of simian virus 5 differentiates its ability to block interferon signaling in human and murine cells. *J Virol* 75:3363–3370.
- Young DE, Didcock L, Goodbourn S, Randall RE. 2000. Paramyxoviridae use distinct virus-specific mechanisms to circumvent the interferon response. *Virology* 269:383–390.

Address correspondence to:
 Dr. Curt M. Horvath
 Pancoe-ENH Research Pavilion
 Northwestern University
 2200 Campus Drive
 Evanston, IL 60208
 E-mail: horvath@northwestern.edu

Received 29 June 2009/Accepted 29 June 2009