Published in final edited form as:

Proc Soc Exp Biol Med. 1968 May ; 128(1): 247–250.

Changes in Serum Haptoglobin and Group Specific Component after Orthotopic Liver Homotransplantation in Humans*

Nobour Kashiwagi, Carl G. Groth, and Thomas E. Starzl

Department of Surgery, University of Colorado School of Medicine; and the Denver Veterans Administration Hospital, Denver, Colorado 80220

Successful clinical liver transplantation should make it possible to more incisively explore certain aspects of hepatic function in man. In the present study this has been done by examining the haptoglobin (Hp) and group specific component (Gc) of the α_2 globulin serum fraction of 4 infant recipients of orthotopic liver homografts and their donors. The results establish that both protein moieties are manufactured exclusively by the liver. They also provide evidence that liver homografts retain the metabolic specificity of the donor after transfer to a new host.

The previous reports of Smithies on Hp (1,2) and of Hirschfeld on Gc (3,4) provided the basis for the experimental approach. They showed that 3 kinds of both Hp and Gc were identifiable in the human population, that the type present in any individual was subject to genetic control, and that the phenotypic expression could be identified with electrophoretic techniques.

Materials and Methods

The 4 liver recipients were females, aged 13–21 months at the time of transplantation. The first patient had a hepatoma; the other 3 had congenital biliary atresia. The cadaveric donors were 2 males and 2 females, aged 14–20 months. Three of the 4 children are still alive, now 180, 136, and 56 days after operation; the fourth died 61 days posttransplantation. A more general account of these cases is published elsewhere (5).

Serum samples were taken from both the donors and recipients prior to transplantation; in Case 1, the parents' sera were also analyzed in order to confirm the Hp genotype. Postoperatively, recipient samples were obtained at variable intervals. The serum Hp genotypes were determined by electrophoresis in a vertical starch gel¹ (6). The wells were charged with a solution consisting of 15 parts of undiluted serum and one part of 10% human hemoglobin solution. The initial current was 6 mA. A constant voltage of 120 was maintained for 24 hours. The gel was removed from the supporting tray and divided horizontally into slices 2–3 mm thick. The gel slice was stained in 0.1% *o*-toluidine in 1% acetic acid and a few drops of 30% H₂O₂ were added.

The determination of serum Gc genotypes was made by Hirschfeld's modification (7) of the microimmunoelectrophoresis² technique of Scheidegger (8). Commercial horse anti-Gc antiserum and rabbit anti-Gc antiserum were used³. Electrophoresis was at 220 volts for 2.5 hours with a discontinuous buffer system (pH 8.6, $\mu = 0.05$ in agar⁴ on plate and pH 8.6, $\mu =$

*Supported by United States Public Health Service Grants AM 06283, AM 06344, HE 07735, AM 07772, AI 04152, FR 00051, FR 00069, FO5 TW 1154.

Introduced by D. W. Talmage

¹From Otto Hiller, Madison, Wisconsin.

²Equipment from LKB-Produkter, Stockholm, Sweden.

³From Hoechst Pharmaceutical Co., Kansas City, Missouri.

0.1 in buffer tank). Agar plates charged with antiserum were incubated in a humid chamber at 37°C for 24 hours and stained with amido black.

Results

The Hp type of both the donors and recipients was 2-2 in Cases 1 and 4. In Case 1, the same type was present after transplantation in samples obtained on post-operative days 10, 52, and 102. Patient 4 was studied 6, 12, 40, and 58 hours posttransplantation and after 4 and 6 days. During the first 12 hours, there was a 2-1 Hp type mixed with the 2-2, presumably as the result of 12C0 ml blood transfusion during operation. The subsequent samples were 2-2 (Fig. 1.)

Patients 2 and 3 had preoperative Hp types 2-2 and 2-1 respectively, whereas their donors were 1-1 and 2-2. After transplantation, both recipients reverted to the Hp types of their donors for the period of their study. In Case 2, the sera tested on days 8, 25, 38, and 64 showed only Hp type 1-1 and no sign of the original Hp 2-2. Patient 3 was studied on days 1, 2, 3, 5, 10, 15, and 31. On all occasions, the only Hp detectable was the donor type 2-2 (Fig. 1).

Patients 3 and 4, who received homografts from donors of the same Gc type, did not have a change in group specific component. In Cases 1 and 2, however, the Gc type of the recipients converted to that of the donors. In patients 1 and 2, the alterations were complete by 8 and 10 days, respectively, when the first analyses were made. These remained unchanged for the subsequent follow-up periods of 4 and 3 months (Fig. 2).

Discussion

There have been previous descriptions of changes in these 2 identifiable proteins. In 1964, we reported that an adult recipient of a liver homograft had converted by the second postoperative day to the Hp type of the organ donor (9,10). However, liver function was poor and after the fourth postoperative day, all detectable haptoglobin had disappeared from the serum. Hirschfeld *et al.* (11) observed that a newborn infant's serum after exchange transfusion contained both his own Gc and that of the blood donor.

In the reports of the earlier liver transplantation (9,10), it was postulated that the liver was the only source of Hp, a suggestion which was: somewhat weakened by the transient nature of the observations. Studies by Prunier *et al.* (12) on various human tissue cultures supported the hypothesis that both Hp and Gc were manufactured solely by the liver. The data obtained in our most recent cases seem to establish this fact beyond doubt.

Nyman (13) has estimated the 50% turn-over rate of Hp to be 5.4 days. It is clear that this is accelerated under the special conditions of liver transplantation since all trace of the original recipient Hp was gone at least in one case within 24 hours. The Hp combines with free hemoglobin to form a complex which cannot be filtered by the kidney (14,15); it has been suggested that this reaction not only prevents excessive losses of endogenous iron, but also shields the tubules from free hemoglobin deposition (16). Thus, hemolysis from massive intraoperative blood transfusion was probably of major importance in increasing Hp utilization. An additional factor could be an increased rate of red cell destruction which is known to be caused by the presence of liver homografts (17). Trauma and steroid therapy also influence Hp metabolism (16,18). It is probable that the conversion of the Gc phenotype was also rapid. There is no information on the biologic function of Gc protein.

⁴Special agar-Noble from Difco Laboratories, Detroit, Michigan.

A practical implication of the demonstration that the liver retains its metabolic specificity after transplantation is that certain liver-based metabolic diseases might be treatable by such a procedure. This concept has been conclusively tested by Kuster *et al.* (19). Using mongrel canine donors, they were able to cure the gout present in Dalmatian recipients. Conversely, the transplantation of Dalmatians' livers conferred the defect in uric acid metabolism upon mongrel recipients.

Summary

In human recipients of orthotopic liver homografts with different haptoglobin (Hp) and group specific component (Gc) types than their respective donors, the donor phenotype permanently replaced that previously present. The findings prove that the liver is the sole source of Hp and Gc, and that it retains its metabolic specificity after transplantation to a new host.

Acknowledgments

We thank Mrs. J. Guptill, M. T., for excellent technical assistance.

References

1. Smithies O. Biochem J 1955;61:629. [PubMed: 13276348]
2. Smithies O, Walker NF. Nature 1956;178:694. [PubMed: 13369501]
3. Hirschfeld J. Acta Pathol Microbiol Scand 1959;47:160. [PubMed: 14402000]
4. Hirschfeld J, Jonsson B, Rasmuson M. Nature 1960;185:931.
5. Starzl TE, Groth CG, Brettschneider L, Moon JB, Fulginiti VA, Cotton EK, Porter KA. Surgery 1968;63:549. [PubMed: 4171413]
6. Smithies O. Biochem J 1959;71:585. [PubMed: 13638269]
7. Hirschfeld J. Acta Pathol Microbiol Scand 1959;46:224.
8. Scheidegger JJ. Intern Arch Allergy Appl Immunol 1955;7:103.
9. Starzl TE, Marchioro TL, Rowlands DT Jr, Kirkpatrick CH, Wilson WEC, Rifkind LO, Waddell WR. Ann Surg 1964;160:411. [PubMed: 14206848]
10. Merrill DA, Kirkpatrick CH, Wilson WEC, Riley CM. Proc Soc Exptl Biol Med 1964;116:748. [PubMed: 14194652]
11. Hirschfeld J, Nilsson BA. Acta Pathol Microbiol Scand 1962;56:471. [PubMed: 13961208]
12. Prunier JH, Bearn AG, Cleve H. Proc Soc Exptl Biol Med 1964;115:1005. [PubMed: 14166544]
13. Nyman M. Scand J Clin Lab Invest Suppl 1959;39.
14. Laurell CB, Nyman M. Blood 1957;12:493. [PubMed: 13436506]
15. Allison AC, Rees W. Brit Med J 1957;2:1137. [PubMed: 13472068]
16. Laurell CB, Gronvall C. Advan Clin Chem 1962;5:135.
17. Starzl TE, Marchioro TL, Porter KA, Taylor PD, Faris TD, Herrmann TJ, Hlad CJ, Waddell WR. Surgery 1965;58:131. [PubMed: 14305148]
18. Jayle MF, Badin J. Presse Med 1953;61:343. [PubMed: 13073976]
19. Kuster G, Shorter RG, Dawson B, Hallenbeck GA. Surg Forum 1967;18:360.

Fig. 1.

Effect of liver transplantation upon serum haptoglobin (Hp). Patients 2 (top) and 3 (center) had preoperative Hp types 2-2, and 2-1, whereas their donors had 1-1 and 2-2, respectively. After transplantation, only the donor type was detectable in the recipient sera. In Case 4 (bottom), both the donor and recipient were 2-2. For the first 12 hours after transplantation, the recipient serum, was found to contain Hp 2-1 mixed with 2-2, presumably a consequence of massive blood transfusion during the operation. All subsequent samples were 2-2.

Fig. 2.
In Cases 1 (top) and 2 (bottom), the group specific component (Gc) types of the recipients which were 2-1 and 1-1, were converted after transplantation to the 1-1 and 2-1 types of their donors.