Published in final edited form as:

Org Lett. 2010 November 5; 12(21): 4936–4939. doi:10.1021/ol1022036.

One-Pot Three-Step Synthesis of 1,2,3-Triazoles by Copper-Catalyzed Cycloaddition of Azides with Alkynes formed by a Sonogashira Cross-Coupling and Desilylation

Frédéric Friscourt^a and Geert-Jan Boons^{*,a}

^aComplex Carbohydrate Research Center, University of Georgia, 315 Riverbend Road, Athens, GA 30602 USA

Abstract

A microwave-assisted one-pot, three-step Sonogashira cross coupling-desilylation-cycloaddition sequence was developed for the convenient preparation of 1,4-disubstituted 1,2,3-triazoles starting from a range of halides, acyl chlorides, ethynyltrimethylsilane and azides.

The exquisite regioselectivity, high tolerance to the presence of functional groups and exceptional high yields of Cu(I)-catalyzed 1,3-dipolar cycloadditions of azides with terminal alkynes (CuAAC) to give 1,4-disubstituted triazoles¹ have made it a powerful approach for bioconjugation,² construction and modification of materials³ and the parallel combinatorial synthesis of libraries of compounds for drug discovery.⁴ Recent efforts to further develop CuAAC have focused on one-pot multi-component reactions in which azides are prepared *in-situ* prior to the cycloaddition. The attraction of such an approach is that it minimizes time-consuming work-up and purification protocols and avoids handling of potentially explosive azides. Most methods for *in-situ* preparation of azides take advantage of the nucleophilicity of sodium azide towards electrophiles such as aromatic fluorides, iodides, benzylic bromides as well as epoxides and boronic acids.⁵ Organic azides have also been generated *in-situ* by Cu(II)-catalyzed diazo-transfer of triflyl azide or azidotrimethylsilane to amines.⁶ Surprisingly, one-pot multi-step reactions involving *in-situ* generation of terminal alkynes for click chemistry have not been reported. The attraction of such an approach is that a wide variety of alkynes can be generated from simple and readily available starting materials, such as aromatic halides.

Herein, we describe a microwave-assisted one-pot three-step procedure entailing Sonogashira cross-coupling of aromatic halides with ethynyltrimethylsilane (TMS-acetylene) to give TMS-protected alkynes, which were desilylated and then employed in

* gjboons@ccrc.uga.edu.

Supporting Information Available: Experimental procedures, characterization data, and copies of ¹H NMR and ¹³C NMR spectra of all products are available. This material is available free of charge via the Internet at <http://pubs.acs.org>.

1,3-dipolar cycloadditions with various azides to give a range of 1,2,3-triazoles (Scheme 1). In addition, it was found that the use of aromatic acyl chlorides in the Sonogashira cross-coupling gave TMS-protected ynones, which after desilylation and Cu(I)-catalyzed cycloadditions with azides led to the regioselective formation of 1-substituted 4-phenylacyl-1*H*-1,2,3-triazoles.

First, optimal reaction conditions were established for a fast and high yielding one-pot desilylation-cycloaddition reaction sequence (Table 1). It has been reported that TMS-modified acetylenes can readily be deprotected using reagents such as tetrabutylammonium fluoride (TBAF)⁷ or silver salts⁸ to give terminal alkynes, which can then undergo cycloadditions with azides. It was expected that these two steps can be performed as a one-pot process and accelerated by microwave irradiation.⁹ A temperature survey revealed that reaction of trimethyl(phenylethynyl)silane (**1a**) with benzyl azide (**2a**) in the presence of TBAF (1 equiv), CuI (10 mol %) and *N,N*-diisopropylethylamine (20 mol %) in methanol under microwave irradiation at 120 °C was complete within 20 min to afford the 1,4-disubstituted triazole **3a** in an almost quantitative yield as a single regioisomer (Table 1, entry 1, method A). A similar reaction without TBAF resulted in no triazole formation, clearly establishing that the TMS-alkyne must be deprotected to undergo a 1,3 dipolar-cycloaddition. Furthermore, when the transformation was performed in the absence of CuI, only a trace amount of triazole was isolated as a mixture of regioisomers.¹⁰ The desilylation-cycloaddition reaction could also be performed at ambient temperature, however, in this case a prolonged reaction time of 12 h was required to achieve complete conversion of the starting materials.

We have found that CuF₂ is an efficient reagent for deprotecting TMS-modified alkynes and promoting 1,3-dipolar cycloadditions with azides to give 1,4-disubstituted triazoles.¹¹ As expected, the use of CuF₂ (2 equiv) also gave the 1,2,3-triazole **3a** in an excellent yield (Table 1, entry 1, method B). It was found that the scope of the one-pot process is excellent and 1,4-disubstituted triazoles containing electron donating (entry 2), withdrawing (entry 3) and bulky functionalities (entry 6) could be obtained in excellent yield. Furthermore, the use of CuF₂ was found to be most convenient and gave in general slightly higher yields of product compared to the use of TBAF/CuI. It is also possible to employ a catalytic amount of CuF₂ (10 mol %) however in this case the addition of TBAF (1 equiv) is required to facilitate desilylation.

TMS-substituted alkynes are readily accessible by Sonogashira cross-coupling between aromatic halides and ethynyltrimethylsilane.¹² Furthermore, it has been found that microwave irradiation can significantly reduce the reaction time of metal-catalyzed cross-coupling reactions.¹³ Thus, we explored whether Sonogashira cross-coupling, desilylation and cycloaddition can be performed under microwave conditions as a one-pot procedure. Various aromatic iodides (Table 2, entries 1-6 and 10-11) were reacted with ethynyltrimethylsilane using a catalytic amount of Pd(PPh₃)₄ (5 mol %) and CuI (10 mol %) in the presence of *N,N*-diisopropylethylamine (2 equiv) in methanol under microwave irradiation at 120 °C for 20 min, followed by the introduction of an azide and TBAF/CuI or CuF₂ and further microwave irradiation at 120 °C for 20 min. Gratifyingly, the one-pot three-step process produced the expected 1,4-disubstituted triazoles (**3**) in most cases in high yield. As expected, aromatic bromides (entries 7-9), which are known to couple poorly under classical Sonogashira cross-coupling conditions, required optimization of the palladium source, choice of amine and reaction time. It was found that triazoles were obtained in acceptable yields when a mixture of an aromatic bromide, ethynyltrimethylsilane, Pd(PPh₃)₂Cl₂ (5 mol %) and CuI (10 mol %) in Et₃N was heated under reflux prior to the desilylation-cycloaddition protocol (entries 8 and 9).

The scope of the one-pot three-step protocol was further extended to the synthesis of 1-substituted 4-acyl-1*H*-1,2,3-triazoles by using acyl chlorides as the starting material in the Sonogashira cross-coupling reaction (7, Table 3).^{12b,14} Acyl chlorides did not tolerate the use of microwave irradiation,¹⁵ however, the desired TMS-protected ynones could be obtained by simply reacting an acyl chloride (6) with ethynyltrimethylsilane in the presence of Pd(PPh₃)₂Cl₂ and triethylamine in THF at room temperature for 1 h. The resulting compounds were not isolated but immediately submitted to the deprotection-cycloaddition protocol and reaction with benzyl azide in the presence of CuF₂ or the TBAF/CuI afforded 1-benzyl,4-acyl-1*H*-1,2,3-triazoles (7). The Sonogashira cross-coupling reaction was found to be sensitive to the nature of the employed acyl chloride and the use of electron-rich acyl chlorides (entries 2 and 3 vs entry 4) gave the best conversions of the starting materials. Full conversions of the 1-substituted 4-acyl-1*H*-1,2,3-triazoles were achieved when TMS-modified ynones were purified prior to reaction with azides, establishing the Sonogashira coupling of the acyl chlorides with ethynyltrimethylsilane as the yield-limiting step of the one-pot, three-step protocol.

It is important to note that 1-substituted 4-acyl-1*H*-1,2,3-triazoles have rarely been prepared by CuAAC due to side product formation.¹⁶ Previous syntheses of this class of triazoles involved classical thermal Huisgen 1,3-dipolar cycloadditions with the inconvenience of separating the two triazole regioisomers^{17a} and condensation of amines with α -diazo-1,3-dicarbonyls.^{16b} The microwave conditions reported here provide easy and regioselective access to this class of biomedically important compounds.

In conclusion, an expedient one-pot, three-step protocol has been developed for the preparation of 1,4-disubstituted triazoles involving Sonogashira cross-coupling to generate *in-situ* a variety of TMS-protected alkynes, which could immediately be desilylated and reacted with azides in the presence of a mixture of TBAF/CuI or CuF₂. The reaction sequence could be completed within 1 h by employing microwave-assisted heating. Furthermore, we have shown that 2-yn-1-ones, which were obtained by Sonogashira cross-coupling of aromatic acyl chlorides with ethynyltrimethylsilane, undergo Cu-catalyzed cycloadditions with azides when subjected to microwave heating to give regioselective formation of 1-substituted 4-phenylacyl-1*H*-1,2,3-triazoles. The new one-pot three-step process will make it possible to rapidly prepare compound libraries for drug discovery programs as it avoids time consuming and costly purification protocols of synthetic intermediates. Furthermore, it is to be expected that the one-pot procedure can be combined with *in-situ* formation of azides⁵ thereby further increasing the convenience of CuAAC.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by the National Cancer Institute of the US National Institutes of Health (Grant No. R01 CA88986, G.-J.B.).

References

1. For the original publications, see: a) Tornøe CW, Christensen C, Meldal M. J. Org. Chem. 2002; 67:3057. [PubMed: 11975567] b) Rostovtsev VV, Green LG, Fokin VV, Sharpless KB. Angew. Chem. Int. Ed. 2002; 41:2596.
2. a) Pieters RJ, Rijkers DTS, Liskamp RM. J. QSAR Comb. Sci. 2007; 26:1181. b) Nájera C, Sansano JM. Org. Biomol. Chem. 2009; 7:4567. [PubMed: 19865689] c) van Dijk M, Rijkers DTS, Liskamp RMJ, van Nostrum CF, Hennink WE. Bioconjugate Chem. 2009; 20:2001.

3. a) Lutz J-F. *Angew. Chem. Int. Ed.* 2007; 46:1018. b) Golas PL, Matyjaszewski K. *Chem. Soc. Rev.* 2010; 39:1338. [PubMed: 20309490]
4. a) Moorhouse AD, Moses JE. *Chem. Med. Chem.* 2008; 3:715. [PubMed: 18214878] b) Tron GC, Pirali T, Billington RA, Canonico PL, Sorba G, Genazzani AA. *Med. Res. Rev.* 2008; 28:278. [PubMed: 17763363] c) Mamidyala SK, Finn MG. *Chem. Soc. Rev.* 2010; 39:1252. [PubMed: 20309485]
5. a) Dururgkar KA, Gonnade RG, Ramana CV. *Tetrahedron.* 2009; 65:3974. b) Feldman AK, Colasson B, Fokin VV. *Org. Lett.* 2004; 6:3897. [PubMed: 15496058] c) Ackermann L, Potukuchi HK, Landsberg D, Vicente R. *Org. Lett.* 2008; 10:3081. [PubMed: 18549230] d) Appukkuttan P, Dehaen W, Fokin VV, Van der Eycken E. *Org. Lett.* 2004; 6:4223. [PubMed: 15524448] e) Yadav JS, Reddy BVS, Reddy M, Chary DN. *Tetrahedron Lett.* 2007; 48:8773. f) Campbell-Verduyn LS, Szymanski W, Postema CP, Dierckx RA, Elsinga PH, Janssen DB, Feringa BL. *Chem. Commun.* 2010; 46:898. g) Tao C-Z, Cui X, Li J, Liu A-X, Liu L, Guo Q-X. *Tetrahedron Lett.* 2007; 48:3525.
6. a) Beckmann HSG, Wittmann V. *Org. Lett.* 2007; 9:1. [PubMed: 17192070] b) Barral K, Moorhouse AD, Moses JE. *Org. Lett.* 2007; 9:1809. [PubMed: 17391043] c) Lee C-T, Huang S, Lipshutz BH. *Adv. Synth. Catal.* 2009; 351:3139.
7. Valverde IE, Delmas AF, Aucagne V. *Tetrahedron.* 2009; 65:7597.
8. Aucagne V, Leigh DA. *Org. Lett.* 2006; 8:4505. [PubMed: 16986936]
9. a) Kappe CO, Van der Eycken E. *Chem. Soc. Rev.* 2010; 39:1280. [PubMed: 20309486] b) Appukkuttan P, Mehta V, Van der Eycken E. *Chem. Soc. Rev.* 2010; 39:1467. [PubMed: 20419202]
10. Heating under microwave irradiation trimethyl(phenylethynyl)silane (**1a**) with benzyl azide (**2a**) in the presence of TBAF (1 equiv) at 120 °C for 20 min resulted in less than 5% yield of the corresponding triazole in a 1:1 mixture of regioisomers.
11. Mechanistic studies and DFT calculations support the notion that CuF₂ catalyzes the cycloaddition. Friscourt, F.; Ledin, PA.; Boons, G-J. Abstracts of Papers, 239th ACS National Meeting; San Francisco, CA, USA. March 21-25; 2010. manuscript in preparation
12. a) Sonogashira K, Tohda Y, Hagihara N. *Tetrahedron Lett.* 1975; 16:4467. b) Tohda Y, Sonogashira K, Hagihara N. *Synthesis.* 1977:777.
13. a) Kappe CO. *Angew. Chem. Int. Ed.* 2004; 43:6250. b) Nilsson P, Olofsson K, Larhed M. *Top Curr. Chem.* 2006; 266:103.
14. Karpov AS, Muller TJJ. *Org. Lett.* 2003; 5:3451. [PubMed: 12967297]
15. Sashida H. *Synthesis.* 1998:745.
16. Synthesis of 1-substituted 4-acyl-1*H*-1,2,3-triazoles using CuAAC exhibits moderate yield (< 60%), see: a) Chassaing S, Kumarraja M, Sido ASS, Pale P, Sommer J. *Org. Lett.* 2007; 9:883. [PubMed: 17286410] b) Xie J, Seto CT. *Bioorg. Med. Chem.* 2007; 15:458. [PubMed: 17046267] c) Pardin C, Roy I, Lubell WD, Keillor JW. *Chem. Biol. Drug Des.* 2008; 72:189. [PubMed: 18715232]
17. a) Calderone V, Giorgi I, Livi O, Martinotti E, Mantuano E, Martelli A, Nardi A. *Eur. J. Med. Chem.* 2005; 40:521. [PubMed: 15922836] b) Dabak K, Sezer O, Akar A, Anaç O. *Eur. J. Med. Chem.* 2003; 38:215. [PubMed: 12620665]

Scheme 1.

One-pot three-step synthesis of 1,2,3-triazoles by *in-situ* formation of a terminal alkyne by Sonogashira cross-coupling and desilylation followed by a cycloaddition with an azide.

Table 1

Microwave-assisted one-pot TMS-deprotection of compounds **1a-g** and cycloaddition with azides **2a-c** to give triazoles **3a-i**

entry	R	R'	method A ^a yield (%)	method B ^b yield (%)
1	C ₆ H ₅	Bn	98	98 ^c
2	4-MeO-C ₆ H ₄	Bn	97	96
3	4-CF ₃ -C ₆ H ₄	Bn	98	97
4	4-Cl-C ₆ H ₄	Bn	89	95
5	4-Br-C ₆ H ₄	Bn	81	91
6	2-Br-C ₆ H ₄	Bn	89	90
7	CH ₂ OH	Bn	82	98
8	C ₆ H ₅	4-MeO-Bn	87	98
9	C ₆ H ₅	4-NO ₂ -Bn	98	99

^a A mixture of TBAF (1.0 M, 2.0 equiv) and CuI (10 mol %) and DIPEA (20 mol %) was used

^b CuF₂ (2.0 equiv) was used

^c **3a** was isolated in yields of 22% and 57% when the reaction was performed at 60 °C or 90 °C, respectively.

Table 2

One-pot, three-step synthesis of 1,4-disubstituted triazoles 3

entry	Ar-X	R	method A ^a yield (%)	method B ^b yield (%)
1	Ph-I	Bn	96	91
2	4-MeO-C ₆ H ₄ -I	Bn	41	54
3	4-CF ₃ -C ₆ H ₄ -I	Bn	97	98
4	4-F-C ₆ H ₄ -I	Bn	98	98
5	4-Me-C ₆ H ₄ -I	Bn	93	95
6	2-Me-C ₆ H ₄ -I	Bn	97	98
7	Ph-Br	Bn	42	65
8	4-MeO-C ₆ H ₄ -Br	Bn	13	15(57 ^c)
9	4-NO ₂ -C ₆ H ₄ -Br	Bn	12	11(53 ^c)
10	Ph-I	4-MeO-Bn\	84	91
11	Ph-I	4-NO ₂ -Bn	97	98

^a A mixture of TBAF (1.0 M, 2.0 equiv) and CuI (10 mol %) was used^b CuF₂ (2.0 equiv) was used^c The Sonogashira cross coupling was performed using Pd(PPh₃)₂Cl₂ (5 mol %), CuI (10 mol %) in refluxing Et₃N for 10 h.

Table 3One-pot, three-step synthesis of 1-substituted 4-acyl-1*H*-1,2,3-triazoles **7a-e**

entry	Ar	R	method A ^a yield (%)	method B ^b yield (%)
1	Ph	Bn	40	42
2	4-MeO-C ₆ H ₄	Bn	80	83
3	4-F-C ₆ H ₄	Bn	82	88
4	4-NO ₂ -C ₆ H ₄	Bn	32	36
5	2-Me-C ₆ H ₄	Bn	35	34

^a A mixture of TBAF (1.0 M, 2.0 equiv) and CuI (10 mol %) was used^b CuF₂ (2.0 equiv) was used.