Published in final edited form as:

Neurosci Lett. 2010 December 17; 486(3): 207–210. doi:10.1016/j.neulet.2010.09.054.

Psychophysical stress increases the expression of phospho-CREB, Fos protein and neurokinin-1 receptors in superficial laminae of trigeminal subnucleus caudalis in female rats

Sara. L. Duenes, Randy Thompson, Zheng Chang, Keiichiro Okamoto*, and David. A. Bereiter

Dept. Diagnostic and Biological Sciences, University of Minnesota School of Dentistry, 18-214 Moos Tower, 515 Delaware St SE, Minneapolis, MN 55455

Abstract

Psychological stress and estrogen status are risk factors to develop painful temporomandibular joint disorders (TMJD); however, the neural basis for this relationship is not known. This study tested the hypothesis that repeated forced swim stress and estradiol treatment alter the phosphorylation of cAMP responsive element binding protein (pCREB) in trigeminal subnucleus caudalis (Vc), the initial site of sensory input from the TMJ. Ovariectomized female rats were given low or high dose estradiol and subjected to repeated forced swim stress for three days and on day 4 an intra-TMJ injection of mustard oil or vehicle was given. Forced swim alone increased the number of pCREB-positive neurons, independent of estradiol treatment or TMJ stimulation, in superficial and deep laminae of Vc. Forced swim also increased the number of Fos-positive neurons in superficial laminae and neurokinin-1 receptor mRNA in whole dorsal Vc, independent of estradiol treatment. These results indicated that persistent psychophysical stress alone was sufficient to increase the expression of pCREB and downstream regulated genes associated with enhanced excitability in the caudal medullary dorsal horn, a brainstem region thought to be critical for TMJD pain.

Keywords

trigeminal subnucleus caudalis; forced swim stress; pCREB; Fos; neurokinin-1 receptor; estradiol

Temporomandibular joint disorders (TMJD) are common conditions characterized by spontaneous pain in the temporomandibular joint (TMJ) region and muscles of mastication. Efforts to understand the neural mechanisms for persistent painful TMJD have been limited due to lack of suitable animal models. Although diverse etiologies may contribute to the development of TMJD, psychological stress and estrogen status are recognized as significant risk factors [10,14,24]. Persistent psychological stress exacerbates evoked nociceptive behavior in animal models for spinal [11], visceral [3] and TMJD pain [8]; however, most studies have used only male animals. The present study used repeated forced swim stress (FS) in female rats, an established psychophysical stress that causes persistent cutaneous and

*Correspondence: Keiichiro Okamoto, D.D.S., Ph.D., Dept. of Diagnostic and Biological Sciences, University of Minnesota School of Dentistry, 18-214 Moos Tower, 515 Delaware St SE, Minneapolis, MN, 55455. Phone: (612) 625-4447; Fax: (612) 626-2651, okamo007@umn.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

muscle hyperalgesia in male rats [22,25]. In spinal dorsal horn, phosphorylation of the transcription factor, cAMP response element-binding protein (pCREB), regulates the expression of the immediate early gene *c-fos* [13] and late response genes such as the neurokinin-1 (NK-1) receptor [23] and underlies long-term changes in synaptic activity [4]. Blockade of pCREB formation at spinal levels prevents hyperalgesia in animal models of muscle pain [9], while destruction of spinal NK-1 receptors prevents the sensitization of dorsal horn neurons and hyperalgesia to cutaneous stimuli [17]. The present study determined if repeated FS and estradiol treatment interact to alter the expression of pCREB and downstream regulated genes at the trigeminal subnucleus caudalis/cervical spinal cord (Vc/C₁₋₂) junction, a medullary dorsal horn region that processes direct input from the TMJ region in an estrogen-dependent manner [20,29] in female rats.

The experimental protocols were approved by the Institutional Animal Care and Use Committee of the University of Minnesota and conformed to the established guidelines set by The National Institute of Health guide for the care and use of laboratory animals (Publications No. 99-158, revised 2002). Ovariectomized (OvX) female rats (225-300 g, Sprague-Dawley, Harlan, Indianapolis, IN) were injected with low (LE, 3 µg/kg) or high (HE, 30 µg/kg, sc) dose 17β-estradiol-3-benzoate (E2, Sigma, St. Louis, MO) dissolved in sesame oil for three days. The estrogen status of each rat was confirmed on the day of the experiment by the cytology of a vaginal smear sample; LE rats had mainly (> 80%) small nucleated leukocytes and HE rats displayed mostly large nucleated epithelial cells. The FS protocol involved placing the rat in a plastic cylinder (30 × 50 cm) containing 20 cm water (25°C) for 10 min per day for three days [22], while sham controls were placed in an empty cylinder. Immobility time, an index of behavioral stress, was measured during each swim session [21], whereas forelimb grip force, an estimate of muscle pain or weakness [16] was measured before each swim session.

Rats received daily injections of E2, grip force measurements and a session of FS or sham swimming for 3 days. On day 4, rats were anesthetized with pentobarbital sodium (65 mg/kg, ip) and those animals assayed for pCREB were further given an intra-TMJ injection of the small fiber excitant mustard oil (20% v/v) or vehicle (mineral oil) and allowed to survive for 30 min. Animals assayed for the immediate early gene product, Fos protein, or NK-1 receptor mRNA received no TMJ injection and were killed under anesthesia. Rats processed for pCREB and Fos immunohistochemistry were perfused through the heart with heparinized saline, followed by cold fixative (4 % paraformaldehyde, 0.1 M phosphate, pH 7.4), the lower brainstem was removed and post-fixed overnight at 4°C. Transverse sections (50 µm) were cut on a vibratome, blocked in a solution of 0.1 M PBS, 0.3% Triton X-100, and 5% normal donkey serum for 60 min and incubated in primary antiserum for pCREB (Upstate, Temecula, CA; 1:3500) or Fos protein (Ab-5, Millipore; 1: 15000) for 40 h at 4°C. After rinsing in PBS for 20 min, sections were incubated with biotinylated donkey anti-rabbit antibodies (Chemicon-Millipore, Temecula, CA, USA; 1:300, 115 min) and avidin-biotin-peroxidase complex (Vector, Burlingame, CA, USA; 60 min). pCREB- and Fos-positive neurons were visualized by incubation in a 0.025% diaminobenzidine solution containing 0.01% nickel ammonium sulfate and 0.125% cobalt chloride, activated by 0.01% hydrogen peroxide. Specific staining for pCREB and Fos was abolished by omission of primary antisera. pCREB-positive neurons were counted at 200× using a calibrated reticle of 640 µm² positioned over the medial superficial or deeper laminae at the trigeminal subnucleus caudalis/upper cervical cord dorsal horn (Vc/C₁₋₂) region (see Fig 2A). We selected specific areas of the Vc/C₁₋₂ region to assess the effects of forced swimming conditioning on TMJ-evoked pCREB induction, because these areas are involved in TMJ-evoked nociceptive processing [28,29]. As a control for general pCREB expression we also counted cells in ventral horn at the same Vc/C₁₋₂ region. Five sections separated by 100 µm at the Vc/C₁₋₂ region were counted from each treatment group. The Vc/C₁₋₂ region is

located approximately 4.5 to 6.5 mm caudal to the obex [30]. In separate rats, Fos-positive neurons at the Vc/C₁₋₂ region were counted in superficial laminae (I-II), deeper laminae (III-V) using established cytological landmarks [19]. Cell counts for pCREB and Fos were compared across groups by analysis of variance (ANOVA) without prior knowledge of treatment. Individual comparisons were made by the Newman-Keuls test after ANOVA. Possible relationships between pCREB cell counts and grip force in FS and sham swim animals were assessed by Spearman's rank-order correlation. All data are presented as mean $\pm$ SEM.

NK-1 receptor mRNA was assessed by quantitative RT-PCR from fresh tissue. Under deep anesthesia the lower brainstem was removed, dorsal horn tissue dissected bilaterally and total RNA extracted (Absolutely RNA Miniprep Kit, Stratagene, La Jolla, CA). RNA was quantified by spectrophotometry, reverse transcribed (iScript cDNA synthesis Kit, Bio Rad, Hercules, CA) and quantified by a Chromo4 DNA cycler using IQ SYBR green Supermix (Bio Rad). Samples were run in triplicate 25 μ L reactions using 2 μ L of the 20 μ L cDNA synthesis as template. Genes of interest were run against a normalizing gene (GAPDH) allowing utilization of the ΔC_T method for relative quantification of gene expression. PCR reaction parameters were: initial denaturation at 95°C for 3 min, 40 cycles of 95°C for 10 s, 58.5°C for 30 s, and 72°C for 45 s. A final melting curve from 55°C to 90°C was performed for each reaction to ensure amplicon validity. Primers were designed using Primer3 software (http://frodo.wi.mit.edu/cgi-bin/primer3/primer3_www.cgi), synthesized by Sigma Genosys (St. Louis, MO) and checked against the NCBI database to ensure gene and species specificity. Primer sets employed were: GAPDH (F: 5' agacagccgcattcttctgt 3', R: 5' cttgccgtggtagagtcac 3') and NK-1 receptor (F: 5' gctgtcttccttctggaac 3' R: 5' ccacacaggaagtgggaagt 3').

Repeated FS reduced forelimb grip force by day 4 ($F_{3,69} = 8.06$, $P < 0.001$) similarly in LE- and HE-treated rats (Fig 1A). Sham conditioning did not affect forelimb grip force ($P > 0.05$). Although a reduction in grip force may have been due to muscle pain or weakness, we cannot exclude that other factors such as local release of agents in working muscle or central drive were involved [1,16]. Immobility times also were increased maximally by day 3 of FS conditioning in both LE and HE rats (data not shown). The number of pCREB-positive neurons in superficial laminae at the Vc/C₁₋₂ region increased significantly ipsilateral to intra-TMJ injection of mustard oil in sham swim rats and bilaterally after FS alone or FS plus TMJ stimulation (Fig 3A, $F_{7,24} = 18.6$, $p < 0.001$). As seen in the example of Fig 2B, FS also increased the number of pCREB-positive cells in deep laminae after FS (Fig 3B, $F_{7,24} = 6.07$, $p < 0.001$). However, TMJ stimulation in sham swim animals did not cause further increases pCREB expression in deep laminae. By contrast, estrogen status did not influence pCREB in superficial or deep laminae in any treatment group (Fig 3). It was not likely that FS caused widespread increases in pCREB throughout the brainstem since cell counts in ventral horn at the Vc/C₁₋₂ region were similar to that of sham swim rats (41 ± 2 versus 40 ± 1 pCREB cells/640 μ m²). Correlation analyses revealed a significant inverse relationship between pCREB cell counts in superficial laminae and maximum forelimb grip force on the day of the experiment (Fig 1B, $r_s = -.556$, $p < 0.005$). A similar analysis between pCREB cell counts in deep laminae and grip force also indicated a significant but weak relationship ($r_s = -.423$, $p < 0.03$). To assess the effect of FS on downstream genes regulated by pCREB, we found that the number of Fos-positive neurons in superficial ($F_{3,12} = 10.85$, $p < 0.001$), but not deep laminae, was increased after FS independent of estrogen status ($F_{3,12} = 0.65$, $p > 0.05$, Fig 4A). Similarly, NK-1 receptor expression in whole dorsal horn was increased significantly after FS alone, but was not affected by estrogen status (Fig 4B).

The main finding of this study was that repeated FS alone was sufficient to markedly increase the number of pCREB-positive neurons bilaterally in superficial (3-fold) and deep laminae (2-fold) at the Vc/C₁₋₂ junction. Although TMJ injection of mustard oil also increased pCREB ipsilateral to the stimulus in superficial laminae of sham swim rats, the number of pCREB cells in superficial and deep laminae after FS plus TMJ stimulation was not different from that seen after FS alone. This suggested that repeated psychophysical stress and TMJ nociceptors targeted a similar subpopulation of neurons at the Vc/C₁₋₂ junction, a dorsal horn region closely associated with the integration of sensory input from the TMJ [2,28,29].

By contrast, estrogen status did not affect the number of pCREB-positive neurons produced after FS or TMJ stimulation. This differed somewhat from previous reports in which we found that TMJ-evoked unit activity [29] and the number of Fos-positive neurons in superficial laminae at the Vc/C₁₋₂ junction of HE-treated rats were increased compared to LE rats [20]. The reason for the lack of an estrogen effect on pCREB staining at the Vc/C₁₋₂ junction is not certain. The pCREB response to E2 was regionally selective in different limbic brain areas and depended on the duration of treatment [5,7]. Time course studies revealed that E2-induced pCREB formation in forebrain was transient and reached a peak within 1 h and returned towards control levels by 4 h [27]. Since the last E2 injection was given ~24 h prior to brainstem removal, we cannot exclude that differential effects of HE and LE treatment on pCREB formation may have occurred at earlier times. Similarly, noxious stimulus-induced pCREB also was produced rapidly and transiently in spinal dorsal horn of male rats [15,18]. The transient nature of pCREB formation in response to other treatments suggested that repeated FS must exert a persistent influence on neurons at the Vc/C₁₋₂ junction since the last swim session was 24 h prior to tissue collection.

One possible interpretation for FS-induced pCREB formation in superficial laminae neurons at the Vc/C₁₋₂ junction concerns the recruitment of descending nociceptive controls. This was supported by findings that the number of pCREB-positive cells in superficial laminae was inversely correlated with grip force, whereas Fos-positive neurons in superficial laminae and NK-1 receptor expression in whole dorsal horn samples were increased after repeated FS. In spinal dorsal horn there was a close relationship between NK-1 receptor-positive neurons and Fos production after noxious stimuli [6], whereas in Vc many NK-1 receptor cells projected to periaqueductal gray a key relay nucleus for descending nociceptive controls [18]. Selective destruction of NK-1 receptor positive spinal lamina I neurons greatly reduced nociceptive behavior, a result reproduced by pharmacological blockade of 5HT₃ receptors in controls but not NK-1 receptor lesioned rats [26]. Chronic stress increased the number of NK-1 receptor positive neurons in superficial laminae of spinal dorsal horn in a visceral pain model [3], while repeated FS enhanced the late phase of formalin-evoked nocifensive behavior and was prevented by rostroventromedial medullary lesion [12]. Upregulation of pCREB underlies many forms of long-term changes in synaptic plasticity throughout the brain [4]. The present study suggests that pCREB formation may play a critical role in modulating dorsal horn neural activity in an animal model for TMJD pain.

Research Highlights

Psychophysical stress increases pCREB expression at trigeminal subnucleus caudalis. Increased pCREB expression after stress is not affected by E2 status and TMJ injury. Psychophysical stress alone increases Fos and NK1R mRNA expression. Psychophysical stress causes persistent activation at trigeminal caudalis neurons.

Acknowledgments

This study was supported by a grant from the National Institute of Dental and Craniofacial Research DE12758 and the Office of Research on Women's Health (ORWH).

References

1. Amann M, Proctor LT, Sebranek JJ, Pegelow DF, Dempsey JA. Opioid-mediated muscle afferents inhibit central motor drive and limit peripheral muscle fatigue development in humans. *J Physiol.* 2009; 587:271–83. [PubMed: 19015193]
2. Bereiter DA, Benetti AP. Amino acid release at the spinomedullary junction after inflammation of the TMJ region in male and female rats. *Pain.* 2006; 126:175–183. [PubMed: 16901647]
3. Bradesi S, Kokkotou E, Simeonidis S, Patierno S, Ennes HS, Mittal Y, McRoberts JA, Ohning G, McLean P, Marvizon JC, Sternini C, Pothoulakis C, Mayer EA. The role of neurokinin 1 receptors in the maintenance of visceral hyperalgesia induced by repeated stress in rats. *Gastroenterology.* 2006; 130:1729–1742. [PubMed: 16697737]
4. Carlezon WA Jr, Duman RS, Nestler EJ. The many faces of CREB. *Trends Neurosci.* 2005; 28:436–45. [PubMed: 15982754]
5. Carlstrom L, Ke ZJ, Unnerstall JR, Cohen RS, Pandey SC. Estrogen modulation of the cyclic AMP response element-binding protein pathway. Effects of long-term and acute treatments. *Neuroendocrinology.* 2001; 74:227–43. [PubMed: 11598379]
6. Doyle CA, Hunt SP. Substance P receptor (neurokinin-1) expressing neurons in lamina I of the spinal cord encode for the intensity of noxious stimulation: a c-fos study in rat. *Neuroscience.* 1999; 89:17–28. [PubMed: 10051214]
7. Fan L, Hanbury R, Pandey SC, Cohen RS. Dose and time effects of estrogen on expression of neuron-specific protein and cyclic AMP response element-binding protein and brain region volume in the medial amygdala of ovariectomized rats. *Neuroendocrinology.* 2008; 88:111–26. [PubMed: 18446018]
8. Gameiro GH, Andrade Ada S, de Castro M, Pereira LF, Tambeli CH, Veiga MC. The effects of restraint stress on nociceptive responses induced by formalin injected in rat's TMJ. *Pharmacol Biochem Behav.* 2005; 82:338–44. [PubMed: 16213578]
9. Hoeger-Bement MK, Sluka KA. Phosphorylation of CREB and mechanical hyperalgesia is reversed by blockade of the cAMP pathway in a time-dependent manner after repeated intramuscular acid injections. *J Neurosci.* 2003; 23:5437–45. [PubMed: 12843242]
10. Huang GJ, LeResche L, Critchlow CW, Martin MD, Drangsholt MT. Risk factors for diagnostic subgroups of painful temporomandibular disorders (TMD). *J Dent Res.* 2002; 81:284–288. [PubMed: 12097315]
11. Imbe H, Iwai-Liao Y, Senba E. Stress-induced hyperalgesia: animal models and putative mechanisms. *Front Biosci.* 2006; 11:2179–92. [PubMed: 16720304]
12. Imbe H, Okamoto K, Donishi T, Senba E, Kimura A. Involvement of descending facilitation from the rostral ventromedial medulla in the enhancement of formalin-evoked nociceptive behavior following repeated forced swim stress. *Brain Res.* 2010; 1329:103–12. [PubMed: 20226771]
13. Ji RR, Rupp F. Phosphorylation of transcription factor CREB in rat spinal cord after formalin-induced hyperalgesia: relationship to c-fos induction. *J Neurosci.* 1997; 17:1776–1785. [PubMed: 9030636]
14. Kanehira H, Agariguchi A, Kato H, Yoshimine S, Inoue H. Association between stress and temporomandibular disorder. *Nihon Hotetsu Shika Gakkai Zasshi.* 2008; 52:375–80. [PubMed: 18678971]
15. Kawasaki Y, Kohno T, Zhuang ZY, Brenner GJ, Wang H, Van Der Meer C, Befort K, Woolf CJ, Ji RR. Ionotropic and metabotropic receptors, protein kinase A, protein kinase C, and Src contribute to C-fiber-induced ERK activation and cAMP response element-binding protein phosphorylation in dorsal horn neurons, leading to central sensitization. *J Neurosci.* 2004; 24:8310–21. [PubMed: 15385614]
16. Kehl LJ, Trempe TM, Hargreaves KM. A new animal model for assessing mechanisms and management of muscle hyperalgesia. *Pain.* 2000; 85:333–43. [PubMed: 10781907]

17. Khasabov SG, Rogers SD, Ghilardi JR, Peters CM, Mantyh PW, Simone DA. Spinal neurons that possess the substance P receptor are required for the development of central sensitization. *J Neurosci.* 2002; 22:9086–98. [PubMed: 12388616]
18. Li JL, Ding YQ, Xiong KH, Li JS, Shigemoto R, Mizuno N. Substance P receptor (NK1)-immunoreactive neurons projecting to the periaqueductal gray: distribution in the spinal trigeminal nucleus and the spinal cord of the rat. *Neurosci Res.* 1998; 30:219–25. [PubMed: 9593332]
19. Molander C, Xu Q, Rivero-Melian C, Grant G. Cytoarchitectonic organization of the spinal cord in the rat: II. The cervical and upper thoracic cord. *J Comp Neurol.* 1989; 289:375–385. [PubMed: 2808773]
20. Okamoto K, Bereiter DF, Thompson R, Tashiro A, Bereiter DA. Estradiol replacement modifies c-fos expression at the spinomedullary junction evoked by temporomandibular joint stimulation in ovariectomized female rats. *Neuroscience.* 2008; 156:729–736. [PubMed: 18765271]
21. Porsolt RD, Le Pichon M, Jalfre M. Depression: a new animal model sensitive to antidepressant treatments. *Nature.* 1977; 266:730–2. [PubMed: 559941]
22. Quintero L, Moreno M, Avila C, Arcaya J, Maixner W, Suarez-Roca H. Long-lasting delayed hyperalgesia after subchronic swim stress. *Pharmacol Biochem Behav.* 2000; 67:449–58. [PubMed: 11164071]
23. Seybold VS, McCarron KE, Mermelstein PG, Groth RD, Abrahams LG. Calcitonin gene-related peptide regulates expression of neurokinin1 receptors by rat spinal neurons. *J Neurosci.* 2003; 23:1816–24. [PubMed: 12629185]
24. Slade GD, Diatchenko L, Bhalang K, Sigurdsson A, Fillingim RB, Belfer I, Max MB, Goldman D, Maixner W. Influence of psychological factors on risk of temporomandibular disorders. *J Dent Res.* 2007; 86:1120–5. [PubMed: 17959908]
25. Suarez-Roca H, Quintero L, Arcaya JL, Maixner W, Rao SG. Stress-induced muscle and cutaneous hyperalgesia: differential effect of milnacipran. *Physiol Behav.* 2006; 88:82–7. [PubMed: 16624346]
26. Suzuki R, Morcuende S, Webber M, Hunt SP, Dickenson AH. Superficial NK1-expressing neurons control spinal excitability through activation of descending pathways. *Nat Neurosci.* 2002; 5:1319–26. [PubMed: 12402039]
27. Szego EM, Barabas K, Balog J, Szilagyi N, Korach KS, Juhasz G, Abraham IM. Estrogen induces estrogen receptor alpha-dependent cAMP response element-binding protein phosphorylation via mitogen activated protein kinase pathway in basal forebrain cholinergic neurons in vivo. *J Neurosci.* 2006; 26:4104–4110. [PubMed: 16611827]
28. Takeshita S, Hirata H, Bereiter DA. Intensity coding by TMJ-responsive neurons in superficial laminae of caudal medullary dorsal horn of the rat. *J Neurophysiol.* 2001; 86:2393–2404. [PubMed: 11698529]
29. Tashiro A, Okamoto K, Milam SB, Bereiter DA. Differential effects of estradiol on encoding properties of TMJ units in laminae I and V at the spinomedullary junction in female rats. *J Neurophysiol.* 2007; 98:3242–3253. [PubMed: 17928557]
30. Yoshida A, Dostrovsky JO, Sessle BJ, Chiang CY. Trigeminal projections to the nucleus submedius of the thalamus in the rat. *J Comp Neurol.* 1991; 307:609–625. [PubMed: 1714465]

Figure 1.

Repeated FS conditioning reduces maximum forelimb grip force. A. Grip force was reduced similarly in LE- and HE-treated rats. Sample sizes: LE = 12; HE = 14. * $p < 0.05$, ** $p < 0.01$ versus day 1 value. B. Reduction in grip force after FS was inversely correlated with the number of pCREB-positive neurons in superficial laminae. $n = 26$.

Figure 2. Examples of pCREB staining at the Vc/C₁₋₂ junction in LE-treated rats. A. Control. B. Repeated FS condition. Boxed areas represent the superficial and deep laminar regions for cell counting (640 μm^2). Calibration = 150 μm .

Figure 3.

Summary of pCREB cell counts at the Vc/C₁₋₂ junction. A. superficial laminae. B. deeper laminae. Abbreviations: E2, estradiol benzoate; FS, forced swim; HE, high E2; LE, low E2; MO, mustard oil. * $p < 0.05$, ** $p < 0.01$ ipsilateral versus contralateral; a = $p < 0.05$, b = $p < 0.01$ 0% versus 20% MO for the equivalent E2-treated non-FS rats; ## = $p < 0.01$ versus corresponding sham swim groups. n = 4-6 rats per group.

Figure 4.

Repeated FS increases the number of Fos-positive neurons in superficial laminae at the Vc/C₁₋₂ junction (A) and NK-1 mRNA in whole Vc/C₁₋₂ tissue samples (B). a = $p < 0.05$, b = $p < 0.01$ versus sham swim groups. n = 4-6 rats per group.