CORRESPONDENCE

Radon in Indoor Spaces—An Underestimated Risk Factor for Lung Cancer in Environmental Medicine

by PD Dr. med. Klaus Schmid, Prof. Dr. med. Torsten Kuwert,
Prof. Dr. med. Hans Drexler in volume 11/2010

3. Schmid K, Kuwert T, Drexler H: Radon in indoor spaces—An underestimated risk factor for lung cancer in environmental medicine. *Dtsch Arztebl Int* 2010; 107(11): 181–6.

Dr. med. Helmut Schlarb
Antoniusplatz 1
49661 Cloppenburg, Germany

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Radon Balneology

Curative waters were regarded as containing radon only if the concentration was higher than 18 nanocuries. For inhalation, spa towns were still happy with an aerosol concentration of about 5 nanocuries/liter. Gas was released from natural spring water, and people taking spa cures were placed under perspex covers at their inhalation seats. The main indications were rheumatic disorders (1, 2).

The Erlangen-Nürnberg working group pointed out twice in their article (3) that emanations other than from uranium mines were not their subject. However, they none the less requested a rigorous risk-benefit analysis of such emanations. However, if they then categorize radon as “the second most important cause of lung cancer [after cigarette smoking] in the general population without occupational exposure,” then readers of medical journals might expect to see a clear position on curative treatments using radon.

In my place of birth we often played in a radon cave, whose entrance had been destroyed in the second world war. We grew up in a wine growing area, and instead of torches we used candles in the long tunnel (where concentrations of carbon dioxide are high the flame will extinguish). Later at school we learnt that if spring water flows over radium containing rock (Ra 223/half life 1600 years), the first resulting breakdown product to become dissolved in the water is radon (Rn 222). The noble gas radon has a half life of 3.28 days and is said to be soluble not only in water and blood but also in fat. Depending on the breathing techniques of test volunteers and on environmental temperatures, concentrations of between 0.6 and 1.6 picocuries/ml blood have been measured in the Bockstein thermal healing cave at Bad Gastein.

DOI: 10.3238/arztebl.2010.0730a

REFERENCES

1. Dirnagl K: Die Technik der Radoninhalation. *Naheland-Kalender* 1984; 169–70.
2. Braunbehrens H v.: Über Radonkuren. *Naheland-Kalender* 1984; 171–2.

Radon in Drinking Water

Firstly, we wish to express to the authors our appreciation of and thanks for their article. They showed the risk for lung cancer while focusing in their discussion mainly on inhaled radon, which penetrates into buildings and residential spaces from underground.

We wish to point out a further source of cancer risk, which we think has not received enough attention in the public discussion—namely, the release of radon that is dissolved in drinking water. In Germany, the mean radon activity in drinking water is about 6 Bq/L, although regionally it may exceed 100 Bq/L and even 1000 Bq/L. What happens if the gas is released from the water into the shower cubicle during a warm shower and then inhaled? For a concentration of 100 Bq/L, activity in the shower cubicle will rise by 100 Bq per liter of water used if the radon is fully released (activity describes the number of radioactive breakdowns per second [1 Bq=1 breakdown/second]). For the purposes of describing exposures, using Bq/L or Bq/m³ as the unit of measurement seems appropriate [2]). A few years ago, an American brochure caused shockwaves by reporting values of several kBq/m³ that had been measured in shower cubicles after 15 minutes—a value that is entirely plausible according to the calculation in our example. Is the risk posed by radon inhaled during a shower really negligible, as is often claimed, especially as ventilating the cubicle or cabin during the shower is usually difficult?

In the kitchen, the evaporation of steam and the radon concentrated in natural gas constitute an additional radon exposure. Opening the windows is a useful counter-measure in this context. In modern residences with tightly sealed windows, however, thorough ventilation is more problematic.

We would welcome a short comment from the authors, outlining their views on the problems of “radon in shower cubicles” and “lacking ventilation in modern residential buildings.” As a result of insufficient ventilation, several other causes of health risks have made the newspaper headlines, including pathogenic indoor climates, concentration of allergens, or fungal (mould) overgrowth.

DOI: 10.3238/arztebl.2010.0730b

REFERENCES

1. Schmid K, Kuwert T, Drexler H: Radon in indoor spaces—An underestimated risk factor for lung cancer in environmental medicine. *Dtsch Arztebl Int* 2010; 107(11): 181–6.
2. www.gesundheitsamt.de/alle/umwelt/physik/strahl/ion/ra/m

Prof. Dr. rer. nat. Hermann Dertinger

Prof. Dr. med. Konrad Andrassy

Institut für Biologische Grenzflächen

KIT – Campus Nord (vormals Forschungszentrum Karlsruhe)

Postfach 3640

76021 Karlsruhe, Germany

hermann.dertinger@kit.edu

Conflict of interest statement

The authors declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Risk Associated With Radon Is Overestimated

The main limitation of the article is the fact that radioactivity in tobacco smoke was not even mentioned. According to the US National Council on Radiation Protection and Measurement (NCRP report No. 95), such activity accounts for a radiation exposure of smokers' lungs of more than 100 mSv. This means that even non-smokers are exposed to notable radiation in buildings where smoking is permitted. The substantial emission of plutonium by coal fired power stations, surely another possible factor of influence in radon studies (and an important task in occupational medicine) was not mentioned at all.

In the cited meta-analysis by Darby, the risk of lung malignancies in non-smokers aged 75 was reported to be 0.4%, 0.5%, and 0.7% for radon concentrations of 0 Bq/m³, 100 Bq/m³, and 400 Bq/m³; in smokers, the corresponding risk was about 25 times as high, at 10%, 12%, and 16%. For non-smokers, the difference is only 0.1% for exposures of 0–100 Bq/m³. The authors reported an average radon exposure in Germany of only 49 Bq/m³ (the natural values of outdoor air differ by up to 80 Bq/m³, according to the Federal Office for Radiation Protection (Bundesamt für Strahlenschutz), which means that a relevant influence in non-smokers cannot be shown and the role of radon as a relevant co-carcinogenic substance is questionable even in smokers.

As far as the allegedly linear dose-effect-relation is concerned, I missed a discussion of why in Iran, whose population's natural radiation exposure reaches up to 200 mSv (the equivalent in Germany is 1–6 mSv), no increase in the incidence of malignancies has been found.

The measures stipulated for building construction do not have any relation to the threat posed by radon. The subsidies given to these would be better spent on smoking cessation programmes, although this would be associated with additional financial efforts, lower revenues from tobacco taxes, lacking stimulation of the building construction sector, and a higher burden on the pension and retirement funds—all owing to a relevant increase in life expectancy.

DOI: 10.3238/arztebl.2010.0731a

REFERENCES

1. Schmid K, Kuwert T, Drexler H: Radon in indoor spaces—An underestimated risk factor for lung cancer in environmental medicine. *Dtsch Arztebl Int* 2010; 107(11): 181–6.

PD Dr. med. Ernst Eising

Elper Weg 66

45657 Recklinghausen, Germany

e-g-e@web.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Creating Awareness

I read the article about indoor radon with great interest and noted that the topic was comprehensively dealt with: in addition to health political aspects, construction interventions, and political necessities—such as a radon protection law—were mentioned. About a month after the manuscript had been accepted the World Health Organization published a handbook on the topic (1), in which a reference value of 100 Bq/m³ is recommended, as it is justified from an epidemiological perspective. *Deutsches Ärzteblatt* reported this on 22 September 2009 (2).

The International Radon Project, which started in 2005 and includes more than 100 radon experts and scientists from almost 40 member states, WHO is trying to create awareness for the problems associated with radon. WHO is particularly interested in informing those in positions of political responsibility about the health effects of long term radon exposure, so as to drive forward the implementation of the necessary legal framework conditions at the national level. Additionally, specific professional groups, such as building professionals, need to be addressed, because by means of modified building techniques and interventions they are able to influence the concentrations of radon in residential dwellings. Finally, the medical community needs to be included in these efforts to be able to inform about existing health risks due to radon. For these reasons I think that the article was excellent.

DOI: 10.3238/arztebl.2010.0731b

REFERENCES

1. WHO handbook on indoor radon: a public health perspective: (eds.) Zeeb H, Shannoun F, WHO, Geneva, 2009. www.who.int/ionizing_radiation/env/radon/en/index1.html
2. www.aerzteblatt.de/v4/news/news.asp?id=38209&src=suche&p=strahlenschutz
3. Schmid K, Kuwert T, Drexler H: Radon in indoor spaces—An underestimated risk factor for lung cancer in environmental medicine. *Dtsch Arztebl Int* 2010; 107(11): 181–6.

Dr. P. H. Ferid Shannoun

Department of Public Health and Environment

World Health Organization (WHO)

20 Avenue Appia

CH-1211 Geneva 27

Switzerland

shannounf@who.int

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

In Reply:

The lively debate surrounding our article (1) confirms our assessment of the situation—namely, that the problem of radon in indoor spaces will have to be given closer consideration in the current debate on environmental policy in Germany.

Our article aimed to present the problems of radon pollution in indoor spaces from the perspective of environmental medicine. Radioactive pollution from power stations, radioactive contamination of tobacco leaves, or radioactivity in Iran were not subjects dealt with in our article; they would have exceeded the scope.

In the same way as any medical therapeutic measure, the use of radon in the context of radon balneology requires a careful assessment of risk versus benefits on the basis of strictly scientific criteria. However, this would require separate scientific article.

In response to Dr Schlarb, we wish to point out that the unit curie has been replaced by Becquerel (Bq), with 1 curie corresponding to 37 gigabecquerel. Radon activity concentrations in “curative” waters are usually 1.5–181 MBq/m³ of water. The release of radon from thermal spa water may result in activity concentrations in the immediate surrounding environmental air of up to 100 000 Bq/m³; such concentrations have been measured in radon healing galleries used for radon inhalation therapy (2).

We did mention potential radon content of drinking water and natural gas. The estimated mean values for radon concentrations in water in the west German states is about 400–4000 Bq/m³ (2). If drinking water contains high enough concentrations of radon then relevant exposures may be possible while taking a shower. However, the usual length of stay in a narrow shower cubicle is so limited that the contribution to the overall exposure would be limited too.

The lacking air exchange and ventilation in modern residential buildings has an important role in environmental medicine. Many pollutants (not only radon) may concentrate in indoor spaces as a result. The environmental medical demand is therefore obviously to ensure a satisfactory rate of ventilation, if required by means of technical air conditioning.

With regard to the letter by Dr Eising, we’d like to maintain that the cited epidemiological assessments were conducted in full consideration of smoking

behaviors. A large number of highly potent chemical carcinogenic substances are found in the mainstream smoke and sidestream smoke (3). The radiation pollution due to radioactive isotopes—for example, polonium-210—in tobacco leaves and thus tobacco smoke will thus also contribute to the carcinogenic effect (4, 5).

After careful checking of the scientific literature we conclude that radon in indoor spaces is undoubtedly an independent risk factor for lung cancers. We are happy to support Dr Eising’s demand for smoking cessation programs, since active cigarette smoking is definitely the most important risk factor for the development of lung cancer. However, doing one thing should never mean omitting to do another. We emphatically thank Dr Shannoun for mentioning the current handbook on radon in indoor spaces and the World Health Organization’s international radon project. We agree with him that interventions are needed with regard to radon in indoor spaces.

DOI: 10.3238/arztebl.2010.0732

REFERENCES

- Schmid K, Kuwert T, Drexler H: Radon in indoor spaces—An underestimated risk factor for lung cancer in environmental medicine. *Dtsch Arztebl Int* 2010; 107(11): 181–6.
- Strahlenschutzkommission: Radon und Lungenkrebs. In: Veröffentlichungen der Strahlenschutzkommission Band 62: Einfluss der natürlichen Strahlenexposition auf die Krebsentstehung in Deutschland. Berlin: H. H. Hoffmann GmbH Fachverlag 2008; 116–66.
- Greim H, Deutsche Forschungsgemeinschaft: Senatskommission zur Prüfung gesundheitsschädlicher Arbeitsstoffe (ed.): Passivrauchen am Arbeitsplatz; Änderung krebserzeugender Arbeitsstoffe. Weinheim: WILEY-VCH 1999; 1–36.
- Winters TH, DiFrenza J: Radioactivity and lung cancer in active and passive smokers. *Chest* 1983; 84: 653–4.
- Muggli ME, Ebbert JO, Robertson C, Hurt RD: Waking a sleeping giant: the tobacco industry’s response to the polonium-210 issue. *Am J Public Health* 2008; 98: 1643–50.

PD Dr. med. Klaus Schmid

Prof. Dr. med. Hans Drexler

Institut für Arbeits-, Sozial- und Umweltmedizin
der Universität Erlangen-Nürnberg
Schillerstr. 25
91054 Erlangen, Germany
klaus.schmid@rzmail.uni-erlangen.de

Prof. Dr. med. Torsten Kuwert

Nuklearmedizinische Klinik der Universität Erlangen-Nürnberg
Krankenhausstr. 12
91054 Erlangen, Germany

Conflict of interest statement

Professor Kuwert has received honoraria for speaking from Siemens Medical Solutions. Professor PD Dr Schmid and Professor Drexler declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.