Published in final edited form as:

J Acquir Immune Defic Syndr. 2010 December 1; 55(4): 516–523. doi:10.1097/QAI.0b013e3181f275e0.

Persistence of racial differences in attitudes toward homosexuality in the United States

Sara Nelson Glick, MPH and Matthew R. Golden, MD, MPH

Departments of Epidemiology (Ms. Glick) and Medicine (Dr. Golden), University of Washington, Seattle, WA; HIV/STD Program, Public Health – Seattle & King County (Dr. Golden)

Abstract

Background—Stigma may mediate some of the observed disparity in HIV infection rates between African American and white men who have sex with men (MSM).

Methods—We used data from the General Social Survey to describe race-specific trends in the U.S. population's attitude toward homosexuality, reporting of male same-sex sexual behavior, and behaviors that might mediate the relationship between stigma and HIV transmission among MSM.

Results—The proportion of African Americans who indicated that homosexuality was “always wrong” was 72.3% in 2008, largely unchanged since the 1970s. In contrast, among white respondents, this figure declined from 70.8% in 1973 to 51.6% in 2008, with most change occurring since the early 1990s. Participants who knew a gay person were less likely to have negative attitudes toward homosexuality (RR=0.60, 95% CI: 0.52–0.69). Among MSM, twice as many African American MSM reported that homosexuality is “always wrong” compared to white MSM (57.1% vs. 26.8%, $p=0.003$). MSM with unfavorable attitudes toward homosexuality were less likely to report ever testing for HIV compared to MSM with more favorable attitudes (RR=0.50, 95% CI: 0.31–0.78).

Conclusions—U.S. attitudes toward homosexuality are characterized by persistent racial differences, which may help explain disparities in HIV infection rates between African American and white MSM.

Keywords

stigma; homosexuality; men who have sex with men; race; General Social Survey

Introduction

Despite the fact that men who have sex with men (MSM) comprise <5% of the male U.S. population, 57% of all new HIV diagnoses in the U.S. occurred in MSM in 2007^{1–3}. Among MSM, HIV disproportionately affects African Americans. African Americans comprise approximately 13% of the total U.S. population and nearly an identical proportion of MSM in the General Social Survey (GSS). However, in 2007, 24.5% of reported HIV cases among

Corresponding Author: Sara Nelson Glick, MPH, University of Washington, Center for AIDS and STD, Harborview Medical Center, 325 9th Avenue, Box 359931, Seattle, Washington 98104. Phone: 206-744-8055, Fax: 206-744-3693, sjnelson@u.washington.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Previous Presentations:

Portions of these data were presented at the 17th *Conference on Retroviruses and Opportunistic Infections* held February 16–19, 2010, in San Francisco, CA.

MSM occurred in African Americans, and some CDC surveillance data suggest that HIV prevalence in African American MSM may be more than twice that observed in white MSM⁴. The reasons for this disparity are unclear. African American MSM do not clearly have higher levels of sexual risk behavior than white MSM^{5, 6}. However, African American MSM tend to preferentially choose other African American partners^{7, 8} and, compared to HIV-infected white MSM, HIV-infected African American MSM are diagnosed later in the course of infection and less frequently initiate antiretroviral therapy^{4, 9, 10}. Because knowing one's HIV status leads to substantial changes in sexual risk behavior and antiretroviral treatment probably decreases the risk of transmitting HIV, these factors may partially explain the relatively high rates of HIV transmission among African American MSM.

The underlying social factors that may promote the observed racial disparities among MSM remain ill-defined, but stigma toward same-sex behavior has been proposed as one explanation^{11–14}. This stigma manifests as sexual prejudice¹⁵ from the larger heterosexual population as well as a person's own incorporation of society's negative views toward homosexuality (internalized homophobia^{16, 17}). Previous population-based research showed declines in negative opinions about homosexuality in the 1990s¹⁸. Studies also found that African Americans tended to have less favorable attitudes toward homosexuality than white respondents, and that the gap between African American and white populations diverged in the 1990s^{18, 19}. Several of these studies used the GSS, a repeated cross-sectional survey of U.S. households that has included a question about attitudes toward homosexuality since 1973.

We analyzed the GSS using contemporary data (through 2008) to describe trends and correlates of attitudes regarding homosexuality. In addition, the GSS provides one of the only relatively large population-based samples of MSM, thus we also explored trends in the reporting of male homosexual behavior. Because of the increasing racial disparities in HIV among MSM, we were specifically interested in how attitudes might vary by race and in identifying behaviors that might mediate the relationship between stigma and HIV transmission.

Methods

Data

The GSS is a nationally representative, door-to-door survey of the demographics, attitudes, and behaviors of U.S. adults²⁰. Conducted annually or semiannually since 1972, the GSS consists of a household-based probability sample of individuals aged 18 and older. The average response rate²¹ has been 76% (range: 70.0–82.4%) since its inception. Each survey includes permanent items asked of all participants each year, supplemental items asked to all participants for only one year, and rotating items that appear every year but only to a random subset of surveys. The majority of the survey is interviewer-administered, although the rotating sexual behavior module is a self-administered questionnaire. Historically, the self-administered questionnaire used a pencil-and-paper format, but has been computer-administered since 2002.

Measures

In most survey cycles, the GSS has included a question about the respondent's attitude toward homosexuality: "What about sexual relations between two adults of the same sex – do you think it is always wrong, almost always wrong, wrong only sometimes, or not wrong at all?" Unless otherwise indicated, for this analysis we coded the variable as "always wrong" vs. not "always wrong"¹⁹. (Combining "almost always wrong" with "always wrong" did not appreciably change the study findings.)

Until 2000, a respondent's race was coded as white, African American, or other based on the interviewer's observation. If the interviewer was unsure, respondents identified their own race.

Subsequent surveys asked all respondents to self-identify their race and included a separate question about Hispanic ethnicity.

Since 1989, a subsample of GSS respondents answered two questions that identify MSM, including the gender of sex partners in the last 12 months and the number of male sex partners since age 18. Starting in 1991, the GSS included a question about the gender of sex partners in the last 5 years. (The GSS does not distinguish between oral and anal sex partners.) Unless otherwise noted, we defined MSM using the 5-year variable because it is likely a more specific measure of male homosexuality than any male partner since age 18. The 5-year measure is also more sensitive and provides more analytical power than the measure that captures MSM behavior in the past year. Nevertheless, the three measures were relatively interchangeable and provided comparable results.

The GSS asked a subset of participants in 2006 about the number of the respondent's acquaintances in various settings who are "gay men or women" – e.g., general acquaintances, relatives, neighbors, co-workers, or members of volunteer associations. We coded any response >0 as being indicative of knowing someone who is gay. To explore the association between sexual risk behavior and attitudes toward homosexuality among MSM, we used a question from the 2006 and 2008 surveys that asked if the respondent had ever been tested for HIV.

Finally, to assess the independent association between race and attitudes toward homosexuality, we conducted multivariate analyses adjusted for year of survey and demographic variables that have been consistent across all GSS surveys: gender, age, marital status, education, religion, family income (adjusted to 2000 dollars), region of the U.S., and size of community.

Analysis

We could not identify Hispanic ethnicity prior to 2000, thus our analyses only included results for respondents who are coded as white or African American, regardless of ethnicity. To explore the overall prevalence and temporal trends in attitudes toward homosexuality, we calculated overall, year-specific, and race-specific proportions and 95% confidence intervals (CI). We also conducted univariate and multivariate analyses to identify independent correlates of unfavorable attitudes. Given the changes in attitudes over time, we included an indicator variable for each survey year in the multivariate model. Because the outcome is common, we used Poisson regression with a log link to produce unbiased estimates of the risk ratios. We explored other correlates of attitudes toward homosexuality using chi-square tests.

We hypothesized that the temporal trend toward greater acceptance of homosexuality might be most pronounced among the youngest persons, reflecting different initial perspectives formed during adolescence, and that recent birth cohorts would reflect more marked changes in attitude with aging. We conducted exploratory analyses to test for interactions between age and year of interview, and between birth cohort and year of interview, on attitudes toward homosexuality. We used likelihood ratio tests to assess whether the inclusion of each set of interaction terms was statistically significant in the multivariate Poisson model.

We also estimated the prevalence of same-sex sexual behavior among male respondents and explored correlates of MSM behavior using univariate and multivariate logistic regression and chi-square tests.

Although the GSS utilizes a full probability sample of U.S. households, we used weights to account for sub-sampling of non-respondents, the oversampling of African Americans, and household size. All analyses were conducted using Intercooled Stata 9.2 (College Station, TX).

Results

Overall prevalence and correlates of unfavorable attitudes toward homosexuality

Since 1972, there have been a total of 27 GSS cycles and 53,043 respondents. Consistent with demographic changes in the U.S., over time GSS participants were increasingly non-white, never married, college educated, non-religious, from the South and Mountain West, and from suburban or exurban areas. (See Supplemental Table.) We excluded 2,406 (4.5%) participants who reported that their race was not white or African American. The survey has included a question about attitudes toward homosexuality to a subset of participants in all but 5 of the surveys. Thus, the final dataset for this analysis included 30,837 respondents. Combining all surveys, 68.0% (95% CI: 67.5–68.6%) of participants reported that homosexuality was “always wrong.”

On multivariate analysis (Table 1), the belief that homosexuality is “always wrong” was significantly associated with African American race; male gender; older age; being married, divorced, separated, or widowed (vs. never married); lower educational attainment; lower income; living in the South, Midwest or Mountain West (vs. Northeast); and not living in a city. Compared to Protestant affiliation, all other religions were associated with more favorable attitudes toward homosexuality with the strongest associations among Jewish participants and those with no religious affiliation. Further adjustment by Protestant denomination (Baptist, Methodist, Lutheran, Presbyterian, Episcopal, other, none) did not result in a meaningful change in the estimates, although we were not able to further refine these categories (e.g., Evangelical or not). Including strength of religious affiliation resulted in a slight attenuation of the association between African American race and reporting that homosexuality is “always wrong” (RR=1.11, 95% CI: 1.08–1.14). There was no significant change in the proportion of respondents who were African American between 1998 and 2000, the year that the GSS began asking participants to identify their own race. However, the adjusted association between African American race and unfavorable attitudes toward homosexuality was stronger in 2000–2008 (RR=1.28, 95% CI: 1.21–1.36) than before 2000 (RR=1.11, 95% CI: 1.09–1.14) (p-value for interaction<0.001).

Using data available only in 2006, we found that knowing someone who is gay was strongly associated with more accepting attitudes toward homosexuality. Overall, 54.9% (95% CI: 51.6–58.2%) of respondents reported knowing someone who is gay. This proportion was somewhat lower among African American compared to white respondents, though this difference was not significant (p=0.09). In univariate analysis, knowing a gay person was significantly less likely among the oldest participants and more likely among women and respondents with more than a high school education, no religious affiliation, or income $\geq$ \$50,000 (Table 2). Respondents who reported knowing a gay person were significantly less likely to indicate that homosexuality was “always wrong” compared to those who did not (RR=0.60, 95% CI: 0.52–0.69). While African Americans who reported knowing someone who is gay were less likely to report unfavorable attitudes toward homosexuality (RR=0.85, 95% CI: 0.69–1.04), this effect was stronger among white participants (RR=0.56, 95% CI: 0.47–0.66) (p-value for interaction=0.002).

Temporal trends in unfavorable attitudes toward homosexuality

The question about attitudes toward homosexuality was first asked in 1973, at which time 72.5% (95% CI: 69.9–74.9%) of respondents indicated that homosexuality was “always wrong”. This proportion increased through the 1970s and 1980s, peaked in 1988 at 77.8% (95% CI: 74.7–80.5%), and declined rapidly to 61.0% (95% CI: 58.5–63.5%) in 1996. The estimate continued to decline slightly into the early 2000s and has remained relatively stable through 2008 at 54.7% (95% CI: 51.9–57.4%).

To assess if age at interview and birth cohort modified the observed calendar year trends, we constructed two multivariate models that included all demographic variables listed in Table 1, as well as the interaction between age at interview and 5-year survey intervals and the interaction between 10-year birth cohort and 5-year survey intervals. The interaction between age and calendar year was not significant ($p=0.131$), suggesting that the association between more accepting attitudes to homosexuality associated with youth were stable over time, though the differences between age groups were less pronounced in the 1980s when the negative attitudes among all age groups increased. The interaction between birth cohort and calendar year was also non-significant ($p=0.706$). There were slight decreases in negative attitudes toward homosexuality in all birth cohorts between the 1980s and 2000s, but there were no meaningful differences in the magnitude of those changes between cohorts over time.

Non-uniformity in temporal trends in attitudes toward homosexuality

In contrast to the experience of white respondents, the proportion of African Americans who indicated that homosexuality was “always wrong” decreased relatively little between the early 1970s and 2008 (Figure 1). For example, among African Americans, 84.2% (95% CI: 77.8–89.0%) reported that homosexuality is always wrong in 1973 and 72.3% (95% CI: 65.2–78.5%) reported this in 2008, a 14% decline. In contrast, among white respondents, the proportion that disapproved of homosexuality declined from 70.8% (95% CI: 68.0–73.4%) to 51.7% (95% CI: 48.7–54.7%) between 1973 and 2008, a 27% drop. The observed racial differences in attitudes toward homosexuality diverged dramatically in the 1990s. In 1990, the gap between African American and white respondents’ attitudes was 6.6 percentage points. It increased to 25.3 in 2004, but has closed somewhat in the two most recent GSS cycles.

For comparison, we also plotted the response that homosexuality is “not wrong at all” for each year (Figure 1). The trends roughly correspond to the inverse of the response that homosexuality is “always wrong”. After nearly two decades of little difference in the proportion (10–20%) of white and African American respondents reporting that homosexuality is “not wrong at all,” in 2008 this estimate increased only slightly among African Americans to 21.0% (95% CI: 15.6–27.7%) and more dramatically among white respondents to 37.6% (95% CI: 34.8–40.6%).

Prevalence of male homosexual behavior and attitudes toward homosexuality among MSM

Between 1989 and 2008, 5.4% (95% CI: 4.9–5.9%) of male respondents reported a male sex partner since the age of 18, 3.9% (95% CI: 3.4–4.4%) in the past 5 years, and 3.2% (95% CI: 2.8–3.6%) in the past year. Since 2000, there has been an increase in the proportion of men reporting same-sex behavior since age 18 but a slight decline in the proportion reporting this in the past 1 or 5 years (Figure 2). There was no significant change in the proportion of men reporting MSM behavior between 2000 and 2002, which was the year that the GSS implemented a computer-based survey for sexual behavior questions. African American men were significantly more likely to have a male sex partner in the past year (4.4%) compared to white men (3.1%) ($p=0.037$), but there was no significant difference between racial groups in the probability of reporting a male partner in the past 5 years (4.9% vs. 3.7%, $p=0.114$) or since age 18 (6.6% vs. 5.2%, $p=0.088$). In univariate analyses of men surveyed since 1989, no religious affiliation was associated with increased likelihood of being a MSM (in the past 5 years); while income $\geq \$50,000$, living in the southern U.S., or living outside of a city was associated with not reporting MSM behavior (Table 3). After adjusting for all demographic variables and year of interview, only age 35–54 (vs. 18–24), lower income, and living in a city were associated with being a MSM in the past 5 years.

Among men interviewed between 1991 and 2008, nearly one-third (30.5%, 95% CI: 24.0–38.0%) of MSM and 63.5% (95% CI: 62.1–65.0%) of non-MSM indicated that they believe

that homosexuality is “always wrong.” Among MSM, attitudes toward homosexuality differed significantly by race with nearly twice as many African American MSM reporting that homosexuality is “always wrong” compared to white MSM (57.1% vs. 26.8%, $p=0.003$).

MSM who believed that homosexuality is “always wrong” were less likely to report ever testing for HIV (36.0%) compared to MSM who did not believe this (72.6%) (RR=0.50, 95% CI: 0.31–0.78), and this association remained unchanged after adjustment for income and education. (For this analysis only, we defined MSM as same-sex behavior since age 18 due to small cell sizes with the other measures. The other MSM measures showed a similar association.) Our analysis lacked sufficient power to look at this association by race.

Discussion

The proportion of U.S. adults with more accepting attitudes toward homosexuality grew substantially over the last 35 years. After an increase in negative attitudes during the 1980s, the percentage of white and African American GSS respondents who indicated that homosexuality is “always wrong” dropped by approximately 25 points. Nevertheless, the majority (55%) of the population continues to disapprove of homosexuality. Moreover, a profound gap in attitudes between white and African American adults developed in the 1990s^{18, 19}, and our findings demonstrate that this difference, though no longer clearly growing, is now large and stable. We also observed that internalized homophobia was more common among African American than white MSM, and that MSM who reported that homosexuality was wrong were less likely to test for HIV. Furthermore, having a gay acquaintance was strongly associated with more accepting attitudes towards homosexuality, though the magnitude of that association was significantly smaller among African Americans than among whites.

Our findings, along with prior research, suggest a possible, at least partial, explanation for the observed disparity in HIV between African American and white MSM. Negative attitudes toward homosexuality at the population level may inhibit MSM from coming out. Because greater acceptance of homosexuality may be fostered by social interactions with gay individuals^{22–24}, widespread disapproval of homosexuality – like that observed among African Americans – is self-perpetuating. Insofar as this disapproval also promotes internalized homophobia among MSM themselves, it inhibits men from adopting behaviors, like HIV testing, that are likely to decrease HIV transmission at the population level^{12, 25–29}. The fact that MSM, like heterosexuals, preferentially choose sex partners of the same race^{7, 8}, further magnifies the population-level impact of what might otherwise be relatively small differences in behavior. Nevertheless, given the cross-sectional nature of each GSS survey, we cannot say that the associations we observed between knowing someone who is gay and less disapproval of homosexuality, and between internalized homophobia and decreased HIV testing, are causal. While we observed an association between knowing someone who is gay and more approving attitudes toward homosexuality, this association was significantly weaker among African Americans, highlighting that the increased visibility of sexual minorities alone is not likely to be sufficient to ameliorate this stigma. The GSS also did not begin collecting data on coming out until 2008, so there were too few data to explore the potential impact of attitudes toward homosexuality on disclosure of sexual orientation. Finally, we cannot rule out that the association between stigma and HIV testing may be confounded by access to health care, although this association was not attenuated after adjustment for income and education. (The GSS does not routinely ask about health insurance coverage.).

We did not observe an ongoing trend toward greater levels of homosexual behavior in the male population. Using earlier GSS data, Anderson and Stall reported that MSM behavior in the past year increased between 1996 and 2000². Population data from the United Kingdom also

demonstrated an increase in male same-sex behavior between 1990 and 2000¹. We found that the proportion of men reporting same-sex sexual behavior since age 18 increased steadily over the past 10 years, while the proportion reporting this in the past 1 and 5 years did not display this trend. Like other self-reported behaviors, it is difficult to know if these trends represent a true reflection of behavior in the population or are a function of more forthcoming reporting over time. This difficulty is exacerbated by the fact that the GSS changed from using a self-administered to computer-administered format in 2002, perhaps increasing reporting of stigmatized behaviors. The absence of a clear and sustained trend in MSM behavior in the prior 1 or 5 years in our analysis emphasizes the uncertainty of whether there is true trend toward greater homosexual behavior in the population, particularly given the very small number of MSM that participate in any individual wave of GSS or other population-based studies.

The primary strength of this study is the use of a frequent, population-based survey. The large annual sample size allowed us to explore trends within subgroups over 35 years. Nevertheless, our study has several limitations. First, our measure of attitudes toward homosexuality is based on a single question as opposed to a more comprehensive measure^{30–32}, and many researchers have advocated for implicit, rather than explicit, measures for socially sensitive attitudes³³. Our analysis also does not account for the finding that African Americans are more likely than whites to report supporting laws prohibiting antigay discrimination¹⁹. Furthermore, the negative framing of the question may have led to an overestimate of negative attitudes toward homosexuality in the population. However, since the wording has remained constant over time, the temporal trend patterns should be valid. Second, not all questions were asked in all surveys resulting in small cell sizes in some analyses. Most notably, the HIV testing question was only recently added to the GSS and there were too few observations to stratify these findings by race among MSM, thus additional data are needed. Third, many respondents reported not knowing their specific religious denomination, thus we were not able to adjust for liberal and conservative denominations. Fourth, the categorization of race was crude prior to 2000. Undoubtedly, there was misclassification due to the interviewer's presumption about the respondent's race. Finally, the primary aim of the GSS is to provide data on changing social attitudes, thus the survey includes only a limited number of sexual behavior questions. Therefore, more definitive studies of attitudes toward homosexuality on HIV risk among MSM may need to employ studies with larger numbers of MSM (e.g., National HIV Behavioral Surveillance).

We acknowledge that this study cannot address with certainty if or how attitudes toward homosexuality in the general population affect HIV risk among MSM in general, or African American MSM in particular. However, our analysis provides a contemporary description of these attitudes in the U.S. population and suggests at least one plausible causal pathway for how attitudes toward homosexuality might result in the dramatic racial disparities in HIV rates observed in U.S. MSM. Given the tragic persistence of a profound HIV epidemic in MSM and the paucity of effective interventions to control that epidemic, we believe that attempting to alleviate the stigma associated with homosexuality should be considered as HIV prevention interventions. Our finding that acquaintanceship with someone who is gay is associated with decreased homophobia suggests that increasing the visibility of MSM may be an effective way to diminish homophobia, though this requires additional study. Beyond this, our observations suggest a need for ongoing surveillance related to homophobia, better research on how sexual prejudice influences sexual risk and health, intervention studies to define if and how public health efforts can diminish homophobia at the population-level, and what impact such efforts might have on the health of MSM.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Sources of Support:

Ms. Glick is supported by the University of Washington STD/AIDS Research Training Program (T32 AI007140) from the National Institutes of Health, U.S. Public Health Service.

Disclosure of Funding:

This research was supported by a pre-doctoral training grant funded by the National Institutes of Health.

References

1. Mercer CH, Fenton KA, Copas AJ, et al. Increasing prevalence of male homosexual partnerships and practices in Britain 1990–2000: evidence from national probability surveys. *Aids* 2004 Jul 2;18(10):1453–1458. [PubMed: 15199322]
2. Anderson JE, Stall R. Increased reporting of male-to-male sexual activity in a national survey. *Sex Transm Dis* 2002 Nov;29(11):643–646. [PubMed: 12438899]
3. CDC. Cases of HIV Infection and AIDS in the United States and Dependent Areas 2007. Atlanta, Georgia: 2007.
4. CDC. HIV prevalence, unrecognized infection, and HIV testing among men who have sex with men--five U.S. cities, June 2004–April 2005. *MMWR Morb Mortal Wkly Rep* 2005 Jun 24;54(24):597–601. [PubMed: 15973239]
5. Millett GA, Flores SA, Peterson JL, Bakeman R. Explaining disparities in HIV infection among black and white men who have sex with men: a meta-analysis of HIV risk behaviors. *Aids* 2007 Oct 1;21(15):2083–2091. [PubMed: 17885299]
6. Millett GA, Peterson JL, Wolitski RJ, Stall R. Greater risk for HIV infection of black men who have sex with men: a critical literature review. *Am J Public Health* 2006 Jun;96(6):1007–1019. [PubMed: 16670223]
7. Mimiaga MJ, Reisner SL, Cranston K, et al. Sexual mixing patterns and partner characteristics of black MSM in Massachusetts at increased risk for HIV infection and transmission. *J Urban Health* 2009 Jul;86(4):602–623. [PubMed: 19466554]
8. Berry M, Raymond HF, McFarland W. Same race and older partner selection may explain higher HIV prevalence among black men who have sex with men. *Aids* 2007 Nov 12;21(17):2349–2350. [PubMed: 18090287]
9. Palacio H, Kahn JG, Richards TA, Morin SF. Effect of race and/or ethnicity in use of antiretrovirals and prophylaxis for opportunistic infection: a review of the literature. *Public Health Rep* 2002 May-Jun;117(3):233–251. discussion 231–232. [PubMed: 12432135]
10. Hall HI, Byers RH, Ling Q, Espinoza L. Racial/ethnic and age disparities in HIV prevalence and disease progression among men who have sex with men in the United States. *Am J Public Health* 2007 Jun;97(6):1060–1066. [PubMed: 17463370]
11. Peterson JL, Jones KT. HIV prevention for black men who have sex with men in the United States. *Am J Public Health* 2009 Jun;99(6):976–980. [PubMed: 19372510]
12. Huebner DM, Davis MC, Nemeroff CJ, Aiken LS. The impact of internalized homophobia on HIV preventive interventions. *Am J Community Psychol* 2002 Jun;30(3):327–348. [PubMed: 12054033]
13. Meyer, IH.; Dean, L. Internalized homophobia, intimacy, and sexual behavior among gay and bisexual men. In: Herek, GM., editor. *Stigma and sexual orientation: Understanding prejudice against lesbians, gay men and bisexuals*. Thousand Oaks, CA: Sage; 1998.
14. Stokes JP, Peterson JL. Homophobia, self-esteem, and risk for HIV among African American men who have sex with men. *AIDS Educ Prev* 1998 Jun;10(3):278–292. [PubMed: 9642425]
15. Herek GM. The psychology of sexual prejudice. *Curr Dir Psychol Sci* 2000;9(1):19–22.
16. Smolenski DJ, Ross MW, Risser JM, Rosser BR. Sexual compulsivity and high-risk sex among Latino men: the role of internalized homonegativity and gay organizations. *AIDS Care* 2009 Jan;21(1):42–49. [PubMed: 19085219]

17. Ross MW, Rosser BR. Measurement and correlates of internalized homophobia: a factor analytic study. *J Clin Psychol* 1996 Jan;52(1):15–21. [PubMed: 8682906]
18. Loftus J. America's liberalization in attitudes toward homosexuality, 1973 to 1998. *American Sociological Review* 2001 Oct;66(5):762–782.
19. Lewis GB. Black-white differences in attitudes toward homosexuality and gay rights. *Public Opinion Quarterly* 2003 Spr;67(1):59–78.
20. Davis JA, Smith TW. General social surveys, 1972–2008 [machine-readable data file]; Principal Investigator, James A. Davis; Director and Co-Principal Investigator, Tom W. Smith; Co-Principal Investigator, Peter V. Marsden; Sponsored by National Science Foundation. --NORC ed.-- Chicago: National Opinion Research Center [producer]; Storrs, CT: The Roper Center for Public Opinion Research, University of Connecticut [distributor]; 2009
21. American Association for Public Opinion Research. Standard Definitions: Final Disposition of Case Codes and Outcome Rates for Surveys. 2006
22. Herek GM, Capitanio JP. "Some of My Best Friends": Intergroup Contact, Concealable Stigma, and Heterosexuals' Attitudes Toward Gay Men and Lesbians. *Personality and Social Psychology Bulletin* 1996;22(4):412–424.
23. Herek GM, Glunt EK. Interpersonal Contact and Heterosexuals Attitudes toward Gay Men - Results from a National Survey. *Journal of Sex Research* 1993 Aug;30(3):239–244.
24. Heinze JE, Horn SS. Intergroup contact and beliefs about homosexuality in adolescence. *J Youth Adolesc* 2009 Aug;38(7):937–951. [PubMed: 19636737]
25. Williamson IR. Internalized homophobia and health issues affecting lesbians and gay men. *Health Educ Res* 2000 Feb;15(1):97–107. [PubMed: 10788206]
26. Newcomb ME, Mustanski B. Moderators of the Relationship Between Internalized Homophobia and Risky Sexual Behavior in Men Who Have Sex with Men: A Meta-Analysis. *Arch Sex Behav*. 2009 Nov 4;
27. Shoptaw S, Weiss RE, Munjas B, et al. Homonegativity, substance use, sexual risk behaviors, and HIV status in poor and ethnic men who have sex with men in Los Angeles. *J Urban Health* 2009;86 Suppl 1:77–92. [PubMed: 19526346]
28. Cassels S, Menza TW, Goodreau SM, Golden MR. HIV serosorting as a harm reduction strategy: evidence from Seattle, Washington. *Aids* 2009 Nov 27;23(18):2497–2506. [PubMed: 19834319]
29. Wilson DP, Hoare A, Regan DG, Law MG. Importance of promoting HIV testing for preventing secondary transmissions: modelling the Australian HIV epidemic among men who have sex with men. *Sex Health* 2009 Mar;6(1):19–33. [PubMed: 19254488]
30. Stoever CJ, Morera OF. A confirmatory factor analysis of the Attitudes Toward Lesbians and Gay Men (ATLG) measure. *J Homosex* 2007;52(3–4):189–209. [PubMed: 17594977]
31. Herek GM. Heterosexuals Attitudes toward Lesbians and Gay Men - Correlates and Gender Differences. *Journal of Sex Research* 1988 Nov;25(4):451–477.
32. Mayfield W. The development of an Internalized Homonegativity Inventory for gay men. *J Homosex* 2001;41(2):53–76. [PubMed: 11482428]
33. Greenwald AG, Poehlman TA, Uhlmann EL, Banaji MR. Understanding and using the Implicit Association Test: III. Meta-analysis of predictive validity. *J Pers Soc Psychol* 2009 Jul;97(1):17–41. [PubMed: 19586237]

Figure 1.

Proportion of adults who indicated that homosexuality is “always wrong” or “not wrong at all”, General Social Survey, 1973–2008. **Key:** African American respondents: solid lines; white respondents: dashed lines.

Figure 2.

Proportion of male adults reporting same-sex behavior; General Social Survey, 1989–2008 data only. **Key:** MSM since age 18: solid line; MSM in past 5 years: long dashed lines; MSM in past 1 year: short dashed lines.

Table 1

Univariate and multivariate correlates of adults reporting that homosexuality is “always wrong,” General Social Survey, 1973–2008 (n=30,837).

Variable	Unweighted Prevalence [% (95% CI)]	Weighted Prevalence [% (95% CI)]	Univariate RR [RR (95% CI)]	Multivariate RR* [RR (95% CI)]
Race				
White	65.8 (65.2–66.4)	66.5 (65.9–67.2)	1.00	1.00
African American	78.7 (77.5–79.9)	78.7 (77.3–80.0)	1.18 (1.16–1.21)	1.15 (1.12–1.18)
Gender				
Female	67.0 (66.3–67.7)	66.7 (65.9–67.5)	1.00	1.00
Male	68.5 (67.8–69.3)	69.6 (68.8–70.4)	1.04 (1.03–1.06)	1.10 (1.09–1.12)
Age				
18–24	58.3 (56.6–60.0)	59.0 (57.2–60.9)	1.00	1.00
25–34	60.1 (58.9–61.2)	60.5 (59.2–61.8)	1.02 (0.99–1.06)	1.00 (0.96–1.04)
35–44	63.8 (62.6–65.0)	64.9 (63.6–66.1)	1.10 (1.06–1.14)	1.05 (1.01–1.10)
45–54	66.2 (64.9–67.5)	68.1 (66.7–69.5)	1.15 (1.11–1.20)	1.07 (1.03–1.12)
55–64	74.1 (72.7–75.4)	75.5 (74.0–76.9)	1.28 (1.23–1.33)	1.12 (1.08–1.17)
65+	83.3 (82.3–84.2)	83.5 (82.4–84.6)	1.41 (1.37–1.46)	1.19 (1.15–1.24)
Marital Status				
Never married	52.3 (51.1–53.5)	53.9 (52.5–55.4)	1.00	1.00
Separated/Divorced	62.7 (61.4–64.1)	62.5 (60.9–64.0)	1.16 (1.12–1.20)	1.07 (1.03–1.11)
Married	72.0 (71.3–72.7)	72.1 (71.3–72.8)	1.34 (1.30–1.38)	1.21 (1.17–1.25)
Widowed	82.2 (80.9–83.5)	81.6 (80.0–83.1)	1.51 (1.46–1.56)	1.18 (1.13–1.23)
Education				
Less than high school	83.6 (82.8–84.4)	83.1 (82.2–84.1)	1.00	1.00
High school diploma	69.0 (68.3–69.7)	69.4 (68.7–70.2)	0.84 (0.82–0.85)	0.93 (0.91–0.95)
College degree (4 year)	45.8 (44.6–47.0)	46.8 (45.5–48.1)	0.56 (0.55–0.58)	0.67 (0.65–0.70)
Religion				
Protestant	76.2 (75.6–76.8)	76.7 (76.0–77.3)	1.00	1.00
Catholic	63.0 (61.8–64.1)	63.2 (61.9–64.4)	0.82 (0.81–0.84)	0.91 (0.89–0.93)
Jewish	28.7 (25.1–32.3)	28.0 (24.3–32.1)	0.37 (0.32–0.42)	0.43 (0.37–0.51)
None	35.7 (34.0–37.5)	36.4 (34.5–38.4)	0.48 (0.45–0.50)	0.56 (0.53–0.59)
Other	53.9 (50.5–57.3)	56.3 (52.5–59.9)	0.73 (0.69–0.78)	0.87 (0.81–0.93)
Family Income (adjusted to 2000 dollars)				
<\$25,000	73.3 (72.4–74.2)	73.4 (72.4–74.3)	1.00	1.00
\$25,000–49,999	68.2 (67.3–69.2)	70.1 (69.1–71.1)	0.96 (0.94–0.97)	1.01 (0.99–1.03)
\$50,000+	58.5 (57.5–59.5)	60.2 (59.1–61.3)	0.82 (0.80–0.84)	0.94 (0.92–0.97)
Region of US				
Northeast	57.9 (56.6–59.1)	57.9 (56.5–59.3)	1.00	1.00
Midwest	69.2 (68.2–70.1)	69.5 (68.4–70.6)	1.20 (1.17–1.23)	1.12 (1.09–1.16)
South	77.2 (76.4–77.9)	77.6 (76.8–78.5)	1.34 (1.31–1.38)	1.21 (1.18–1.25)
Mountain West	62.8 (60.6–65.1)	64.6 (62.2–67.0)	1.12 (1.07–1.17)	1.13 (1.08–1.18)
Pacific	54.9 (53.3–56.4)	55.9 (54.1–57.6)	0.96 (0.93–1.00)	1.02 (0.98–1.06)

Variable	Unweighted Prevalence [% (95% CI)]	Weighted Prevalence [% (95% CI)]	Univariate RR [RR (95% CI)]	Multivariate RR* [RR (95% CI)]
Size of city				
City	61.7 (60.7–62.7)	61.8 (60.8–63.0)	1.00	1.00
Suburb (incorporated)	62.8 (61.8–63.8)	63.2 (62.2–64.3)	1.02 (1.00–1.05)	1.05 (1.03–1.08)
Exurb (unincorporated)	69.7 (68.2–71.1)	70.0 (68.4–71.5)	1.13 (1.10–1.16)	1.08 (1.05–1.12)
Small city/town	74.7 (73.3–76.0)	74.8 (73.3–76.3)	1.21 (1.18–1.24)	1.10 (1.07–1.13)
Rural or open area	81.9 (80.8–83.0)	81.8 (80.5–83.0)	1.32 (1.29–1.35)	1.13 (1.11–1.16)

*
The multivariate model adjusted for survey year and included all variables listed in Table 1.

Table 2

Univariate and multivariate correlates of knowing someone who is gay among adults, General Social Survey, 2006 data only (n=1,178).

Variable	Unweighted Prevalence [% (95% CI)]	Weighted Prevalence [% (95% CI)]	Univariate RR [RR (95% CI)]	Multivariate RR* [RR (95% CI)]
Race				
White	56.8 (53.8–59.9)	56.0 (52.4–59.5)	1.00	1.00
African American	48.0 (40.5–55.5)	48.0 (39.5–56.6)	0.86 (0.71–1.04)	0.92 (0.75–1.14)
Gender				
Female	57.5 (53.8–61.3)	57.8 (53.5–62.0)	1.00	1.00
Male	52.7 (48.3–57.1)	51.2 (46.0–56.3)	0.89 (0.78–1.00)	0.88 (0.78–1.01)
Age				
18–24	64.4 (54.4–74.5)	58.8 (46.4–70.2)	1.00	1.00
25–34	61.4 (54.3–68.5)	57.8 (48.9–66.2)	0.98 (0.76–1.27)	0.99 (0.75–1.32)
35–44	64.0 (57.7–70.3)	64.1 (56.7–70.9)	1.09 (0.86–1.38)	1.09 (0.83–1.43)
45–54	55.3 (49.2–61.5)	56.2 (49.2–63.1)	0.96 (0.75–1.22)	0.95 (0.72–1.24)
55–64	59.0 (51.7–66.3)	56.2 (47.7–64.3)	0.96 (0.74–1.23)	0.97 (0.73–1.30)
65+	37.3 (31.2–43.4)	38.0 (31.3–45.3)	0.65 (0.49–0.85)	0.68 (0.48–0.96)
Marital Status				
Never married	61.0 (55.2–66.8)	58.5 (51.1–65.6)	1.00	1.00
Separated/Divorced	61.7 (55.3–68.2)	63.9 (56.2–71.0)	1.09 (0.92–1.29)	1.08 (0.89–1.32)
Married	52.8 (48.7–56.9)	52.8 (48.3–57.3)	0.90 (0.78–1.05)	0.90 (0.74–1.09)
Widowed	42.7 (33.3–52.1)	40.9 (30.6–52.1)	0.70 (0.52–0.94)	1.02 (0.73–1.42)
Education				
Less than high school	38.1 (30.5–45.7)	37.5 (29.1–46.7)	1.00	1.00
High school diploma	52.5 (48.7–56.3)	52.9 (48.5–57.2)	1.41 (1.10–1.81)	1.24 (0.97–1.57)
College degree (4 year)	69.0 (64.2–73.8)	66.8 (60.6–72.5)	1.78 (1.38–2.29)	1.51 (1.17–1.94)
Religion				
Protestant	53.0 (49.2–56.8)	52.5 (48.1–56.8)	1.00	1.00
Catholic	50.4 (44.1–56.7)	50.3 (43.0–57.5)	0.96 (0.81–1.13)	0.81 (0.67–0.97)
Jewish	75.0 (51.2–98.8)	68.4 (33.1–90.5)	1.30 (0.81–2.10)	0.88 (0.40–1.93)
None	66.3 (59.8–72.9)	65.9 (57.6–73.3)	1.26 (1.09–1.45)	1.06 (0.90–1.24)
Other	68.9 (54.8–83.0)	66.8 (49.4–80.5)	1.27 (0.99–1.64)	1.12 (0.87–1.44)
Family Income (adjusted to 2000 dollars)				
<\$25,000	48.2 (42.9–53.6)	48.9 (42.7–55.0)	1.00	1.00
\$25,000–49,999	58.2 (52.7–63.7)	55.0 (48.6–61.2)	1.13 (0.95–1.34)	1.02 (0.86–1.21)
\$50,000+	62.8 (57.9–67.6)	60.2 (54.3–65.7)	1.23 (1.05–1.44)	1.07 (0.89–1.28)
Region of US				
Northeast	59.3 (52.0–66.6)	56.6 (47.4–65.3)	1.00	1.00
Midwest	55.8 (50.2–61.5)	54.6 (48.0–60.9)	0.96 (0.79–1.18)	0.98 (0.79–1.21)
South	50.2 (45.5–54.9)	51.5 (46.2–56.8)	0.91 (0.75–1.10)	0.88 (0.71–1.10)
Mountain West	58.6 (49.5–67.7)	54.1 (43.9–64.0)	0.96 (0.75–1.22)	0.86 (0.65–1.13)
Pacific	64.1 (56.1–72.1)	64.3 (54.9–72.7)	1.14 (0.92–1.41)	1.04 (0.93–1.31)

Variable	Unweighted Prevalence [% (95% CI)]	Weighted Prevalence [% (95% CI)]	Univariate RR [RR (95% CI)]	Multivariate RR* [RR (95% CI)]
Size of city				
City	54.9 (49.5–60.3)	53.7 (47.2–60.0)	1.00	1.00
Suburb (incorporated)	61.5 (56.4–66.5)	60.2 (54.1–66.0)	1.12 (0.96–1.31)	1.08 (0.93–1.26)
Exurb (unincorporated)	56.4 (49.5–63.3)	55.8 (47.9–63.5)	1.04 (0.86–1.25)	1.02 (0.84–1.23)
Small city/town	48.7 (39.3–58.0)	48.4 (38.1–58.8)	0.90 (0.70–1.15)	0.90 (0.69–1.18)
Rural or open area	48.0 (40.6–55.5)	48.8 (40.7–57.0)	0.91 (0.74–1.12)	0.96 (0.77–1.20)

*
The multivariate model adjusted for survey year and included all variables listed in Table 2.

Table 3

Univariate and multivariate correlates of reporting same-sex sexual behavior in the past 5 years among male adults, General Social Survey, 1991–2008 data only (n=7,818).

Variable	Unweighted Prevalence [% (95% CI)]	Weighted Prevalence [% (95% CI)]	Univariate OR [OR (95% CI)]	Multivariate OR* [OR (95% CI)]
Race				
White	4.2 (3.7–4.7)	3.7 (3.3–4.3)	1.00	1.00
African American	5.2 (3.8–6.7)	4.9 (3.6–6.6)	1.32 (0.93–1.87)	1.09 (0.74–1.62)
Age				
18–24	3.6 (2.2–4.9)	3.3 (2.2–4.9)	1.00	1.00
25–34	4.8 (3.8–5.8)	4.5 (3.5–5.7)	1.40 (0.85–2.29)	1.70 (0.97–2.96)
35–44	4.8 (3.9–5.8)	4.6 (3.7–5.7)	1.44 (0.89–2.32)	2.11 (1.23–3.63)
45–54	5.0 (3.9–6.1)	4.5 (3.5–5.8)	1.41 (0.86–2.32)	2.29 (1.32–3.98)
55–64	3.5 (2.4–4.7)	2.4 (1.7–3.5)	0.74 (0.43–1.29)	1.25 (0.68–2.31)
65+	2.8 (1.7–3.8)	2.3 (1.5–3.5)	0.70 (0.38–1.27)	1.02 (0.52–1.99)
Education				
Less than high school	3.3 (2.2–4.3)	3.4 (2.4–4.9)	1.00	1.00
High school diploma	4.2 (3.6–4.8)	3.9 (3.3–4.5)	1.13 (0.75–1.69)	1.20 (0.77–1.88)
College degree (4 year)	5.1 (4.3–6.0)	4.2 (3.4–5.1)	1.23 (0.81–1.87)	1.53 (0.94–2.49)
Religion				
Protestant	3.8 (3.2–4.4)	3.4 (2.9–4.0)	1.00	1.00
Catholic	4.0 (3.1–4.9)	3.9 (3.0–5.1)	1.14 (0.82–1.59)	1.03 (0.72–1.48)
Jewish	6.4 (2.7–10.0)	4.6 (2.2–9.2)	1.37 (0.64–2.93)	1.03 (0.49–2.19)
None	6.0 (4.8–7.3)	5.1 (4.0–6.5)	1.53 (1.13–2.09)	1.25 (0.88–1.76)
Other	5.0 (2.7–7.3)	4.5 (2.7–7.2)	1.33 (0.77–2.27)	1.08 (0.62–1.90)
Family Income (adjusted to 2000 dollars)				
<\$25,000	5.8 (4.7–6.8)	5.3 (4.3–6.6)	1.00	1.00
\$25,000–49,999	5.1 (4.2–6.0)	4.4 (3.6–5.4)	0.82 (0.61–1.12)	0.74 (0.54–1.01)
\$50,000+	3.0 (2.4–3.6)	2.8 (2.2–3.5)	0.50 (0.36–0.69)	0.40 (0.28–0.57)
Region of US				
Northeast	5.4 (4.2–6.6)	4.9 (3.8–6.3)	1.00	1.00
Midwest	3.9 (3.1–4.8)	3.7 (2.9–4.7)	0.75 (0.52–1.09)	0.70 (0.46–1.05)
South	3.7 (3.0–4.4)	3.3 (2.6–4.0)	0.66 (0.46–0.94)	0.66 (0.45–0.97)
Mountain	4.0 (2.3–5.6)	3.3 (2.1–5.2)	0.67 (0.39–1.15)	0.58 (0.33–1.02)
Pacific	5.7 (4.3–7.1)	4.7 (3.6–6.2)	0.97 (0.65–1.44)	0.92 (0.60–1.42)
Size of city				
City	7.0 (6.0–8.0)	6.0 (5.1–7.0)	1.00	1.00
Suburb (incorporated)	3.5 (2.9–4.2)	3.2 (2.6–4.0)	0.53 (0.40–0.70)	0.57 (0.41–0.78)
Exurb (unincorporated)	2.6 (1.5–3.6)	2.7 (1.6–4.6)	0.44 (0.25–0.79)	0.58 (0.32–1.04)
Small city/town	3.1 (2.0–4.2)	2.8 (1.9–4.1)	0.45 (0.29–0.70)	0.47 (0.29–0.74)
Rural or open area	2.5 (1.5–3.6)	2.7 (1.7–4.2)	0.43 (0.26–0.72)	0.55 (0.33–0.93)

* The multivariate model adjusted for survey year and included all variables listed in Table 3.