Published in final edited form as:

Org Lett. 2010 October 15; 12(20): 4466–4469. doi:10.1021/ol101482h.

Tricyclic 2'-C-Modified Nucleosides as Potential Anti- HCV Therapeutics

Orrette R. Wauchope¹, Matthew J. Tomney¹, Joseph L. Pepper¹, Brent E. Korba², and Katherine L. Seley-Radtke^{1,*}

¹ Department of Chemistry and Biochemistry, University of Maryland, Baltimore County, 1000 Hilltop Circle, Baltimore, Maryland 21250

² Department of Microbiology and Immunology, Georgetown University Medical Center, 3900 Reservoir Rd., N.W., Washington, DC 20057

Abstract

1

Promising biological activity in a number of therapeutic areas has been reported for both tricyclic nucleosides and 2'-modified nucleosides. In particular, di-substitution at the C-2' position of nucleosides has resulted in significant activity against the Hepatitis C virus (HCV). Combining this with the observation that tricyclic nucleosides developed in our laboratory have been shown to inhibit the RNA-dependent RNA polymerase NS5B led to the design of a series of 2'-modified tricyclic nucleosides. Details of the synthesis, structural characterization and preliminary biological results are reported.

The Hepatitis C virus (HCV) is a blood borne virus that has affected more than 5 million people in the United States and nearly 200 million people worldwide.¹ HCV is a leading cause of long term liver cirrhosis, resulting in liver transplants, liver failure, and in many cases, death. These complications are particularly life-threatening for patients that are coinfectd with the Human Immunodeficiency virus (HIV). HCV consists of several genotypes, with genotype 1 (HCV-GT1) being the most common. The primary treatment for HCV-GT1 currently consists of a combination of pegylated interferon administered with the nucleoside analogue Ribavirin, however the success rate of this treatment is reported to be less than 50%. Although several promising nucleoside analogues are currently in clinical trials, most have failed to make it to market.^{2,3}

The process by which HCV replicates relies on the use of an RNA-dependant RNA-polymerase (RdRp), in particular, NS5B. The NS5B polymerase is known as a “right hand”

kseley@umbc.edu.

Supporting information available: Experimental details, ¹H, ¹³C NMR and HRMS data provided. This material is available free of charge via the Internet at <http://pubs.acs.org>.

polymerase made up of a palm (the active site), fingers and a thumb domain.⁴ To date, NS5B is the only known site in which elongation of new virus occurs, designating it as a prime target for new drug development.

To date a number of 2'-modified nucleosides (Figure 1) have shown potent biological activity against HCV.^{5,6} Moreover, a thiophene-expanded tricyclic purine nucleoside analogue developed in our laboratories has also shown interesting HCV inhibition (Figure 1).⁷

In an effort to further explore the potential for increasing the HCV activity, the tricyclic nucleoside was modified at the C-2' position of the sugar (Figure 2). It was hoped that by combining these two leads, a synergistic increase in inhibition properties against HCV would be realized.

Two synthetic routes were considered in order to realize the target (Figure 2); adding the methyl group to the preconstructed tricyclic nucleoside at the last step or alternatively, adding the methyl group prior to construction of the tricyclic scaffold. Both routes proceed by way of a key intermediate (**2**, Scheme 1), which could be obtained through a 3-step process using the Vorbrüggen⁸ coupling method as previously reported from our laboratory.⁹

The first pathway was quickly abandoned due to the need for additional protection and deprotection steps and very low yields. Turning to the second route, intermediate **2** was bis-protected at the 3'- and 5'-hydroxyls using tetraisopropylidisiloxane chloride (TIPDSCI) in preparation for selective oxidation of the 2'-OH with pyridinium chlorochromate (PCC) (Scheme 2).¹⁰

Intermediate **3** was then treated with trimethylaluminum¹¹ to afford the C-2' methyl substituted TIPS-protected intermediate, which was then immediately deblocked using TBAF to provide **4**. It should be noted that trimethylaluminum was used instead of the more common Grignard and lithium reagents due to reports of higher yields and the stereoselective addition of the methyl to the α -face of the sugar.^{12,13} This approach was desired since the ultimate goal was to replace the hydroxyl group functionality with a fluorine, which is known to proceed through inversion of configuration by the replacement of the "up" OH.⁶

It should be noted that while the use of trimethylaluminum was reported to give stereoselective yields, to the authors' knowledge, no spectroscopic proof for this selective addition has been reported. In an effort to confirm the correct isomer was obtained, a series of 1D-NOE NMR experiments were run. Irradiations of the anomeric proton, 2'-Me, 3'-H, and the 4'-H showed similar correlations thus providing insight as to the stereochemistry of the sugar. Specifically, the anomeric proton was found to have the strongest correlation to the 2'-Me group. This signal was normalized to 1 as a reference for measuring the strength of other correlations. The anomeric proton showed a relative abundance of 100% (normalized), the 4'-H showed a relative abundance of 38%, while the 3'-H showed relatively no correlation to the 2'-Me (less than 1%). These correlations strongly suggest that the Me addition occurred to the α -face of the C-2'-position as expected.

Proceeding with the construction of the tricyclic base, blocking the hydroxyl groups was undertaken next. Employing the benzyl (Bn) group, the protected intermediate was obtained in a 65% yield as shown in Scheme 2. The remaining steps proceeded as previously reported for the guanosine ribose analogue.⁹ Replacement of the C-5 iodo with the formyl group followed by conversion to the oxime, which was then dehydrated to give the cyano modified **5**.

Derivatization of C-4 with thioglycolamide then afforded an intermediate which was then treated with base to give the bicyclic analogue **6**. Closure of the third ring with standard conditions is known to lead to a competitive formation of the desired tricyclic guanosine analogue (**1**) and the undesired tricyclic xanthosine analogue (**8**), a problem also encountered with the formation of guanosine. Previously we reported¹⁴ a mechanistic study that favors the preferential formation of the guanosine analogue when the sugar is a ribose⁹, however this did not prove fruitful with the C-2'- modification and the xanthosine was the major product in all cases.

In examining the reaction more closely, it appeared that the water present in the 30% aqueous hydrogen peroxide solution used was displacing the intermediate sulfenic (SOH) or sulfinic (SO₂H) acid group on **7** (Scheme 2) more rapidly than the ammonia, thereby resulting in Bn-protected xanthosine **8** as the major product. In an effort to overcome this obstacle, use of a non-nucleophilic peroxide was tried, and indeed, proved highly successful. Thus, by substituting tert-butyl hydrogen peroxide in the ring closure reaction, the ratio of the tricyclic guanosine to xanthosine was increased significantly from 1:20 to 1.2:1.15. Deblocking as before provided **1** and **8**, with the desired guanosine **1** being the major product.

Compounds **1** and **8** were then tested for anti-HCV activity (genotypes 1a and 1b) and cytotoxicity in two separate HCV replicon-containing cell lines using an established¹⁶ standardized three day assay. Unfortunately, neither compound showed any significant activity at concentrations up to 10 μ M. It should also be noted that neither compound exhibited cytotoxicity at concentrations of 100 μ M. While initial results against HCV were disappointing, the compounds will be tested against other viruses as soon as sufficient amounts can be obtained.

In summary, two novel tricyclic C-2' modified nucleosides were successfully obtained and screened for potential activity against HCV. Although the compounds did not show activity, possibly due to their poor solubility in the assay system, the new ring closure protocol significantly increased the ratio of the desired guanosine analogue so that scaleup can now be accomplished. Future efforts are focused on pursuit of the analogous 2'-fluoro, 2'-methyl compounds. The results of those studies will be reported elsewhere as they become available.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was generously supported by the National Institutes of Health (R01 GM076345, KSR). The anti-HCV assays were supported by NIAID contract N01-AI-30046 to GUMC (BEK). The authors are thankful for this support.

References

1. Alter MJ. Gastroenterol. 2007; 13:2436.
2. Foster G, Mathurin P. Antiviral Ther. 2008; 13:3.
3. Macor, JE. Annual Reports in Medicinal Chemistry. Macor, JE., editor. Vol. 44. Elsevier; Oxford, UK: 2009. p. 398Chapter 20
4. Qin W, Luo H, Nomura T, Hayashi N, Yamashita T, Murakami S. J Biol Chem. 2002; 277:2132–2137. [PubMed: 11673460]

5. Eldrup AB, Prhavc M, Brooks J, Bhat B, Prakash TP, Song Q, Bera S, Bhat N, Dande P, Cook PD, Bennett F, Carroll S, Ball RG, Bosserman M, Burlein C, Colwell LF, Fay JF, Flores OA, Getty K, LaFemina RL, Leone J, MacCoss M, McMasters DR, Tomassini JE, Langen DV, Wolanski B, Olsen DB. *J Med Chem.* 2004; 47:5284–5297. [PubMed: 15456273]
6. Clark JL, Hollecker L, Mason JC, Stuyver LJ, Tharnish PM, Lostia S, McBrayer TR, Schinazi RF, Watanabe KA, Otto MJ, Furman PA, Stec WJ, Patterson SE, Pankiewicz KW. *J Med Chem.* 2005; 48:5504–5508. [PubMed: 16107149]
7. Seley-Radtke K, Zhang Z, Wauchope O, Zimmermann S, Ivanov A, Korba B. *Nucleic Acids Symp.* 2008; 52:635–636.
8. Niedballa U, Vorbrüggen H. *J Org Chem.* 1974; 39:3654–3660. [PubMed: 4464327]
9. Seley KL, Zhang L, Hagos A, Quirk S. *J Org Chem.* 2002; 67:3365–3373. [PubMed: 12003548]
10. **2-(4,5-Diiodo-imidazol-1-yl)-5,5,7,7-tetraisopropyl-1- β -D-ribofuranose (3):**Compound **2** (10.0 g, 22.1 mmol) previously dried under high vacuum was treated with anhydrous pyridine (200 mL) and the solution stirred for 15 min, at which point 1,3-dichloro-1,1,3,3-tetraisopropylidisiloxane (8.5 mL, 26.5 mmol) was added and the reaction stirred for 14 h. The solvent was removed under reduced pressure and the crude compound purified by column chromatography eluting with 6:1 hexanes:ethyl acetate to provide **3** as a sticky white foam (12.5 g, 17.3 mmol, 81.4%). The intermediate (9.0 g, 13.0 mmol) was stirred with PCC (7.0 g, 32.4 mmol) in anhydrous CH₂Cl₂ (200 mL) over oven 4Å dried molecular sieves. The reaction turned from orange to a dark red-brown color within the first 30 min. The reaction was stirred for 12 hours at rt at which point TLC analysis confirmed the absence of starting material. The crude reaction mixture was filtered over a pad of celite and the solvent removed under reduced pressure. The crude compound was purified by column chromatography eluting with 6:1 hexanes:ethyl acetate to provide **3** as a light yellow foam (6.0 g, 8.7 mmol, 66.9%). ¹H NMR (CDCl₃) δ 0.99 (m, 28H), 3.75 (dd, 1H), 4.00 (dd, 1H), 4.09 (d, 2H), 4.77 (d, 1H), 5.59 (s, 1H), 7.82 (s, 1H). ¹³C NMR (CDCl₃) δ 12.3 (multiple peaks), 17.5 (multiple peaks), 60.3, 72.4, 78.7, 84.4, 92.5, 99.2, 139.8, 204.5. HRMS calculated for C₂₀H₃₄I₂N₂O₅Si₂ [MH]⁺ 693.0175; Found, 693.0175.
11. **2-(4,5-Diiodo-imidazol-1-yl)-5,5,7,7-tetraisopropyl-3-methyl-1- β -D-ribofuranose (4):**Intermediate **3** (4.0 g, 5.8 mmol) was added to anhydrous CH₂Cl₂ (100 mL) and cooled to 0 °C. Trimethylaluminum (3M in hexanes, 6.7 mL, 20.2 mmol) was added and the mixture vigorously stirred at rt for 2 h. The solution was then cooled in an ice bath, quenched with saturated NaHCO₃ solution, extracted with CH₂Cl₂ (3 $\times$ 150 mL), the organic layers combined, dried over MgSO₄ and the solvent removed under reduced pressure. The crude compound was then purified by column chromatography eluting with 6:1 hexanes:EtOAc to afford **4** as a white solid (3.0 g, 4.2 mmol, 73.2%). The intermediate (2.0 g, 2.8 mmol) was dissolved in anhydrous THF (75 mL) and tetrabutylammonium fluoride (1M in THF, 11.3 mL, 11.3 mmol) was added. The mixture was allowed to stir at rt for 5 h before the solvent was removed under reduced pressure. The crude compound was purified by column chromatography eluting with 10:1 EtOAc:MeOH to give **26** as an off white hygroscopic solid (1 g, 2.15 mmol, 76.3%). ¹H NMR (CD₃OD) δ 1.28 (s, 3H), 3.82 (dd, 1H), 3.95 (d, 2H), 4.08 (d, 1H), 5.67 (s, 1H), 8.12 (s, 1H). ¹³C NMR (CD₃OD) δ 17.6, 61.4, 75.4, 79.4, 81.2, 85.3, 93.8, 96.5, 142.4. HRMS calculated for C₉H₁₂I₂N₂O₄ [MH]⁺ 466.8966; Found, 466.8966.
12. Li NS, Piccirilli JA. *J Org Chem.* 2006; 71:4018–4020. [PubMed: 16674089]
13. Li NS, Lu J, Piccirilli JA. *J Org Chem.* 2009; 74:2227–2230. [PubMed: 19173609]
14. Zhang Z, Wauchope O, Seley K. *Tetrahedron.* 2008; 64:10791–10797. [PubMed: 19946353]
15. New procedure for **1-[(5-Aminoimidazo[4',5':4,5]thieno[3,2-d]pyrimidin-3-yl-7-one)]-b-D-ribofuranose 2,3,5-Triol (1)** and **1-[(5-hydroxyimidazo[4',5':4,5]thieno[3,2-d]pyrimidin-3-yl-7-one)]-b-D-ribofuranose (8):**In a steel reaction vessel, compound **6** (2.62 g, 4.38 mmol) was dissolved in 40 mL anhydrous MeOH, NaOH (1.76 g, 44 mmol) added and the mixture stirred at room temperature for 30 min until a clear solution was obtained. CS₂ (1.6 mL, 26.3 mmol) was added, the vessel sealed, and the reaction heated in an oil bath at 145°C for 18 h. The vessel was cooled, and the solvent removed under reduced pressure to give an orange solid to which THF (80 mL) was added. The mixture was cooled to 0°C and 11.4 mL tert-butyl hydroperoxide (5M in decane) was added. This mixture was stirred at 0°C for 4 h until TLC analysis confirmed the disappearance of the starting material. The reaction mixture was then transferred to a steel reaction vessel, cooled to –78°C, and NH₃ bubbled into the solution for 15 min. The reaction vessel was

sealed and heated in oil bath at 120°C for 18 h. The vessel was cooled, the reaction mixture evaporated to dryness under vacuum to give a yellow solid. The crude mixture was purified by column chromatography eluting with hexanes:EtOAc (3:1) to afford the guanosine and xanthosine analogues as colorless oils. The crude compounds were then dried before anhydrous CH₂Cl₂ (15 mL) was added. At this point 12 molar eq. of BF₃•Et₂O and 12 molar eq. of EtSH were added to the solution dropwise. The reaction was allowed to proceed for 48 hours at room temperature before TLC analysis confirmed complete product formation. The solvent was removed under reduced pressure to give a crude solid. The crude compound was purified via column chromatography using a solvent system of 10:1 MeOH:EtOAc. This provided the guanosine analogue **1** and the xanthosine analogue **8** in a 1.2:1 ratio respectively. Each final compound was rinsed 3 times with CH₂Cl₂, rinsed 3 times with EtOAc, and finally 3 times with a 5:1 mixture of EtOAc:MeOH. Guanosine **1**: ¹H NMR (DMSO-d₆) δ 1.44 (s, 3H), 3.65 (dd, 1H), 4.20 (d, 2H), 4.39 (d, 1H), 5.43 (broad m, 3H), 5.72 (s, 1H), 6.51 (s, 2H), 8.35 (s, 1H), 10.98 (broad s, 1H). Xanthosine **8**: ¹H NMR (DMSO-d₆) δ 1.12 (s, 3H), 3.82–3.93 (m, 1H), 4.01–4.11 (m, 1H), 5.40 (broad m, 2H), 5.83 (s, 1H), 8.45 (s, 1H), 10.68 (s, 1H), 11.42 (s, 1H).

16. Korba BE, Montero AB, Farrar K, Gaye K, Mukerjee S, Ayers M, Rossignol J-F. *Antivir Res.* 2007; 77:56–63. [PubMed: 17888524]

Figure 1.
Modified nucleoside analogues exhibiting anti-HCV properties.

Figure 2.
Tricyclic C-2' modified target.

Scheme 1.
Initial route to the C-2'-Me intermediate.

Scheme 2.
Synthetic route to **1**.