Published in final edited form as:

Transplantation. 1991 October ; 52(4): 723–724.

SPLENIC ARTERY ANEURYSMS OCCURRING IN LIVER TRANSPLANT RECIPIENTS¹

Oscar Bronshter, Hadar Merhav, David Van Thiel, and Thomas E. Starzl²

Department of Surgery, University Health Center of Pittsburgh, University of Pittsburgh, Veterans Administration Medical Center, Pittsburgh, Pennsylvania

Splenic artery aneurysms (SAA)* are found in 0.1% of the cases in large autopsy series (1) and in 0.8% of unselected abdominal aortograms (2). The incidence of SAA is higher in patients with portal hypertension, in whom it is reported to occur in 8.8% to 50% of cases (3, 4). Rupture of SAA carries a high mortality rate.

Rupture of SAA in liver transplant recipients has been reported (4,5). To assess the importance of this complication in the liver transplant population, we reviewed the medical records of all liver recipients whose 1311 transplants were performed at the Presbyterian University Hospital from January 1, 1988 until July 1, 1990. Of 5 patients with ruptured SAA, 4 died. An additional patient was recognized to have a SAA following his second orthotopic liver transplantation and this was removed electively by splenectomy and distal pancreatectomy. Summaries of these 6 cases are in Table 1.

SAAs are the most common visceral arterial aneurysms and account for 60% of all aneurysms found within the splanchnic arterial bed (2). The pathogenesis of SAA is multifactorial, and Stanley et al. (2) have recognized 4 conditions that place patients at high risk: (1) arterial dysplasia, (2) portal hypertension, (3) focal arterial inflammatory processes, and (4) multiparity in women. Anatomically, about 70% of the SAAs in patients with cirrhosis and portal hypertension are located in the distal third of the artery, and half of the aneurysms are multiple (2,3). In our series, 4 of the 5 liver transplant patients had aneurysms greater than 2 cm in diameter, and one had multiple aneurysms.

Multiple factors could contribute to the higher incidence and larger size of SAAs in patients with chronic liver disease and portal hypertension. These include: increased splenic and overall splanchnic blood flow secondary to arteriovenous shunts and collateral formation; dilatation and elongation of the splenic artery (6,7); increased cardiac output and splanchnic vasodilatation from hyperglucagonemia (8); and vascular changes caused by other hormone changes, such as those which “feminize” male cirrhotic patients (9).

Whatever the explanation, the impact of SAA in liver transplantation needs emphasis. In a recent study at the Mayo Clinic (4), 60 patients with portal hypertension who were being considered for OLT were submitted to routine preoperative celiac angiography, and 5 (8.3%) were found to have SAA 8 to 25 mm in diameter. A sixth patient in this series developed a

¹This work was supported by research grants from the Veterans Administration and by project grant No. DK 29961 from the National Institutes of Health, Bethesda, Maryland.

Copyright © 1991 by Williams & Wilkins

²Address correspondence to: Thomas E. Starzl, M.D., Ph.D., Department of Surgery, 3601 Fifth Avenue, University of Pittsburgh, Pittsburgh, PA 15213.

* Abbreviation: SAA, splenic artery aneurysm.

SAA 3 months postoperatively. The size at which an asymptomatic SAA should arouse alarm has been reported to be 15 mm (4,5). There have been no reported ruptures of smaller SAAs in liver transplant recipients. Whatever the size, most SAAs are asymptomatic, as in 4 of our 5 patients. Pain in the mid upper or left upper quadrant of the abdomen is an ominous portent of imminent or contained rupture (2,10,11).

The question of critical size of SAA in liver transplant candidates or recipients should be left open until there is more information. The incidence of rupture of documented SAA in nontransplant patients is 3% to 10% (2), but extra risk factors in liver recipients could include the higher rupture rate following any intraabdominal operation (12), abrupt changes in celiac trunk blood flow caused by OLT (6,7), the addition of postoperative corticosteroids, inadvertent trauma to the aneurysm intraoperatively, opening of the retroperitoneal space, and the coagulopathy that often is a feature of perioperative recovery.

Only one previously reported patient has survived a post-transplant SAA rupture (4), and in our series, the mortality rate following rupture was 80%. Improvement will require identification of the pathology during pretransplantation workup. MRI is the most discriminating procedure, and we recommend it routinely. Doppler ultrasound of the splenic artery is less discriminating, and angiography is too dangerous in many patients with end-stage liver disease. With MRI, other essential information about liver size, portal vein patency, and the structure and flow patterns of the visceral arterial supply are obtained at the same time (13).

Operative management should include ligation of the splenic artery distal and proximal to the aneurysm and resection if feasible (5). Proximal splenic artery ligation alone is apt to be ineffective because of rich collateral arterial supply. A delayed operation may be indicated if multiple or large distal SAAs are found that can not be ligated without splenectomy at the time of orthotopic liver transplantation. Although there may be a role for splenic artery embolization before or after transplantation, we have not had personal experience. Splenic infarction and the formation of a splenic abscess is a potential complication of either splenic artery ligation without splenectomy or of embolization.

REFERENCES

1. Moore SW, Lewis RJ. Splenic artery aneurysm. *Ann Surg* 1961;153:1033. [PubMed: 13772355]
2. Stanley SW, Fry WJ. Pathogenesis and clinical significance of splenic artery aneurysms. *Surgery* 1974;6:898. [PubMed: 4428355]
3. Puttini M, Aseni P, Brambilla G, et al. Splenic artery aneurysms in portal hypertension. *J Cardiovasc Surg* 1982;23:490. [PubMed: 7153237]
4. Ayalon A, Wiesner RH, Perkins JD, et al. Splenic artery aneurysms in liver transplant patients. *Transplantation* 1988;45:386. [PubMed: 3278432]
5. Brems JJ, Hiatt JR, Klein AS, et al. Splenic artery aneurysm rupture following orthotopic liver transplantation. *Transplantation* 1988;45:1136. [PubMed: 3289154]
6. Nishida O, Moriyasu F, Nakamura T, et al. Hemodynamics of splenic artery aneurysms. *Gastroenterology* 1986;90:1042. [PubMed: 3512354]
7. Gittin N, Grahame GR, Kreel L, et al. Splenic blood flow and resistance in patients with cirrhosis before and after portocaval anastomosis. *Gastroenterology* 1970;19:208.
8. Lee SS, Morean R, Hadenque A, et al. Glucagon selectively increases splanchnic blood flow in patients with well-compensated cirrhosis. *Hepatology* 1988;8:1501. [PubMed: 3192162]
9. Van Thiel DH, Gavalier JJ, Cobb CF, et al. Is feminization in diabetic men due in part to portal hypertension? A rat model. *Gastroenterology* 1980;78:81. [PubMed: 7350040]
10. Trastek VF, Paivalero PC, Joyce JW, et al. Splenic artery aneurysm. *Surgery* 1982;91:694. [PubMed: 7079972]

11. Salo JA, Salmenkivi K, Tenhunen A, et al. Rupture of splanchnic artery aneurysms. *World J Surg* 1986;10:123. [PubMed: 3962324]
12. Swanson RJ, Littog FN, Hunt TK, et al. Laparotomy as a precipitating factor in the rupture of intraabdominal aneurysms. *Arch Surg* 1980;115:299. [PubMed: 7356384]
13. Edelman RR, Mattie HP, Atkinson DJ, et al. MR angiography. *Am J Radiol* 1990;154:937.

TABLE 1

Summary

OLTx No.	Age (yrs)	Sex	Diagnosis ^a	Time of rupture post OLTx	Outcome	LFT's at time of presentation	Pathology	Comments
1921	32	F	HVC hepatitis	45 days	Died	SGOT SGPT PHOS GGT Bili 104 IU/L 62 IU/L 156 IU/L 103 IU/L 11.2 mg%	3-cm solitary SAA with necrotic wall	1 Elective splenectomy 8 years pre-OLT 2 Splenic artery ligated as treatment for rupture 3 Second OLT postoperative day 50 4 Died postoperative day 124
1725	43	F	Autoimmune hepatitis	49 days	Died	SGOT SGPT PHOS Bili 157 IU/L 151 IU/L 210 IU/L 1.3 mg%	3-cm solitary SAA, splenic artery tortuous and thin-walled	1 Sudden death in hospital 2 Autopsy showed rejection of liver
1897	37	M	Sclerosing cholangitis	56 days	Died	SGOT SGPT PHOS GGT Bili 18 IU/L 23 IU/L 80 IU/L 76 IU/L 10mg%	No size of aneurysm reported	1 First OLT primary graft dysfunction 2 Retransplanted on postoperative day 3 3 Explored for bleeding on postoperative day 4 4 Died suddenly at home on postoperative day 56
1552	37	F	HBV hepatitis	10 days	Died	SGOT SGPT GGT PHOS Bili 23 IU/L 33 IU/L 37 IU/L 46 IU/L 0.9mg%	2-cm SAA with arterial sclerosis and perforation	1 Collapsed in the hospital 2 Died of irreversible shock 24 hr after splenectomy
1352	22	M	HVC hepatitis	–	Alive	SGOT SGPT PHOS GGT Bili 31 IU/L 4 IU/L 147 IU/L 57 IU/L 2.1 mg%	Tortuous artery with multiple aneurysms, largest 2 cm	1 First OLT lost 14 months postoperative to recurrent HVC hepatitis 2 Following second OLT complained of persistent left upper quadrant pain 3 Angiography revealed lesion 4 Treated with splenectomy and distal pancreatectomy
1568	36	F	Sclerosing cholangitis	6 days	Alive	SGOT SGPT PHOS GGT Bili 67 IU/L 416 IU/L 187 IU/L 190 IU/L 5.9 mg%	No size of aneurysm reported	1 Collapsed in the hospital 2 Underwent emergency splenectomy 3 Alive and well

^a All had chronic liver disease and cirrhosis. HVC, C hepatitis; HBV, B hepatitis; PNC, postnecrotic cirrhosis; OLT, orthotopic liver transplantation.