Published in final edited form as:

Biol Psychiatry. 2010 December 1; 68(11): 1055–1060. doi:10.1016/j.biopsych.2010.08.014.

Infralimbic D2 receptors are necessary for fear extinction and extinction-related tone responses

Devin Mueller, Christian Bravo-Rivera, and Gregory J. Quirk

Departments of Psychiatry and Anatomy & Neurobiology, University of Puerto Rico School of Medicine, P.O. Box 365067, San Juan, Puerto Rico 00936-5067

Abstract

Background—Fear extinction is dependent on plasticity in the infralimbic prefrontal cortex (IL), an area heavily innervated by midbrain dopaminergic inputs. Dopamine D2 receptors are concentrated in infralimbic output neurons that are involved in the suppression of conditioned fear after extinction. Here we examined the specific role of the D2 receptor in mediating associative learning underlying fear extinction, using the selective D2 antagonist raclopride.

Methods—Raclopride was administered systemically or infused into IL prior to fear extinction, and extinction retention was tested the following day. Rats were also prepared for single unit recording in IL to assess the effect of raclopride on firing properties.

Results—We found that systemic injection of raclopride given prior to extinction impaired retrieval of extinction when tested drug free the next day, but also induced catalepsy during extinction training. To determine whether impaired extinction was due to impaired motor function or disruption of extinction consolidation, we infused raclopride directly into IL. Raclopride infused immediately before extinction training did not produce motor deficits, but impaired recall of extinction when tested drug free. Furthermore, in animals that underwent extinction training, systemic raclopride reduced the tone responsiveness of IL neurons in layers 5/6, with no changes in average firing rate.

Conclusion—We suggest that D2 receptors facilitate extinction by increasing the signal-to-noise of IL neurons that consolidate extinction.

Keywords

fear conditioning; medial prefrontal cortex; dopamine; raclopride; anxiety disorder; schizophrenia

Introduction

The D2 family of dopamine receptors is intimately involved in affective, motor, and cognitive functions, many of which are mediated by the prefrontal cortex (1,2). D2 receptor signaling in prefrontal cortex has been linked to working memory function (3), reversal learning (4), and

© 2010 Society of Biological Psychiatry. Published by Elsevier Inc. All rights reserved.

Correspondence should be addressed to: Devin Mueller, Ph.D., Department of Psychology, University of Wisconsin-Milwaukee, P.O. Box 413, Milwaukee, WI 53201-0413, USA, Tel: (414) 229-6113, Fax: (414) 229-5219, devinm@uwm.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosure/Conflict of Interest

All authors report no biomedical financial interests or potential conflicts of interest.

behavioral flexibility (5), and blocking these receptors plays a key therapeutic role in diminishing the positive symptoms of schizophrenia (6,7). Previous work has shown that D2 actions in prefrontal circuits are associated with motor control functions such as the suppression of prepotent responses (8). Similar suppression of behavioral responses is observed after extinction of conditioned fear, and converging evidence implicates the medial prefrontal cortex, particularly the infralimbic subregion (IL), in the consolidation of fear extinction (9–11). Whether D2 receptors in IL contribute to extinction consolidation, however, remains unknown.

D2 receptor density is particularly high in IL as compared to other prefrontal regions (12), and IL is heavily innervated by dopaminergic projections from the ventral tegmental area (13). D2 receptor binding and transcripts are most prominent in Layer V neurons (14), which are the output neurons of the cortex. These neurons are thought to inhibit fear after extinction training by impeding amygdala output (15,16). The localization of D2 receptors on IL output neurons and the function of these neurons in fear suppression suggest that D2 receptors have a significant function in fear extinction. Thus, D2 receptor activity may critically modulate IL output, and extinction-related tone responses generated in IL are hypothesized to suppress fear responses (10). Moreover, fear extinction induces dopamine release in IL (17) suggesting that dopamine is involved in the consolidation of extinction.

While one might expect D2 receptor antagonists to impair extinction, a previous study in mice showed that antagonizing D2 receptors with systemic administration of sulpiride facilitated extinction (18). Therefore, to clarify the role of D2 receptors in extinction, we used raclopride, which has greater specificity and is more potent at antagonizing D2 receptors than sulpiride (19). We administered raclopride both systemically and intra-IL to determine the role of D2 receptors in the acquisition and retrieval of extinction. We also evaluated the effect of raclopride on the firing properties of IL neurons, specifically on tone responses after extinction.

Methods and Materials

Subjects

Male Sprague-Dawley rats (270–320g) were obtained, housed, and handled as described previously (20). Rats were restricted to 18 g of standard laboratory rat chow daily, and were subsequently trained to bar press for food pellets on a variable interval schedule (VI60). Throughout behavioral experiments, rats were able to press for food to maintain a constant level of activity against which freezing could be reliably measured (20). All procedures were approved by the Institutional Animal Care and Use Committee at the University of Puerto Rico in compliance with NIH guidelines.

Surgery

For infusion experiments, rats were implanted with a single 26-gauge stainless steel guide cannula (Plastics One, Roanoke, VA) aimed at IL (AP: +2.9, ML: –1.0, DV: –4.1 mm relative to bregma, angled 11° towards the midline in the coronal plane). After behavioral testing, rats were perfused with 0.9% saline followed by 10% buffered formalin. Brains were removed and stored in a 30% sucrose/10% formalin solution. 40 µm coronal sections were cut, mounted on slides, and stained for Nissl bodies in order to visualize the injector tip location.

A separate group of rats were surgically implanted with recording electrodes that consisted of drivable bundles of 16 microwires (22 µm, Stablohm 650; California Fine Wire, Grover Beach, California) as previously described (21). Electrodes were aimed at IL, located 2.9 mm anterior, 0.6 mm lateral, and 5.0 mm ventral from bregma. At the conclusion of the experiment, lesions were made at the tip of the recording wires by passing an anodal current of 25 µA for 18 s.

Rats were then perfused with 10% buffered formalin, and the brains were removed to mark the microlesions with a blue reaction of 6% ferrocyanide while fixing the tissue in 30% sucrose/10% buffered formalin. Locations of lesions were reconstructed onto coronal drawings adapted from Paxinos and Watson (22) from 40 μ m Nissl-stained sections.

Fear conditioning and extinction

All fear conditioning and extinction procedures were carried out in four identical operant boxes (Colbourn Instruments, Allentown, PA), located within sound-attenuating chambers. Between rats, shock grids were cleaned with soap and water and conditioning chamber walls were wiped clean. On day 1, rats received seven conditioning trials in which a tone (30 s, 4 kHz, 75 dB) co-terminated with a mild footshock (0.5 mA, 0.5 s). The intertrial interval varied, averaging 3 min. On day 2, rats were injected with either saline ($n = 9$) or raclopride (0.3 mg/kg, IP, $n = 11$) 10 min before extinction training (Experiment 1), saline ($n = 4$) or raclopride (0.1 mg/kg, IP, $n = 5$) 10 min before extinction training (Experiment 2) or infused with saline ($n = 9$) or raclopride (5 μ g in 0.5 μ l, $n = 11$) into IL before extinction training (Experiment 3) which consisted of 15 tone presentations in the absence of footshock. On day 3, rats were tested drug free with 15 presentations of the tone alone. In experiment 4, we infused saline ($n = 5$) or raclopride ($n = 6$) into IL 10 min prior to test (Day 3). Conditioned fear was assessed by measuring the percentage of time spent freezing during the tone.

Open field testing

To test the effects of raclopride on locomotor activity, rats were given injections of saline or raclopride (0.1 or 0.3 mg/kg, IP) 10 min before testing in an open field ($n = 4$ per group). Grid lines drawn on the floor of the arena (91.5 $\times$ 91.5 $\times$ 61 cm) divided it into a peripheral region (within 15.25 cm of the walls) and central region (61 $\times$ 61 cm) of approximately equal area. The number of line crosses and time spent in the central region were scored by an observer blind with respect to experimental groups.

Infusions of raclopride in IL

Infusions of raclopride (5 μ g, Sigma, St. Louis, MO) dissolved in saline or saline alone were made 10 minutes prior to extinction training in a volume of 0.5 μ l at a rate of 0.2 μ l/min. Following infusions, injectors were left in place for 2 min to allow the drugs to diffuse. The tip of the injection cannula extended 1.0 mm beyond that of the guide cannula. This dose of raclopride was chosen as it impairs acquisition of fear conditioning when infused in the amygdala (23), and has been infused into the medial prefrontal cortex (24).

Behavioral data analysis

Digital video was recorded during the behavioral procedures and was analyzed with Freezescan software (Clever Systems, Reston, VA). Total seconds freezing during the tone presentations were scored for each rat, and this number was expressed as a percentage of the total tone presentation time. All data are included in the analysis, but data are shown in blocks of two trials (necessarily excluding trial 7 of conditioning or trial 15 of extinction). Group comparisons were made using ANOVA or Student's *t*-tests (SPSS for Windows, SPSS Inc., Chicago, IL). Significant main effects were followed by Tukey *post hoc* comparisons.

Multi-channel unit recording

After surgery, rats were allowed 6 days to recover. Rats ($n = 3$) were fear conditioned with 5 tone-shock pairings, followed by extinction training consisting of 20 tone-alone presentations. The next day, rats received an additional 5 tone-alone presentations to assess extinction retention. Rats were then acclimated to recording procedures in the same chambers as in the behavioral experiments, and electrodes were driven in increments of 44 μ m until single units

were isolated with principle components analysis and template matching (Offline Sorter; Plexon, Dallas, Texas). Once cells in IL were well-isolated, we assessed the effects of injections of saline or raclopride (0.3 mg/kg, IP) on spontaneous activity while rats were in the operant chamber pressing for food. Five-minute sessions of spontaneous activity and 4-min sessions consisting of 3 tone CS presentations were recorded at four timepoints: 30 min before and 10 min after saline injection, and 30 min before and 10 min after raclopride injection. Firing rates before and after injections were compared with a Wilcoxon matched-pairs test. After recording the four sessions at a given location, the electrode drive was advanced in 80 μ m steps until new cells were found, and the experiment was repeated. Spike trains were analyzed with NeuroExplorer (NEX Technologies, Littleton, Massachusetts) to obtain firing rate and bursting. Bursts were defined as three or more successive spikes in which the first interspike interval was < 25 ms and subsequent intervals were < 50 ms (21). To measure tone-induced activity of neurons, the firing rate in the first 400 ms bin after tone onset was compared to the firing rate of 20 pre-tone bins of equal duration using a z-score transformation as previously described (10).

Results

Systemic blockade of dopaminergic D2-receptors impairs consolidation of extinction memory

We first examined whether dopamine, acting at D2 receptors, is necessary for extinction learning. Rats were fear conditioned on day 1, and were systemically injected with a D2 receptor antagonist (raclopride, 0.3 mg/kg, i.p.) or saline 10 min prior to extinction training on day 2. Rats injected with raclopride expressed significantly higher levels of freezing on average than rats injected with saline throughout the extinction session (Figure 1A). ANOVA revealed a main effect of group ($F_{1,15} = 15.6, p < 0.001$), but no group by trial interaction ($F_{14,252} = 1.2, p = 0.26$) indicating that raclopride-treated rats expressed more freezing overall during the session than saline-treated controls. Thus, raclopride augmented freezing behavior during extinction training. In a drug-free test the following day, raclopride-treated rats expressed higher levels of freezing to the tone than saline-treated rats. ANOVA revealed a main effect of group ($F_{1,18} = 6.7, p = 0.02$) and a group by trial interaction ($F_{14,252} = 2.3, p = 0.005$), indicating that raclopride-treated rats expressed higher levels of freezing than saline-treated rats at the beginning of the extinction retention test but gradually expressed less freezing across the test. Thus, it would appear that raclopride interfered with both extinction acquisition and consolidation. However, it is well known that D2 antagonists can induce catalepsy (25,26), which could interfere with extinction. Indeed, rats injected with 0.3 mg/kg (i.p.) raclopride showed significantly reduced pre-tone bar-press rates prior to the first extinction trial (Figure 1B; $t_{18} = 2.3, p = 0.03$) and reduced locomotor activity in an open field as compared to saline-injected controls (Figure 1B; $t_6 = 7.3, p < 0.001$). Thus, it is possible that the apparent blockade of extinction acquisition could be due instead to locomotor side effects of raclopride.

In an attempt to dissociate locomotor from extinction effects of raclopride, we repeated the experiment with a lower dose of raclopride (0.1 mg/kg, i.p.; Figure 1C). Raclopride-treated rats expressed similar levels of freezing as saline-treated rats throughout extinction training. ANOVA revealed no effect of group nor a group by trial interaction (all F 's < 1.0, $p > 0.75$). In a drug-free test the following day, both the raclopride- and saline-injected groups showed equally good retention of extinction. ANOVA revealed no effect of group or group by trial interaction on extinction retention (all F 's < 1.5, $p > 0.15$). The lower dose of raclopride also had no effect on pre-tone bar-press rates prior to the first extinction trial ($t_7 = 0.58, p = 0.6$; Figure 1D) and did not affect locomotor activity in an open field as compared to saline-injected controls ($t_6 = 0.30, p = 0.78$; Figure 1D). Thus, increased freezing observed during extinction training with the 0.3 mg/kg dose correlated with the cataleptic effects of the drug. Because we

were unable to dissociate locomotor effects from impaired retrieval of extinction, it remains possible that the apparent extinction deficit induced by systemic raclopride resulted from locomotor side effects.

Blockade of D2-receptors in IL impairs consolidation of extinction memory

D2 antagonists are thought to induce catalepsy by acting in the striatum (27–29). Therefore, to dissociate locomotor effects from effects on extinction consolidation, we infused raclopride directly into IL, a structure strongly implicated in consolidation of fear extinction. Rats were infused with raclopride (5 μ g in 0.5 μ l) or saline into IL 10 min prior to extinction training (see Figure 2A for infusion site for all rats). This dose of raclopride was chosen as it impairs acquisition of fear conditioning when infused in the amygdala (23). Unlike systemic administration, intra-IL raclopride did not affect expression of freezing or within-session extinction (day 2; Figure 2B). ANOVA revealed no effect of group nor a group by trial interaction (all F 's < 1.0 , $p > 0.60$). In addition, raclopride did not affect pre-tone bar-press rates prior to the first extinction trial relative to saline-infused controls (Figure 2C; $t_{18} = 0.06$, $p = 0.95$). The following day, however, raclopride-infused rats showed impaired retrieval of extinction relative to saline-infused rats (Figure 2B). ANOVA revealed a main effect of group ($F_{1,18} = 4.8$, $p = 0.04$) and a group by trial interaction ($F_{14,252} = 1.9$, $p = 0.03$), indicating that raclopride-infused rats expressed higher levels of freezing than saline-infused rats at the beginning of the extinction retention test, and gradually extinguished across the test. In a separate experiment, raclopride infusions prior to test did not impair the recall of extinction (Figure S1 in Supplement 1). Thus, dopaminergic D2 receptor blockade in IL impairs consolidation of fear extinction, but not the expression of extinction previously learned.

Blockade of D2-receptors reduces extinction-related tone responses in IL

Given the preferential expression of D2 receptors in the output layer of cortex (14), we expected that D2 receptors would have distinct effects on the firing properties of IL neurons engaged in fear extinction. To determine the physiological effects of D2 receptor blockade, we recorded from IL neurons within layers 5/6 in behaving rats that had received conditioning and extinction training (procedure shown in Figure 3A and placement of electrodes shown in Figure 3B). All rats showed significant within-session extinction, as demonstrated by an overall effect of trial ($F_{2,38} = 4.0$, $p < 0.001$), and good retention of extinction as they showed similar freezing levels to a tone CS at test as compared to the last trial of extinction ($t_2 = 0.8$, $p = 0.52$). Following the test, single units were isolated and spontaneous activity was recorded for 5 min followed by 3 tone CS presentations. This was repeated 10 min after a saline injection, 30 minutes prior to raclopride injection, and 10 min after raclopride injection, resulting in a within-cell statistical design. A total of 20 single units were analyzed.

Raclopride did not affect averaged firing rate (Wilcoxin matched-pairs test: $Z = 1.53$, $p = 0.13$; Figure 3C) or high-frequency bursting ($t_{19} = 1.46$, $p = 0.16$) as compared to saline. Tone responses in IL neurons reflect extinction learning, so we examined the effect of raclopride administration on the magnitude of extinction-related tone responses. Tone-evoked activity was recorded before and after injection of saline or raclopride (0.3 mg/kg, IP). Following saline administration, a subset of IL neurons (6/20, or 30%) exhibited significant conditioned tone responses within the first 400 ms after tone onset relative to the preceding 8 s prior to tone onset (all z 's > 1.96 , $p < 0.05$). Raclopride administration significantly attenuated tone responses in these IL neurons from an average $z = 5.07$ to $z = 2.50$ ($t_5 = 5.38$, $p = 0.003$; Figure 3D), and reduced the number of neurons exhibiting significant tone responses (4/6, or 67%). The peri-event time histogram for this subset of neurons shows that tone responses were robustly expressed at the onset of the tone, but activity rapidly returned to baseline levels in the first 400 ms of the tone period (Figure 3E). This decrease in tone-evoked activity was independent of changes in firing rate or high-frequency bursting, as raclopride did not affect

firing rate (Wilcoxin matched-pairs test: $Z = 1.57$, $p = 0.12$; Figure 3F) or high-frequency bursting ($t_5 = 0.10$, $p = 0.92$) in these tone-responsive neurons. Furthermore, each individual tone-responsive unit showed a marked decline in tone-evoked activity after raclopride treatment (Figure 3G), and no previously non-responsive units became tone responsive after raclopride treatment. The decline in tone response with raclopride could not be attributed to extinction with repeated testing, because no such decline was observed in saline-treated rats (Figure S2 in Supplement 1). Taken together, these results suggest that blockade of D2 receptors impairs fear extinction by disrupting IL tone responses that arise during extinction (30,31).

Discussion

The present study demonstrates that D2 receptor signaling is necessary for extinction of conditioned fear. Blockade of D2 receptors in IL during fear extinction impairs later retrieval of extinction, implicating D2 receptor signaling in this structure. Although raclopride did not affect firing rate or bursting in IL, it attenuated extinction-related tone responses. Thus, D2 receptor signaling in IL promotes fear extinction, presumably by enabling tone responses during extinction that are necessary for extinction consolidation (10,30,31).

Raclopride-induced impairment of fear extinction generally agrees with previous findings. Haloperidol, a predominantly D2 receptor antagonist with notable affinity for the D1 receptor, was recently shown to impair extinction when given systemically or into the nucleus accumbens (32). In addition, lesions of the dopaminergic terminals in mPFC impair contextual fear extinction (33). Our results implicate dopaminergic actions at the D2 receptor, and parallel the finding that D4 receptor activation in IL is also required for fear extinction (34). Ponnusamy and colleagues (18), however, observed accelerated extinction in mice using a different systemically-administered D2 receptor antagonist, sulpiride. The disparity with our results may be explained by the greater D2-receptor selectivity and binding strength of raclopride over sulpiride (19). Sulpiride is known to bind non-D2 receptor sites, an effect not observed with other neuroleptics (35). It should be noted that sulpiride did not enhance extinction of a cocaine-induced conditioned place preference (36). Our results cannot be explained by state dependent effects as raclopride is quickly metabolized, and infusing the drug prior to extinction retrieval did not impair retrieval of extinction.

We identified IL as a site of action of raclopride in fear extinction, and observed a significant reduction in tone responses of IL neurons after systemic raclopride injections. Tone responses in IL are correlated with extinction retention (10), and electrical stimulation designed to mimic tone responses in this region during extinction enhance extinction retention (10,37). Output from IL neurons inhibits fear after extinction training by impeding amygdala output (15,16). Because D2 receptors are located on IL output neurons, and conditioned responses in these neurons are observed during extinction (31,30), we argue that activation of D2 receptors augments tone responsiveness during extinction learning.

In addition to direct modulation of tone responses, D2 receptor activation also stimulates mitogen-activated protein kinases (MAPKs) (38,39). The MAPK signaling cascade is critical for protein synthesis, and reducing the activity of MAPK (9,40) or blocking protein synthesis (41,42) in IL impairs extinction retention. Thus, activation of D2 receptors on IL output neurons would facilitate consolidation of fear extinction.

Our results indicate that optimal levels of dopamine acting at D2 receptors are required for fear extinction to occur. D2 antagonists are commonly prescribed in psychiatry and are useful for the treatment of schizophrenia, a condition that is often comorbid with anxiety disorders. Our results suggest that the use of these drugs could lead to resistance to extinction, which may

impede exposure therapy for anxiety disorders in patients currently using D2 receptor antagonists. Recently, it was shown that patients with schizophrenia exhibit extinction recall deficits (43), an effect that could be attributed, in part, to the use of medications that block D2 receptors (44). Whether chronic use of these medications reduces their potential to impair extinction learning, however, remains to be determined.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by MH058883, MH081975 and the UPR President's Office to GJQ and a FQRNT (Quebec, Canada) postdoctoral fellowship to DM. We gratefully acknowledge the assistance of Ivan Vidal-Gonzalez with single-unit recording.

References

1. Mehta MA, Swainson R, Ogilvie AD, Sahakian J, Robbins TW. Improved short-term spatial memory but impaired reversal learning following the dopamine D(2) agonist bromocriptine in human volunteers. *Psychopharmacology (Berl)* 2001;159:10–20. [PubMed: 11797064]
2. Usiello A, Baik JH, Rouge-Pont F, Picetti R, Dierich A, LeMeur M, et al. Distinct functions of the two isoforms of dopamine D2 receptors. *Nature* 2000;408:199–203. [PubMed: 11089973]
3. Wang M, Vijayraghavan S, Goldman-Rakic PS. Selective D2 receptor actions on the functional circuitry of working memory. *Science* 2004;303:853–856. [PubMed: 14764884]
4. De Steno DA, Schmauss C. A role for dopamine D2 receptors in reversal learning. *Neuroscience* 2009;162:118–127. [PubMed: 19401217]
5. Floresco SB, Magyar O, Ghods-Sharifi S, Vexelman C, Tse MT. Multiple dopamine receptor subtypes in the medial prefrontal cortex of the rat regulate set-shifting. *Neuropsychopharmacology* 2006;31:297–309. [PubMed: 16012531]
6. Creese I, Burt DR, Snyder SH. Dopamine receptor binding predicts clinical and pharmacological potencies of antischizophrenic drugs. *Science* 1976;192:481–483. [PubMed: 3854]
7. Seeman P, Lee T, Chau-Wong M, Wong K. Antipsychotic drug doses and neuroleptic/dopamine receptors. *Nature* 1976;261:717–719. [PubMed: 945467]
8. Volkow ND, Wang GJ, Ma Y, Fowler JS, Wong C, Ding YS, et al. Activation of orbital and medial prefrontal cortex by methylphenidate in cocaine-addicted subjects but not in controls: relevance to addiction. *J Neurosci* 2005;25:3932–3939. [PubMed: 15829645]
9. Hugues S, Chessel A, Lena I, Marsault R, Garcia R. Prefrontal infusion of PD098059 immediately after fear extinction training blocks extinction-associated prefrontal synaptic plasticity and decreases prefrontal ERK2 phosphorylation. *Synapse* 2006;60:280–287. [PubMed: 16786530]
10. Milad MR, Quirk GJ. Neurons in medial prefrontal cortex signal memory for fear extinction. *Nature* 2002;420:70–74. [PubMed: 12422216]
11. Sotres-Bayon F, Cain CK, LeDoux JE. Brain mechanisms of fear extinction: historical perspectives on the contribution of prefrontal cortex. *Biol Psychiatry* 2006;60:329–336. [PubMed: 16412988]
12. Vincent SL, Khan Y, Benes FM. Cellular distribution of dopamine D1 and D2 receptors in rat medial prefrontal cortex. *J Neurosci* 1993;13:2551–2564. [PubMed: 8501521]
13. Swanson LW. The projections of the ventral tegmental area and adjacent regions: a combined fluorescent retrograde tracer and immunofluorescence study in the rat. *Brain Res Bull* 1982;9:321–353. [PubMed: 6816390]
14. Lidow MS, Wang F, Cao Y, Goldman-Rakic PS. Layer V neurons bear the majority of mRNAs encoding the five distinct dopamine receptor subtypes in the primate prefrontal cortex. *Synapse* 1998;28:10–20. [PubMed: 9414013]
15. Pare D, Quirk GJ, LeDoux JE. New vistas on amygdala networks in conditioned fear. *J Neurophysiol* 2004;92:1–9. [PubMed: 15212433]

16. Quirk GJ, Likhtik E, Pelletier JG, Pare D. Stimulation of medial prefrontal cortex decreases the responsiveness of central amygdala output neurons. *J Neurosci* 2003;23:8800–8807. [PubMed: 14507980]
17. Hugues S, Garcia R, Lena I. Time course of extracellular catecholamine and glutamate levels in the rat medial prefrontal cortex during and after extinction of conditioned fear. *Synapse* 2007;61:933–937. [PubMed: 17663454]
18. Ponnusamy R, Nissim HA, Barad M. Systemic blockade of D2-like dopamine receptors facilitates extinction of conditioned fear in mice. *Learn Mem* 2005;12:399–406. [PubMed: 16077018]
19. Simon VM, Parra A, Minarro J, Arenas MC, Vinader-Caerols C, Aguilar MA. Predicting how equipotent doses of chlorpromazine, haloperidol, sulpiride, raclopride and clozapine reduce locomotor activity in mice. *Eur Neuropsychopharmacol* 2000;10:159–164. [PubMed: 10793317]
20. Quirk GJ, Russo GK, Barron JL, Lebron K. The role of ventromedial prefrontal cortex in the recovery of extinguished fear. *J Neurosci* 2000;20:6225–6231. [PubMed: 10934272]
21. Burgos-Robles A, Vidal-Gonzalez I, Santini E, Quirk GJ. Consolidation of fear extinction requires NMDA receptor-dependent bursting in the ventromedial prefrontal cortex. *Neuron* 2007;53:871–880. [PubMed: 17359921]
22. Paxinos, G.; Watson, C. The rat brain in stereotaxic coordinates. 6th ed. New York: Elsevier/Academic Press; 2007.
23. Greba Q, Gifkins A, Kokkinidis L. Inhibition of amygdaloid dopamine D2 receptors impairs emotional learning measured with fear-potentiated startle. *Brain Res* 2001;899:218–226. [PubMed: 11311883]
24. Capriles N, Rodaros D, Sorge RE, Stewart J. A role for the prefrontal cortex in stress- and cocaine-induced reinstatement of cocaine seeking in rats. *Psychopharmacology (Berl)* 2003;168:66–74. [PubMed: 12442201]
25. Hillegaart V, Ahlenius S. Effects of raclopride on exploratory locomotor activity, treadmill locomotion, conditioned avoidance behaviour and catalepsy in rats: behavioural profile comparisons between raclopride, haloperidol and preclamol. *Pharmacol Toxicol* 1987;60:350–354. [PubMed: 3615345]
26. Wadenberg ML, Kapur S, Soliman A, Jones C, Vaccarino F. Dopamine D2 receptor occupancy predicts catalepsy and the suppression of conditioned avoidance response behavior in rats. *Psychopharmacology (Berl)* 2000;150:422–429. [PubMed: 10958084]
27. Alcock SJ, Hemsley KM, Crocker AD. Atropine acts in the ventral striatum to reduce raclopride-induced catalepsy. *Eur J Pharmacol* 2001;424:179–187. [PubMed: 11672560]
28. Ogren SO, Fuxe K. D1- and D2-receptor antagonists induce catalepsy via different efferent striatal pathways [corrected]. *Neurosci Lett* 1988;85:333–338. [PubMed: 2966309]
29. Sanberg PR. Haloperidol-induced catalepsy is mediated by postsynaptic dopamine receptors. *Nature* 1980;284:472–473. [PubMed: 7189016]
30. Laviolette SR, Lipski WJ, Grace AA. A subpopulation of neurons in the medial prefrontal cortex encodes emotional learning with burst and frequency codes through a dopamine D4 receptor-dependent basolateral amygdala input. *J Neurosci* 2005;25:6066–6075. [PubMed: 15987936]
31. Baeg EH, Kim YB, Jang J, Kim HT, Mook-Jung I, Jung MW. Fast spiking and regular spiking neural correlates of fear conditioning in the medial prefrontal cortex of the rat. *Cereb Cortex* 2001;11:441–451. [PubMed: 11313296]
32. Holtzman-Assif O, Laurent V, Westbrook RF. Blockade of dopamine activity in the nucleus accumbens impairs learning extinction of conditioned fear. *Learn Mem* 2010;17:71–75. [PubMed: 20154351]
33. Fernandez EE. Prefrontocortical dopamine loss in rats delays long-term extinction of contextual conditioned fear, and reduces social interaction without affecting short-term social interaction memory. *Neuropsychopharmacology* 2003;28:490–498. [PubMed: 12629528]
34. Pfeiffer UJ, Fendt M. Prefrontal dopamine D4 receptors are involved in encoding fear extinction. *Neuroreport* 2006;17:847–850. [PubMed: 16708027]
35. Csernansky JG, Csernansky CA, Hollister LE. 3[H]-Sulpiride labels mesolimbic non-dopaminergic sites that bind antidepressant drugs. *Experientia* 1985;41:1419–1421. [PubMed: 4065296]

36. Yim AJ, Andersen ML, Soeiro AC, Tufik S, Oliveira MG. Acute systemic blockade of D2 receptors does not accelerate the extinction of cocaine-associated place preference. *Brain Res* 2009;1304:122–128. [PubMed: 19766608]
37. Milad MR, Vidal-Gonzalez I, Quirk GJ. Electrical stimulation of medial prefrontal cortex reduces conditioned fear in a temporally specific manner. *Behav Neurosci* 2004;118:389–394. [PubMed: 15113265]
38. Neve KA, Seamans JK, Trantham-Davidson H. Dopamine receptor signaling. *J Recept Signal Transduct Res* 2004;24:165–205. [PubMed: 15521361]
39. Beaulieu JM, Sotnikova TD, Marion S, Lefkowitz RJ, Gainetdinov RR, Caron MG. An Akt/beta-arrestin 2/PP2A signaling complex mediates dopaminergic neurotransmission and behavior. *Cell* 2005;122:261–273. [PubMed: 16051150]
40. Hugues S, Deschaux O, Garcia R. Postextinction infusion of a mitogen-activated protein kinase inhibitor into the medial prefrontal cortex impairs memory of the extinction of conditioned fear. *Learn Mem* 2004;11:540–543. [PubMed: 15466305]
41. Mueller D, Porter JT, Quirk GJ. Noradrenergic signaling in infralimbic cortex increases cell excitability and strengthens memory for fear extinction. *J Neurosci* 2008;28:369–375. [PubMed: 18184779]
42. Santini E, Ge H, Ren K, Pena dO, Quirk GJ. Consolidation of fear extinction requires protein synthesis in the medial prefrontal cortex. *J Neurosci* 2004;24:5704–5710. [PubMed: 15215292]
43. Holt DJ, Lebron-Milad K, Milad MR, Rauch SL, Pitman RK, Orr SP, et al. Extinction memory is impaired in schizophrenia. *Biol Psychiatry* 2009;65:455–463. [PubMed: 18986648]
44. Martinez KG, Quirk GJ. Extending fear extinction beyond anxiety disorders. *Biol Psychiatry* 2009;65:453–454. [PubMed: 19236997]
45. Swanson, LW. *Brain maps: structure of the rat brain*. New York: Elsevier; 1992.

Figure 1. Systemic raclopride induces mild catalepsy and impairs long-term retention of fear extinction

(A) Systemic injections of 0.3 mg/kg raclopride (*arrow*) before extinction on day 2 led to a significant increase in percent freezing and impaired retention of extinction on day 3. (B) This dose of raclopride (0.3 mg/kg) reduced pre-tone bar-press rates during extinction and reduced locomotor activity as measured by line crosses in an open field. (C) Systemic injections of 0.1 mg/kg raclopride (*arrow*) before extinction did not affect percent freezing, nor impaired retention of extinction on day 3. (D) This dose of raclopride (0.1 mg/kg) did not affect pre-tone bar-press rates during extinction or locomotor activity in an open field. Data shown in blocks of 2 trials for A and C. * $p < 0.05$, *** $p < 0.001$

Figure 2. Infusion of raclopride into IL impairs long-term retention of fear extinction, without inducing catalepsy

(A) Coronal drawings (bregma, 3.20 mm, adapted with permission from (45)) show placements of injector tips for all rats in IL. (B) Before extinction, rats were infused (*arrow*) with either the D2 receptor antagonist raclopride or saline. Rats infused with raclopride into IL showed normal within-session extinction, but were impaired in their recall of extinction on day 3. Data shown in blocks of 2 trials. (C) Raclopride infusions into IL did not affect pre-tone bar-press rates during extinction. * $p < 0.05$

Figure 3. Raclopride attenuates extinction-related tone-evoked activity in IL units

(A) Coronal drawing shows location of recording electrodes (bregma + 2.80 mm; adapted with permission from (45)). (B) Systemic raclopride had no effect on firing rate relative to saline. (C) In tone-responsive neurons, systemic raclopride attenuated the magnitude of tone-evoked responses in the first 400 ms after tone onset. (D) The reduction in tone responses are shown in the group peri-event time histogram (bin width 100 ms). (E) Raclopride did not alter firing rate of these tone-responsive neurons, but (F) markedly reduced tone responsiveness in each of these individual neurons (dashed line indicates significant tone response, $p < 0.05$). $**p < 0.01$.