A stochastic, evolutionary model for range shifts and richness on tropical elevational gradients under Quaternary glacial cycles

Robert K. Colwell^{1,2,*} and Thiago F. Rangel³

¹*Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT 06269-3043, USA*

²*Center for Macroecology, Evolution and Climate, Department of Biology, University of Copenhagen, Universitetsparken 15, 2100 Copenhagen O, Denmark*

³*Departamento de Ecologia, ICB, Universidade Federal de Goiás, Caixa Postal 131, 74.001-970 Goiânia, GO, Brasil*

Quaternary glacial–interglacial cycles repeatedly forced thermal zones up and down the slopes of mountains, at all latitudes. Although no one doubts that these temperature cycles have left their signature on contemporary patterns of geography and phylogeny, the relative roles of ecology and evolution are not well understood, especially for the tropics. To explore key mechanisms and their interactions in the context of chance events, we constructed a geographical range-based, stochastic simulation model that incorporates speciation, anagenetic evolution, niche conservatism, range shifts and extinctions under late Quaternary temperature cycles along tropical elevational gradients. In the model, elevational patterns of species richness arise from the differential survival of founder lineages, consolidated by speciation and the inheritance of thermal niche characteristics. The model yields a surprisingly rich variety of realistic patterns of phylogeny and biogeography, including close matches to a variety of contemporary elevational richness profiles from an elevational transect in Costa Rica. Mountaintop extinctions during interglacials and lowland extinctions at glacial maxima favour mid-elevation lineages, especially under the constraints of niche conservatism. Asymmetry in temperature (greater duration of glacial than of interglacial episodes) and in lateral area (greater land area at low than at high elevations) have opposing effects on lowland extinctions and the elevational pattern of species richness in the model—and perhaps in nature, as well.

Keywords: biogeography; extinction; mid-domain effect; simulation; speciation; stochastic model

1. INTRODUCTION

From our viewpoint in the present, with anthropogenic global warming challenging the evolutionary adaptations of many species (Clark *et al.* 2003; Feeley *et al.* 2007; Deutsch *et al.* 2008; Sinervo *et al.* 2010) as the Earth's temperature nears its highest point in the past million years (Hansen *et al.* 2006), it is easy to forget that adaptations in contemporary lineages have also been repeatedly tested and shaped by cold during the glacial phases of the past 2.6 Ma (the Quaternary Period). In fact, because of the asymmetry of the peaks and valleys of temperature, the total time elapsed during cold glacial periods is more than twice the time that the Earth has spent in warm interglacials since the advent of dominant 100-thousand-year temperature cycles in the late Pleistocene (Rutherford & D'Hondt 2000; Liu *et al.* 2008; figure 1). Although continental plate movements and mountain uplift were relatively insubstantial over this geologically short period (Garzione *et al.* 2008), the freezing and melting of ice masses

produced profound changes in shorelines, inland lakes, river courses, continental shelf islands and land bridges. Meanwhile, continental ice-sheets repeatedly altered the poleward limits and alpine glaciers the elevational bounds for terrestrial life, and regional patterns of precipitation shifted, sometimes profoundly, under the influence of temperature-driven changes in global circulation (Marchant *et al.* 2009). The biogeographical drama played out on this shifting physiographic and climatic template, among essentially modern lineages, continues to be revealed by stoichiometry, by pollen and phytoliths, by plant and animal macrofossils, and by phylogeographic analysis of living populations (e.g. Colinvaux *et al.* 1996; Schluter & Rambaut 1996; MacFadden *et al.* 1999; Marchant *et al.* 2002, 2009; Bush *et al.* 2004; Hooghiemstra & Van der Hammen 2004; Bush & Flenley 2007; Dick & Heuertz 2008; Blois *et al.* 2010; Voelker 2010).

It is now known that the tropics were not spared (Rutherford & D'Hondt 2000; Marchant *et al.* 2009). Estimates for the scope of terrestrial temperature change in the tropics, at any given elevation, range from 4°C to 10°C between glacial lows and interglacial peaks (Wille *et al.* 2001), accompanied by complex changes in the amount and seasonality of precipitation

* Author for correspondence (colwell@uconn.edu).

One contribution of 16 to a Discussion Meeting Issue 'Biological diversity in a changing world'.

Figure 1. Late Quaternary temperature fluctuations. (a) The EPICA Dome C Antarctic Ice Core 800 kyr dataset (Jouzel *et al.* 2007). (b) The EPICA series resampled at 1605 kyr equal intervals by linear interpolation, rescaled to the estimated Quaternary range of equatorial, continental temperatures at sea level.

(Bush & Silman 2004; Marchant *et al.* 2009) and the advance and retreat of high-elevation glaciers (Hooghiemstra & Van der Hammen 2004). Although mean annual temperature (MAT), at any given elevation, declines almost linearly with latitude outside the tropics, elevation-specific MAT varies little across tropical latitudes (Ahrens 2006), a pattern that has apparently persisted, on average, throughout the Pleistocene (MacFadden *et al.* 1999). By contrast, the decline of temperature with elevation (the lapse rate) is similar at all latitudes (Ahrens 2006). Thus, among temperate and boreal taxa, temperature-driven Quaternary range shifts were both latitudinal and elevational (Davis & Shaw 2001), a pattern now being repeated under current warming trends (e.g. Hickling *et al.* 2006; Moritz *et al.* 2008). By contrast, Quaternary changes in geographical distributions in the tropics were dominated by elevational range shifts and by responses to changing regional gradients in precipitation (Bush & Flenley 2007; Marchant *et al.* 2009). Documented upslope elevational shifts in the tropics are already emerging under current global warming trends (e.g. Chen *et al.* 2009).

Within the time frame of the late Quaternary, contemporary biogeographical patterns of terrestrial species can be viewed as the product of a complex interplay of ecological and evolutionary processes operating largely on pre-existing lineages, constrained by space and climate, all in the context of chance events. In this study, we aim to capture the essence of some of these processes in a simple, stochastic model that simulates both ecological and evolutionary processes affecting the elevational ranges of tropical species through late Quaternary time. Like all models, this one simplifies reality. We model tropical

temperatures realistically (albeit approximately), based on an empirical temperature series (figure 1) and estimated lapse rates, but do not attempt to model precipitation (still a daunting challenge for palaeoclimates; Marchant *et al.* 2009), edaphic factors, habitat specialization, changes in carbon dioxide or changes in sea level. In the model, geographical ranges are represented by their projection on an elevational axis, with their lateral extent modelled as a function of elevational band area. Because temperature is the only environmental variable considered, the evolution of the thermal tolerances of each species, which determine its potential elevational range, is modelled in a one-dimensional Hutchinsonian niche space. We model the tension between thermal niche conservatism, which forces elevational range shifts in response to temperature change, and *in situ* adaptation to shifting temperature, which promotes range stasis. The model tracks species richness and the phylogenetic structure of assemblages over time, as lineages undergo both speciation and extinction. To assess the ability of the model to produce realistic elevational patterns of species richness, we examine the parameter values that optimize fit to the contrasting elevational patterns of species richness of four contemporary species groups that share a 2900 m elevational gradient in Costa Rica. Throughout, we emphasize the role played by stochastic events in producing enduring patterns.

2. MATERIAL AND METHODS

(a) *Model design*

Our model tracks the geographical distribution of species on a bounded, tropical elevational gradient (the geographical *domain*) over the course of temperature cycles of the past 800 kyr (thousand years). The model is explicitly constructed upon the duality between the environmental tolerance limits of species in niche space and the projection of those limits on an environmentally heterogeneous geographical domain (Rangel *et al.* 2007; Colwell & Rangel 2009). In the model, niche space consists of a single environmental axis representing temperature, and the geographical domain (in the simplest version of the model) consists of a single elevational transect somewhere in the wet tropics, from sea level to the top of a mountain massif. Within this spatial domain, only elevation (the geographical *z*-coordinate) is modelled explicitly, and not distances along the ground (geographical *x*- and *y*-coordinates). As an option, we use the areal (*x,y*) extent of land within elevational bands on the massif to model its lateral extent and shape. The niche space is continuous; the geographical domain is discretized into an arbitrary (large) number of ordered, equal elevational steps. Changes in sea level are not considered, and we have not attempted to model patterns of precipitation or carbon dioxide (Marchant *et al.* 2009), short-term, abrupt climate shifts (Alley *et al.* 2003) or climate refuges based on within-elevation spatial heterogeneity (Ashcroft 2010).

The model was implemented in Delphi in the Embarcadero RAD Studio 2010 for Windows

Table 1. Model parameters and settings.

	meaning	figure 3	figures 4 and 6	figure 5
parameters				
F	number of founders	10	10	10
K	species number ceiling	25	100	100
σ	founder niche breadth	1.50	1.50	1.50
Δ	niche centre evolution	0.10	0.10	varies
β	niche breadth evolution	0.10	0.10	0.10
α	extinction	0.10	varies	0.01
γ	speciation	0.06	0.05	0.05
area	lateral area effect	off	on	on
settings				
domain	maximum elevation (m) (minimum is 0)	2900	5000	5000
ElevSteps	number of elevational steps	140	100	100
climate	temperature series over time	EPICA	EPICA	EPICA
min. temp	minimum temperature at sea level at glacial maximum ($^{\circ}\text{C}$)	23	23	23
max. temp	maximum temperature at sea level at glacial minimum ($^{\circ}\text{C}$)	28	28	28
lapse	wet adiabatic lapse rate ($^{\circ}\text{C km}^{-1}$ elevation)	5	5	5

environment, as a multi-threaded, compiled application, using visual display libraries from Spatial Analysis in Macroecology (Rangel *et al.* 2010).

(b) Palaeotemperature data

Palaeotemperatures from 800 kyr before present until 1950 were derived from the EPICA Dome C Antarctic Ice Core dataset (Jouzel *et al.* 2007), which models palaeotemperature based on oxygen isotope ratios (figure 1a). Because the time-spacing of EPICA samples increases sharply with core depth (age), we resampled the original time series (5800 data points) at 160 constant intervals of 5 kyr, by linear interpolation (figure 1b), for use as the discrete time steps required in the model.

Based on estimates of tropical, terrestrial temperatures at the Last Glacial Maximum (LGM; Flenley 1998; Wille *et al.* 2001; Bush *et al.* 2004), the coldest point in the EPICA series, we assumed that sea-level temperature at the equator varied between a minimum of 23°C at the LGM and a maximum of 28°C at the previous (Eemian) interglacial, the warmest point in the EPICA series. For the model, the resampled EPICA series was proportionally rescaled to this range (figure 1b).

The temperature at each elevation on the geographical domain at each point in the resampled, rescaled EPICA series was estimated by assuming a 5°C temperature decline for each 1000 m elevation above sea level, a wet adiabatic lapse rate typical for contemporary tropical climates (e.g. Wille *et al.* 2001; Bush *et al.* 2004; Colwell *et al.* 2008). Although tropical lapse rates may, at times, have been somewhat greater than at present during the late Quaternary and may under some conditions be somewhat larger at high elevations (Wille *et al.* 2001), these differences are not sufficiently well documented or well understood to include them in our model. The scope of niche space represented on the geographical domain over the course of the time series covers the full range of temperatures, from the coldest, full-glacial mountaintop temperature (which depends upon the

height of the modelled massif) to the warmest, interglacial sea-level temperature.

As a control, a flat temperature time series was constructed, the same length as the resampled EPICA series, but representing a constant sea-level temperature of 25.5°C , with corresponding elevation-specific values based on a 5°C lapse rate.

(c) Founders

In the model, before launching the time-based simulation, the geographical domain is first populated with F (a model parameter, table 1) founders, representing pre-existing (Early Pleistocene) lineages that collectively become the ancestors of all subsequent species on the domain. For each potential founder, a niche centre (temperature optimum) is selected at (uniform) random along the thermal niche axis. To allow truncated niches (Colwell & Rangel 2009; Feeley & Silman 2010), the thermal niche axis is modelled to encompass temperatures spanning three times the range of elevational temperatures expressed on the domain at Time 0 in the modelled time series. (The niche axis extends symmetrically both above and below the temperature extremes initially expressed on the domain.)

A thermal niche breadth (in degree-Celsius) for each founder is selected from a normal distribution, with a mean at the niche centre and standard deviation σ (a model parameter, table 1). The niche extends symmetrically to each side of the niche centre. The niche is then projected onto the geographical domain (in its Time 0 climatic state) to determine the initial elevational range of the founder, regardless of whether the niche is fully expressed or, instead, truncated at sea level or mountaintop (Colwell & Rangel 2009; Feeley & Silman 2010). If the modelled niche does not correspond to any portion of the geographical domain, the species is not a successful founder, and a new random niche centre and a new random niche breadth are selected to replace it. This process, which repeats until the specified number of founders is reached, guarantees a stochastically uniform pattern

of species richness across the full domain (model 1 of Colwell & Hurtt 1994), avoiding any mid-domain effect (Colwell & Lees 2000) among the founders. Once the domain has been populated with founders and environmental changes begin, the niche centre, niche breadth and distribution of all species on the domain are completely under the control of the model dynamics, and the initial niche breadth parameter σ no longer plays any role.

(d) *Anagenetic (within-species) adaptive evolution and niche conservatism*

Once the simulation is launched, the temperature at each elevation on the domain changes at each time step in accord with the tropically rescaled EPICA time series (figure 1b). At each step in the simulation, following the change in temperature, each living species is evaluated for discordance between its temperature optimum in niche space (before the environmental temperature change) and the mean temperature of its distribution on the domain after the temperature change.

When niche and distribution are discordant, selection favours an adaptive evolutionary shift in the niche that will decrease or eliminate the difference between the niche optimum and the mean environmental conditions (Smith *et al.* 1995; Kirkpatrick & Barton 1997; Davis *et al.* 2005). On the other hand, an upslope or downslope range shift may also bring the geographical distribution into accord with the niche, without any evolution—but only if the range is not already too close to sea level (in a cooling climate) or mountaintop (in a warming climate; Colwell & Rangel 2009). The model assumes that range shifts into contiguous, thermally suitable portions of the domain are always successful. Sometimes, however, following environmental temperature change in a single time step, the new potential range of a species with narrow thermal tolerances maps on the domain outside its range in the previous time step. As detailed in §2e, these ‘range shift gaps’ (Colwell *et al.* 2008) are assumed to be insurmountable, and extinction ensues.

When ranges track climatic zones with little or no adaptation, the range shifts may be described as evidence for niche conservatism (Martinez-Meyer *et al.* 2004; Losos 2008). Based on an approach introduced by Rangel *et al.* (2007), two niche-evolution parameters control how much adaptive anagenetic evolution is permitted (or conservatism enforced) between time steps in the model. If D_1 is the distance in niche space ($^{\circ}\text{C}$) between the old niche optimum and the current mean environmental temperature of a species’ realized range, and D_2 is the distance between the old niche optimum and the new optimum after selection, then $D_2 = D_1 \times \Delta$. Thus, for perfect niche conservatism, model parameter $\Delta = 0$, whereas for perfect adaptation, $\Delta = 1$, although Δ may take any value between these limits (table 1). A second model parameter, β , regulates evolution or conservatism in niche breadth: $W_2 = W_1 \times |N(1, \beta)|$, where W_1 is a species’ niche breadth before and W_2 is its niche breadth after evolutionary change, and $N(1, \beta)$ is a normal distribution with mean 1 and variance

Figure 2. The extinction function. Based on a single parameter, α , the function scales probability of stochastic extinction, per species per time step, to range size r and geographical domain size v (Rangel *et al.* 2007).

β . Thus, for perfect niche conservatism, parameter $\beta = 0$, whereas values of $\beta > 0$ allow stochastic evolutionary change in niche breadth, which may prove adaptive (table 1).

Once these evolutionary adjustments in each species’ niche have been made, the niche is freshly projected onto the domain to establish the realized geographical range. If little or no evolution has been permitted, the new range will have shifted to track the corresponding upslope or downslope shift of climatic thermal zones, to the degree possible given the bounds of the spatial domain.

(e) *Extinction*

The model incorporates two kinds of extinction. After each step of temperature change and subsequent niche evolution (as described above), each newly re-projected range is considered for *stochastic extinction*, based on its elevational range size as a proxy for population size, which is commonly considered the primary risk factor for such extinctions (Purvis *et al.* 2000). The relationship between range size and probability of extinction is governed by model parameter α (table 1) by evaluating $P_{\text{ext}} = -\ln(r/v)(1/\alpha)$, where P_{ext} is the probability of extinction, r is the range size and v is the full extent of the domain (figure 2; Rangel *et al.* 2007). Variables r and v are measured either in metres of elevation or, as an option, as the areal extent of land within elevational bands bounded by the species’ elevational limits (r) and the sum of all elevational areal bands on the massif (v). To explore the effect of lateral area on model outcomes, we used a right cone with base radius equal to its height (approximating a stratovolcanic cone), as well as an empirical mountain gradient in Costa Rica, as described in a later section.

In contrast with stochastic extinctions, *deterministic extinctions* occur when temperature change during a single time step of the model leaves a species without any portion of the domain that corresponds with the scope of its thermal niche (Williams *et al.* 2007), leaving no population to evolve. Three kinds of deterministic extinctions occur in the model: *mountaintop extinctions* during warming phases, *sea-level extinctions*

during cooling phases and *range-shift gap extinctions* (Colwell *et al.* 2008) that befall species at mid-elevations (described in §2d). Narrow ranges (initially as a result of small σ , but ranges may become stochastically narrow as well) predispose species to deterministic distinctions as well as stochastic ones. Note that the number of time steps (160 in our model) chosen for the EPICA resampling affects the rate of deterministic extinctions as well, by controlling the mean shift in climate in an average step—perhaps, a reasonable proxy for the rate of environmental change in relation to biological response.

(f) *Speciation: cladogenetic adaptive evolution and niche conservatism*

Speciation in the model is conceived as allopatric or parapatric, with the niche and distribution of a child species diverging, stochastically, from its parent species, which itself remains initially unchanged. The child species is envisioned as inhabiting a neighbouring but not overlapping region of the massif, perhaps an adjacent watershed (Jansson & Dynesius 2002; Fjeldså & Rahbek 2006). A parent species is chosen at random from among species that have survived extinction in the present time step, with probability γ (a model parameter, table 1). The degree to which the thermal niche, and therefore the elevational distribution, of the child species resembles its parent's niche and distribution after cladogenesis is governed just as for anagenetic evolution and range shifts between time steps. The child's niche evolves towards the optimum for its realized range to the degree permitted by the niche centre conservatism/evolution parameter Δ and expands or contracts stochastically to the degree allowed by the niche breadth conservatism/evolution parameter β .

As an option, a ceiling of K species may be imposed on the model, imposing a degree of diffuse interspecific competition and forcing a degree of lineage selection on the model (and as a practical matter, preventing the runaway accumulation of species). If a ceiling has been set, once it is reached, speciation occurs only to the degree that extinction reduces the total number of living species below K .

(g) *Model outputs and behaviour*

A model run (realization) begins at EPICA Time 0 (800 kyr ago) and ends at Time 160 (1950), or earlier if all lineages go extinct. The programme maintains a species 'vital statistics' table, recording for each species its time of origination (0 for founders), time and cause (deterministic or stochastic) of extinction, parent species (if not a founder), number of descendant species, and initial and final values for niche size and breadth ($^{\circ}\text{C}$) and range size (in m elevation). Biogeographic data, consisting of the presence or absence of each species at each time step and at each elevation, yield a time versus elevation plot of species richness (figures 3d and 4), as well as time-based trends in patterns of range size and elevational location (figure 5).

From parent–child relationships and times of origination and speciation, a fully dichotomous phylogeny is recorded (in Newick notation) for each

founder's descendants, with time-based branch lengths (figure 3a). On the basis of phylogeny and the modelled biogeographic data, the phylogenetic structure of assemblages at each time step is assessed for each elevation (figure 3e) based on the 'average taxonomic distinctness' (Warwick & Clarke 1998) among species at each elevation at each time step (see figure caption for details). This pattern can be compared with the corresponding pattern of richness (figure 3d) as an indication of lineage selection and lineage mixing.

We combined visualization and statistical tools to explore the behaviour of the model and its parameters (table 1). Simple statistical procedures are documented in context in §3 or in figure captions. The more complex methods we used for pattern matching are explained in §2h.

(h) *Target pattern matching and parameter assessment*

To assess the potential of the model to reproduce empirical richness patterns on a tropical elevational gradient, we used four contrasting elevational records of species richness as 'target patterns' and systematically analysed the model parameter space to maximize the correspondence between observed and modelled patterns. The empirical datasets represented four ecologically and phylogenetically distinct groups of organisms on the same gradient, the Barva Transect in Costa Rica (tables S1–S4 in Colwell *et al.* 2008): 568 species of epiphytes (Cardelús *et al.* 2006), 82 species of understory Rubiaceae (Gilman 2007), 739 species of geometrid moths (Brehm *et al.* 2007) and 495 species of ants (Longino *et al.* 2002).

For all Barva Transect model runs, a maximum elevation of 2900 m was set, to match the actual scope of the transect, with the domain divided into 140 steps of 20 m elevation each, with a lapse rate of 5°C per 1000 m elevation (electronic supplementary material and methods in Colwell *et al.* 2008). To assess the effect of lateral area on extinction and patterns of richness, we applied an estimate of the land area in each 100 m elevational band on the Atlantic slope of Costa Rica (Gilman 2007) to the extinction function (figure 2).

For each of the four target richness patterns, a parameter space comprising nearly 50 000 combinations of parameter values was considered and formally explored, with a set of 10 replicate simulations (a 'replicate set') per parameter value combination (table 2). The results for different replicates within a replicate set vary among replicates only as result of the stochastic elements of the model. We did not attempt to match the absolute number of species in the target patterns, only the shape of the distribution (figure 7). Because it is sensitive to all aspects of the shape of distributions, but not to their absolute magnitude, we minimized the Kolmogorov–Smirnov ($K-S$) statistic (Sokal & Rohlf 1995) as a criterion of fit.

Separately for each target group of organisms, a multi-way analysis of variance (ANOVA) was carried out on the full set of simulation runs, with $K-S$ values as the response variable and 10 runs under

Figure 3. Visualization of results for a single realization of the model driven by the EPICA temperature series. (a) Cladograms for the 10 founders and their descendant species (each clade a different colour). Vertical location of clades in the figure is arbitrary. (b) Extinctions for each 20 kyr interval for the clades in (a). Note that more extinctions occur during cold phases than during interglacials. (c) The resampled EPICA temperature series (from figure 1b). (d) Time-elevation map of species richness for the clades in (a). Cool colours indicate lower richness, hot colours higher richness (0–100 species). (e) Time-elevation map of the phylogenetic composition of local assemblages. Hot colours indicate significant ($p < 0.025$, red) or nearly significant ($p < 0.05$, orange) phylogenetic clustering, whereas dark blue indicates phylogenetic mixing ('overdispersion'), although no assemblages quite reach $p > 0.975$ in this example. The 'average taxonomic distinctness' (Warwick & Clarke 1998) of assemblages was computed for each elevation and time, based on the cladogram in (a). Significance was assessed by a permutation test (500 permutations) for each cell, based on the pool of all living species at that time interval. See table 1 for parameters and settings for this replicate.

identical parameter settings as replicates, to determine which parameters were most important in distinguishing good fit from poor for each target pattern, given the parameter combinations considered. Each parameter (table 1) was treated as a factor and its settings as

levels. Standardized F -statistics were compared among parameters and target groups, but no significance tests were appropriate or performed for the ANOVA.

To identify the parameter value combinations that most accurately reproduced each target pattern, the

Figure 4. The signature of chance in time-elevation maps of species richness. All nine examples share exactly the same values of all settings and parameters except for the extinction parameter, α . The three examples within each column, however, are true replicates, sharing the same value of α as well as all other parameters. See table 1 for other parameters and settings.

Figure 5. Elevational range shifts as a function of niche conservatism/evolution. (a) The three time-elevation maps show representative richness patterns for three values of Δ , the niche centre evolution parameter (table 1). (b) The small graph below each richness map plots, for that example, the mean of the elevational range midpoints for all living species as a function of sea-level temperature, through the full course of the EPICA glacial cycles (figure 1). Strong niche conservatism ($\Delta = 0.1$) forces ranges to shift upslope during warming phases and downslope during cooling phases, whereas rapid niche evolution ($\Delta = 0.9$) permits ranges to remain relatively fixed on the elevation gradient regardless of temperature. In (c), the slopes of ordinary least-squares regressions (the lines in (b)) for 10 replicates at each of seven values of parameter Δ (spanning the range between 0.01 and 1.0) are plotted as a function of Δ , demonstrating a linear decrease in elevational range shift as niche conservatism weakens. See table 1 for other parameters and settings.

Table 2. Parameter space and model selection for Barva Transect richness simulation. (Parameter values that appeared in the most frequent parameter sets among the best-fitting 1% of replicates are indicated by x or **X**. Parameter sets for the examples in figure 7 are marked **X**. Entries in the ANOVA column indicate factor importance averaged across groups. Preliminary explorations having shown that the probability of speciation, as long as not zero or extremely high, matters little to the outcome, γ was fixed at 0.06 for all parameter sets. Model settings as for figure 3 (see table 1)).

parameter	ANOVA	levels	geometrid moths	epiphytes	Rubiaceae	ants
lateral area effect	0.375	off	X			
		on	x	X	X	X
number of founders	0.306	2				X
		10	X	x	X	x
		50		X		
founder niche breadth	0.033	1.00	x	x	x	X
		1.25	X	x	X	x
		1.50	x	X	x	x
		2.00	x			x
		2.50	x		x	x
		3.00	x	x		x
		3.50	x	x	x	
niche centre evolution	0.158	0.05				
		0.10				
		0.30				
		0.50				
		0.70	X	X	x	
		0.90	x	x	X	X
niche breadth evolution	0.375	0		x	x	x
		0.10	x	X	X	X
		0.15	X	x	x	x
		0.20	x			
		0.25				
		0.30				
extinction	0.128	0.01				
		0.05		X	x	x
		0.10	x	X	X	x
		0.15	X	X		X

model outcomes for all replicates (for all parameter values) were first sorted by the K–S statistic, for each target pattern separately. The best-fitting (smallest K–S) 1 per cent were selected for further analysis. Next, to help eliminate chance successes, this subset was searched for the most frequently represented parameter combinations (those closest to 10 replicates per combination) and the parameter settings for these sets were compared among the four target groups. No significance tests were appropriate or performed for the K–S statistics.

3. RESULTS

(a) *Visualization of patterns over time and elevation*

Each run (realization) of the model simulates the evolutionary, ecological and biogeographic fate of one or more founders and all descendant species over the 800 kyr of the EPICA temperature series (figure 1), under some particular fixed value of each of the model parameters and settings (table 1). Another run using exactly the same values of the parameters is considered a ‘replicate’ of this biogeographical experiment. We refer to groups of runs that share the same parameter values and model settings as ‘replicate sets’. Only by running the model repeatedly under the same parameter settings can we distinguish between

the roles of chance (stochasticity) and mechanism. Figure 3 illustrates, in detail, the results for a single replicate.

A key visualization tool for this study is the time versus elevation map, which plots the magnitude of richness for each discrete time step (160 for the EPICA series) and each elevational step (we used 100–140). For example, figure 3d plots local species richness, at each time and elevation, as a function of time (x -axis), and elevation (y -axis) under EPICA temperature cycles (figure 3c), for a single realization of the model. Species richness, which is simply the number of overlapping species ranges at a particular time and elevation, is indexed by colour according to the scale below the time-elevation map (figure 3d).

(b) *Stochastic variation in pattern among replicates*

The stochastic elements of the model (table 1) are limited to (i) initial location of founder niche centres (uniform random), (ii) the magnitude of initial niche breadths (normal distribution tuned by parameter σ), (iii) the niche breadths of child species (normally distributed differences from parent niche breadth, tuned by parameter β), (iv) the identity of species chosen to speciate (uniform random with probability γ), and (v) the identity of species chosen to undergo stochastic

extinction (weighted by range size and tuned by parameter α). All other processes and algorithms in the model are fully deterministic, given the prior settings (table 1) of the model. Nevertheless, temporal and spatial (elevational) variation in the pattern of species richness, range size, range location and local phylogenetic structure among replicates sharing the same parameter values proved to be remarkably rich.

Figure 4 illustrates the striking stochastic variation, among replicates within replicate sets, in species richness patterns as a function of time and elevation, for three levels of the extinction parameter α . The three replicates within each column share all parameter values. These examples show how the stochastic events of history become consolidated by speciation and the inheritance of niche characteristics.

(c) *An emergent pattern: mid-elevation richness peaks*

Despite the variation in spatial and temporal features of richness and ranges among replicates and between parameter settings, consistent patterns do emerge. The most striking is the tendency for richness to reach its maximum near mid-elevations, or at least to decline towards the boundaries of the domain, a pattern frequently seen along elevational transects in nature (Rahbek 2005).

When two or more founders initiate the simulation and a species ceiling is imposed, under most parameter settings subsequent biogeographic and phylogenetic history shows a pattern of domination by one or a very few clades, while others become extinct. The surviving clade or clades tend to be those that either began or (if parameters permit) have shifted towards mid-elevations, away from the extinction perils of the bounds of the domain. For example, in figure 3 (which is typical in this regard), of 10 founders, only two produced lineages that survived to the end of the simulation (figure 3a). Both started out at moderate elevations: 750 m and 1680 m on a 2900 m domain. The phylogenetic composition of local assemblages routinely shows the increasing dominance of the surviving clades as time passes and other clades become extinct. In the example, figure 3e plots the significance level of phylogenetic clustering or mixing in the assemblage at each elevation at each time. Following an early period of coexistence with other clades (see the cladogram in figure 3a), the two surviving clades come to dominate different elevations (red areas of figure 3e), briefly mixing at intermediate elevations (blue areas), eventually broadly overlapping (increasing yellow and green areas) to produce the final richness pattern. A formal, quantitative analysis of such repeated patterns is beyond the scope of the present study, but they strongly suggest lineage-level selection for biogeographic location on the domain.

Moreover, although the EPICA temperature fluctuations certainly strengthen this effect, we found that stochastic extinctions near the domain boundaries, where ranges tend to be smaller, can produce mid-elevation richness peaks even with a flat temperature series replacing EPICA (maintaining the usual lapse-based elevational temperature gradient). Rangel &

Diniz-Filho (2005; further discussed by Colwell & Rangel 2009) showed a similar result for a related, niche-based stochastic model. Previous stochastic models that incorporate speciation within a bounded domain have all yielded mid-domain peaks of richness (Bokma *et al.* 2001; Davies *et al.* 2005; Arita & Vazquez-Dominguez 2008; Connolly 2009). The differential extinction of small populations near domain boundaries was first proposed by Colwell & Hurtt (1994) as a biological explanation for the mid-domain effect (the tendency for larger ranges to overlap towards the middle of a bounded, spatial domain owing to geometric constraints on range location).

(d) *The effect of niche conservatism versus niche evolution on range shifts*

In the model, in the complete absence of anagenetic (within-species) niche evolution ($\Delta = 0$ and $\beta = 0$; table 1), species ranges are forced to shift downslope during cooling phases and upslope during warming phases in lockstep response to the constantly changing temperatures of the EPICA glacial–interglacial cycles. At the other extreme, if a species is capable of rapid evolutionary response to directional selection imposed by discordance between its niche optimum and the current mean temperature within its elevational range ($\Delta = 1$ and $\beta \geq 0$), its range centre remains fixed on the gradient. Figure 5 shows that the model yields a linear decrease in the magnitude of elevational range shift as niche conservatism weakens, for values of Δ spanning the range between 0.01 and 1.0, confirming that the model behaves as expected.

(e) *The effects of niche conservatism, temperature asymmetry and lateral area on extinction*

Extinctions are more likely with strong niche conservatism, for two reasons. First, if little or no adaptive evolution is possible, temperature change may leave a species without any region of the domain that corresponds to its thermal niche (deterministic extinction). Second, species with ranges abutting either boundary of the domain (sea level or mountaintop) may express only a portion of their fundamental niches, guaranteeing discordance between niche optimum and mean environmental conditions in the expressed range. If selection cannot correct this discordance, the range remains small, and is thus susceptible to stochastic extinction. At extremes of the glacial–interglacial cycles, both lowland and mountaintop species in the model are therefore more vulnerable to both kinds of extinction than species with mid-elevation ranges, especially so when niche conservatism is strong. However, two opposing, well-known asymmetries complicate this simple picture.

The first asymmetry lies in the long-term pattern of temperatures during the late Quaternary. For the resampled EPICA series (figure 1b), the temperature lies at or below the middle of its long-term range for 73 per cent of the 5 kyr intervals. At the extremes, the asymmetry is even more striking: 33 per cent of the time intervals lie within the coldest 20 per cent of the EPICA temperature range—the lingering glacial

Figure 6. The effect of lateral surface area on the elevational pattern of richness. The elevational centre of mass for species richness is plotted for 10 replicate simulations for each of four extinction intensities (parameter α , table 1). The extinction function (figure 2) was modelled for a hypothetical mountain shaped as a right cone, 5000 m high with a 5000 m base radius. Species elevational ranges were transformed to the corresponding area (r in the extinction function) of a band on the surface of the cone, as a proxy for relative population size, with the area of the full domain (v in the function) equal to the entire surface of the cone. With increasing extinction pressure, lower-elevation lineages are increasingly favoured by reason of their larger population sizes. The slope of the regression line, which passes the test for linearity, is significant at $p = 0.003$ (Sokal & Rohlf 1995, p. 476). See table 1 for other parameters and settings.

episodes, whereas only 6 per cent of intervals reach the warmest 20 per cent of the temperature range—the relatively brief spikes of the interglacials. Thus, even if extinctions are random across time intervals, far more extinctions would be expected to occur during glacial episodes than during interglacials. The pattern of extinction shown in figure 3*b* illustrates this asymmetry, which would be expected, over time, to impose stronger selection for cold tolerance among lowland lineages than for heat tolerance among mountaintop lineages.

The second asymmetry arises from the shape of mountain massifs, which generally (but not always) offer decreasing land surface area with increasing elevation, yielding smaller populations and thus greater vulnerability to stochastic extinction at higher elevations. We modelled the effect of lateral area on extinction by treating the domain as a cone, reckoning species ranges as the surface area of an elevational band, as a proxy for population size. The net result of the opposing effects on extinction of these two asymmetries is thus a quantitative matter, not predictable from first principles. Under the conditions of the model, figure 6 shows that increasing extinction pressure lowers the elevation of the centre of mass for species richness, evidently overcoming the contrary effect of greater numbers of extinctions in the lowlands during glacial maxima. Figure 4 illustrates a variety of examples from these simulations.

(f) *Target pattern matching for Barva Transect plants, moths and ants*

The empirical patterns of species richness with elevation for four groups of organisms on the Barva

Figure 7. Modelling target richness patterns for four groups of species on the Barva Transect, Costa Rica. The time-elevation maps illustrate the modelled temporal history of richness for each simulation, on a colour scale from 0 (blue) to 100 (red) species. The blue curves on the right show the smoothed empirical profile of richness with elevation for four groups of species, and the red curves show the modelled richness profile at the end (right edge of the richness map) of the simulation, on a relative scale. See the text for methods and table 2 for parameter settings for each group. (a) geometrid moths, (b) epiphytes, (c) Rubiaceae and (d) ants.

Transect (blue profiles in the graphs on the right of figure 7) vary greatly in shape and in the elevation at which species richness peaks, but all four richness patterns can be reasonably well reproduced from the model (red profiles in figure 7). Not surprisingly, given the role of chance in the richness patterns produced by the model (as illustrated in figure 4 for a different analysis), even within a particular replicate set, some replicates fit better than others. Nevertheless, patterns of success and failure do emerge, and the range of parameter settings that yield the best fit differs among the four groups for some parameters (table 2), but not for others. We endeavoured to minimize the hazard of finding one-off fits by considering only parameter sets that repeatedly produced a good fit to the empirical patterns (based on the K–S statistic). Moreover, the examples illustrated in figure 7 were not selected from the realizations of the model used to define ‘best sets’ of parameter values,

but were produced subsequently by independent runs based on one or more of those sets. In effect, the procedure can be viewed as an informal test of the hypothesis that the empirical profiles could have been drawn from the distribution of richness patterns produced by the model.

As table 2 shows, a surprisingly high level of the niche centre evolution parameter, Δ (0.7 or 0.9, where 1.0 represents full adaptive evolution at each time step), proved to be crucial for a good fit of modelled to empirical richness for all groups. By contrast, a good fit called for relatively low levels of the niche breadth evolution parameter, β , although a somewhat higher level of this parameter proved optimal for the broad, centred distribution of the moths. High rates of extinction proved to be optimal for all groups, perhaps serving to enhance the rate of evolution of niche centres under the constraint of the species ceiling (K). Groups differed in the optimal level of founder number (few for the low-elevation ants, many for the broad, convex richness pattern of the epiphytes) and lateral area effect (which tends to produce lower elevation richness peaks, figure 6). The levels for speciation rate (γ , results not shown) and founder niche breadth (σ) proved to matter little, as long as not zero.

4. DISCUSSION

Integrating determinism with chance and ecology with evolution, we have attempted to model a brief moment in the history of the Earth for a simplified biogeographical gradient, based on a single environmental factor. By simplifying both niche space and geographical space, we were able to model and visualize environmental effects on species distributions on a geological time scale (figure 3), taking 'general simulation models' in macroecology in a new direction (Gotelli *et al.* 2009). Although we incorporated lateral extent into the extinction function by taking elevational band area into account, a more explicit accounting for 'x, y space' (where elevation is the z dimension) would clearly be desirable. As for the environmental side of the model, the most significant limitation of our model is its lack of accounting for patterns of precipitation. While no species can escape its limits of thermal tolerance, those limits are of course often modified by water availability, particularly for plants.

To us, the most surprising result was not that the constraints and mechanisms built into the model succeeded in producing reasonable patterns of phylogeny and macroecology, but the remarkably rich variation among replicate realizations under identical parameters, as speciation consolidates the differential survival of lineages and inherited ecological tolerances perpetuate their location in space (figure 3). This result, which is unlikely to be changed in a more complete model, suggests that the search for mechanistic explanations for macroecological patterns may inevitably be constrained by the idiosyncratic nature of historical events, perhaps to a greater degree than many of us have appreciated.

Our attempt to reproduce the empirical elevational richness profiles of four groups of organisms on the Barva Transect within the parameter space of the

model (figure 7 and table 2) is no more than an exploratory first step towards mechanistic modelling of empirical patterns on tropical elevational gradients. Ideally, parameter-set selection should target multiple characteristics (Grimm *et al.* 2005), including range-size frequency distributions, diversity and dispersion fields, characteristics of the phylogeny and the phylogenetic structure of local assemblages (Gotelli *et al.* 2009).

Consistently high rates of niche centre evolution combined with low rates of niche breadth evolution were required for a good fit of modelled to empirical richness profiles, together confining each founder clade to a relatively stable range of elevations, in spite of the repeated temperature cycles. Thus, two founders were more likely to produce a good fit for ants (a single insect family), with their relatively narrow lowland richness peak, whereas 50 founders did a better job for epiphytes (50 plant families; Cardelús *et al.* 2006), with their broad, convex richness profile. Clearly, more than two lineages of ants and more than 50 lineages of epiphytes have contributed to the contemporary biotas of the Costa Rican Atlantic slope, but the qualitative difference in phylogenetic, niche and functional diversity may be realistic.

Our results suggest that niche conservatism for temperature optimum over the time scale of the Quaternary might not be a winning strategy, contrary to other views (e.g. Martinez-Meyer *et al.* 2004; Kozak & Wiens 2010). Experimental and genetic studies of tropical montane species may reveal unexpected levels of intraspecific and intraclade genetic variation within and between elevations (Gilman 2007), just as within-species adaptation to local conditions may play a substantial role in latitudinal range shifts (Davis & Shaw 2001). In fact, although we have consistently referred to the taxonomic units of modelled clades (figure 2), as 'species', we recognize that speciation in most groups of organisms has not proceeded at such a rapid pace within the relatively short span of the Quaternary, although there are exceptions (Richardson *et al.* 2001; Hooghiemstra & Van der Hammen 2004; Kay *et al.* 2005; Fjeldså & Rahbek 2006). However, the same units may instead be viewed as clades at an infraspecific level, with the caveat that, in the model, lineages do not rejoin.

The opposing effects on lowland extinctions of temporal asymmetry in temperature (greater duration of glacial than of interglacial episodes) and spatial asymmetry in lateral area (greater land area at low than at high elevations) have apparently not previously been recognized. Our results, for a hypothetical conical mountain (of a particular height and slope; figure 6), indicate that area comes to the rescue, yielding richness peaks below mid-elevation, but of course that result may be model-dependent and is certainly dependent on the actual elevational profile of particular mountain massifs. We conjecture that, in general, the steeper and more isolated the mountain, the weaker the area effect in countering lowland extinctions during glacial episodes, forcing the richness profile upslope, compared with patterns on broader mountain massifs. In any case, in terms of selection for traits that allow species to tolerate extreme temperatures, the inescapable inference appears to be that

the Quaternary has imposed far more continuous selection for cold-tolerance than for heat-tolerance, at all elevations (Colwell *et al.* 2008). If, indeed, tropical species are better adapted to glacial cold than to interglacial heat, these opposing asymmetries might help explain the 'out of the tropics' (Jablonski *et al.* 2006) Pleistocene expansion of certain lineages towards higher latitudes. Meanwhile, this inference is not good news for tropical species under the current anthropogenic global warming trend, which comes at an unfortunate point in the orbitally forced cycles of the Quaternary.

The comments of R. Burnham, D. Nogués-Bravo, G. Brehm, J. Fjeldsø, C. Rahbek and two anonymous reviewers greatly improved the manuscript. We are grateful to G. Brehm, C. Cardelús, A. Gilman and J. Longino for sharing their Barva Transect data. We acknowledge the Danish National Research Foundation for support to the Center for Macroecology, Evolution and Climate, University of Copenhagen (Denmark), the National Science Foundation (USA, DEB-0639979 and DBI-0851245) and the National Center for Ecological Analysis and Synthesis (NCEAS, USA). T.F.R. was supported by the Department of Ecology and Evolutionary Biology, University of Connecticut (USA), the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior/Fulbright Fellowship (CAPES, Brazil); and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Brazil).

REFERENCES

- Ahrens, C. D. 2006 *Meteorology today*, 8th edn. Brooks/Cole Publishing.
- Alley, R. B. *et al.* 2003 Abrupt climate change. *Science* **299**, 2005. (doi:10.1126/science.1081056)
- Arita, H. & Vazquez-Dominguez, E. 2008 The tropics: cradle, museum or casino? A dynamic null model for latitudinal gradients of species diversity. *Ecol. Lett.* **11**, 653–663. (doi:10.1111/j.1461-0248.2008.01197.x)
- Ashcroft, M. 2010 Identifying refugia from climate change. *J. Biogeogr.* **37**, 1407–1413. (doi:10.1111/j.1365-2699.2010.02300.x)
- Blois, J. L., McGuire, J. L. & Hadly, E. A. 2010 Small mammal diversity loss in response to late-Pleistocene climatic change. *Nature* **465**, 771–774. (doi:10.1038/nature09077)
- Bokma, F., Bokma, J. & Mönkkönen, M. 2001 Random processes and geographic species richness patterns: why so few species in the north. *Ecography* **24**, 43–49. (doi:10.1034/j.1600-0587.2001.240106.x)
- Brehm, G., Colwell, R. K. & Kluge, J. 2007 The role of environment and mid-domain effect on moth species richness along a tropical elevational gradient. *Glob. Ecol. Biogeogr.* **16**, 205–219.
- Bush, M. B. & Flenley, J. R. 2007 *Tropical rainforest responses to climatic change*. Berlin, Germany: Springer.
- Bush, M. B. & Silman, M. R. 2004 Observations on Late Pleistocene cooling and precipitation in the lowland Neotropics. *J. Quaternary Sci.* **19**, 677–684. (doi:10.1002/jqs.883)
- Bush, M. B., Silman, M. R. & Urrego, D. H. 2004 48 000 years of climate and forest change in a biodiversity hot spot. *Science* **303**, 827–829. (doi:10.1126/science.1090795)
- Cardelús, C. L., Colwell, R. K. & Watkins, J. E. 2006 Vascular epiphyte distribution patterns: explaining the mid-elevation richness peak. *J. Ecol.* **94**, 144–156. (doi:10.1111/j.1365-2745.2005.01052.x)
- Chen, I.-C., Shiu, H.-J., Benedick, S., Holloway, J. D., Chey, V. K., Barlow, H. S., Hill, J. K. & Thomas, C. D. 2009 Elevation increases in moth assemblages over 42 years on a tropical mountain. *Proc. Natl Acad. Sci. USA* **106**, 1479–1483. (doi:10.1073/pnas.0809320106)
- Clark, D. A., Piper, S. C., Keeling, C. D. & Clark, D. B. 2003 Tropical rain forest tree growth and atmospheric carbon dynamics linked to interannual temperature variation during 1984–2000. *Proc. Natl Acad. Sci. USA* **100**, 5852–5857. (doi:10.1073/pnas.0935903100)
- Colinvaux, P., De Oliveira, P., Moreno, J., Miller, M. & Bush, M. 1996 A long pollen record from lowland Amazonia: forest and cooling in glacial times. *Science* **274**, 85. (doi:10.1126/science.274.5284.85)
- Colwell, R. K. & Hurtt, G. C. 1994 Nonbiological gradients in species richness and a spurious Rapoport effect. *Am. Nat.* **144**, 570–595.
- Colwell, R. K. & Lees, D. C. 2000 The mid-domain effect: geometric constraints on the geography of species richness. *Trends Ecol. Evol.* **15**, 70–76. (doi:10.1016/S0169-5347(99)01767-X)
- Colwell, R. K. & Rangel, T. F. 2009 Hutchinson's duality: the once and future niche. *Proc. Natl Acad. Sci. USA* **106**, 19 651–19 658. (doi:10.1073/pnas.0901650106)
- Colwell, R. K., Brehm, G., Cardelús, C., Gilman, A. C. & Longino, J. T. 2008 Global warming, elevational range shifts, and lowland biotic attrition in the wet tropics. *Science* **322**, 258–261. (doi:10.1126/science.1162547)
- Connolly, S. R. 2009 Macroecological theory and the analysis of species richness gradients. In *Marine macroecology* (eds J. D. Witman & K. Roy), pp. 279–309. Chicago, IL: University of Chicago Press.
- Davies, T. J., Grenyer, R. & Gittleman, J. L. 2005 Phylogeny can make the mid-domain effect an inappropriate null model. *Biol. Lett.* **1**, 143–146. (doi:10.1098/rsbl.2005.0297)
- Davis, M. B. & Shaw, R. G. 2001 Range shifts and adaptive responses to Quaternary climate change. *Science* **292**, 673. (doi:10.1126/science.292.5517.673)
- Davis, M. B., Shaw, R. G. & Etterson, J. R. 2005 Evolutionary responses to changing climate. *Ecology* **86**, 1704–1714. (doi:10.1890/03-0788)
- Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C. & Martin, P. R. 2008 Impacts of climate warming on terrestrial ectotherms across latitude. *Proc. Natl Acad. Sci. USA* **105**, 6668–6672. (doi:10.1073/pnas.0709472105)
- Dick, C. W. & Heuertz, M. 2008 The complex biogeographic history of a widespread tropical tree species. *Evolution* **62**, 2760–2774. (doi:10.1111/j.1558-5646.2008.00506.x)
- Feeley, K. J. & Silman, M. R. 2010 Biotic attrition from the hot tropics correcting for truncated temperature niches. *Glob. Change Biol.* **16**, 1830–1836. (doi:10.1111/j.1365-2486.2009.02085.x)
- Feeley, K. J., Joseph Wright, S., Nur Supardi, M. N., Kassim, A. R. & Davies, S. J. 2007 Decelerating growth in tropical forest trees. *Ecol. Lett.* **10**, 461–469. (doi:10.1111/j.1461-0248.2007.01033.x)
- Fjeldsø, J. & Rahbek, C. 2006 Diversification of tanagers, a species rich bird group, from lowlands to montane regions in South America. *Integr. Comp. Zool.* **46**, 72–78. (doi:10.1093/icb/icj009)
- Flenley, J. R. 1998 Tropical forests under the climates of the last 30 000 years. *Clim. Change* **39**, 177–197. (doi:10.1023/A:1005367822750)
- Garzone, C. N., Hoke, G. D., Libarkin, J. C., Withers, S., MacFadden, B., Eiler, J., Ghosh, P. & Mulch, A. 2008 Rise of the Andes. *Science* **320**, 1304. (doi:10.1126/science.1148615)

- Gilman, A. 2007 *Tropical plant diversity and species altitudinal range across an elevational gradient in Costa Rica*. Los Angeles, CA: Department of Ecology and Evolutionary Biology, UCLA.
- Gotelli, N. *et al.* 2009 Patterns and causes of species richness: a general simulation model for macroecology. *Ecol. Lett.* **12**, 873–886. (doi:10.1111/j.1461-0248.2009.01353.x)
- Grimm, V. *et al.* 2005 Pattern-oriented modeling of agent-based complex systems: lessons from ecology. *Science* **310**, 987. (doi:10.1126/science.1116681)
- Hansen, J., Sato, M., Ruedy, R., Lo, K., Lea, D. W. & Medina-Elizade, M. 2006 Global temperature change. *Proc. Natl Acad. Sci. USA* **103**, 14 288–14 293. (doi:10.1073/pnas.0606291103)
- Hickling, R., Roy, D. B., Hill, J. K., Fox, R. & Thomas, C. D. 2006 The distributions of a wide range of taxonomic groups are expanding polewards. *Glob. Change Biol.* **12**, 450–455. (doi:10.1111/j.1365-2486.2006.01116.x)
- Hooghiemstra, H. & Van der Hammen, T. 2004 Quaternary Ice-Age dynamics in the Colombian Andes: developing an understanding of our legacy. *Phil. Trans. R. Soc. B* **359**, 173. (doi:10.1098/rstb.2003.1420)
- Jablonski, D., Roy, K. & Valentine, J. W. 2006 Out of the tropics: evolutionary dynamics of the latitudinal diversity gradient. *Science* **314**, 102. (doi:10.1126/science.1130880)
- Jansson, R. & Dynesius, M. 2002 The fate of clades in a world of recurrent climate change: Milankovitch oscillations and evolution. *Ann. Rev. Ecol. Syst.* **33**, 741–777. (doi:10.1146/annurev.ecolsys.33.010802.150520)
- Jouzel, J. *et al.* 2007 Orbital and millennial Antarctic climate variability over the past 800,000 years. *Science* **317**, 793–797. (doi:10.1126/science.1141038)
- Kay, K. M., Reeves, P. A., Olmstead, R. G. & Schemske, D. W. 2005 Rapid speciation and the evolution of hummingbird pollination in Neotropical *Costus* subgenus *Costus* (Costaceae): evidence from nrDNA ITS and ETS sequences. *Am. J. Bot.* **92**, 1899. (doi:10.3732/ajb.92.11.1899)
- Kirkpatrick, M. & Barton, N. H. 1997 Evolution of a species' range. *Am. Nat.* **150**, 1–23.
- Kozak, K. & Wiens, J. 2010 Niche conservatism drives elevational diversity patterns in Appalachian salamanders. *Am. Nat.* **176**.
- Liu, Z., Cleaveland, L. & Herbert, T. 2008 Early onset and origin of 100-kyr cycles in Pleistocene tropical SST records. *Earth Planet. Sci. Lett.* **265**, 703–715. (doi:10.1016/j.epsl.2007.11.016)
- Longino, J., Coddington, J. A. & Colwell, R. K. 2002 The ant fauna of a tropical rainforest: estimating species richness three different ways. *Ecology* **83**, 689–702. (doi:10.1890/0012-9658(2002)083[0689:TAFOAT]2.0.CO;2)
- Losos, J. 2008 Phylogenetic niche conservatism, phylogenetic signal and the relationship between phylogenetic relatedness and ecological similarity among species. *Ecol. Lett.* **11**, 995–1003. (doi:10.1111/j.1461-0248.2008.01229.x)
- MacFadden, B. J., Cerling, T. E., Harris, J. M. & Prado, J. 1999 Ancient latitudinal gradients of C3/C4 grasses interpreted from stable isotopes of New World Pleistocene horse (*Equus*) teeth. *Glob. Ecol. Biogeogr.* **8**, 137–149.
- Marchant, R., Boom, A. & Hooghiemstra, H. 2002 Pollen-based biome reconstructions for the past 450 000 yr from the Funza-2 core, Colombia: comparisons with model-based vegetation reconstructions. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **177**, 29–45. (doi:10.1016/S0031-0182(01)00350-9)
- Marchant, R. *et al.* 2009 Pollen-based biome reconstructions for Latin America at 0, 6000 and 18 000 radiocarbon years ago. *Clim. Past* **5**, 725–767. (doi:10.5194/cp-5-725-2009)
- Martinez-Meyer, E., Peterson, A. T. & Hargrove, W. W. 2004 Ecological niches as stable distributional constraints on mammal species, with implications for Pleistocene extinctions and climate change projections for biodiversity. *Glob. Ecol. Biogeogr.* **13**, 305–314. (doi:10.1111/j.1466-822X.2004.00107.x)
- Moritz, C., Patton, J., Conroy, C., Parra, J., White, G. & Beissinger, S. 2008 Impact of a century of climate change on small-mammal communities in Yosemite National Park, USA. *Science* **322**, 261. (doi:10.1126/science.1163428)
- Purvis, A., Gittleman, J. L., Cowlishaw, G. & Mace, G. M. 2000 Predicting extinction risk in declining species. *Proc. R. Soc. Lond. B* **267**, 1947. (doi:10.1098/rspb.2000.1234)
- Rahbek, C. 2005 The role of spatial scale in the perception of large-scale species-richness patterns. *Ecol. Lett.* **8**, 224–239. (doi:10.1111/j.1461-0248.2004.00701.x)
- Rangel, T. F. L. V. B. & Diniz-Filho, J. A. F. 2005 An evolutionary tolerance model explaining spatial patterns in species richness under environmental gradients and geometric constraints. *Ecography* **28**, 253–263. (doi:10.1111/j.0906-7590.2005.04038.x)
- Rangel, T. F. L. V. B., Diniz-Filho, J. A. F. & Colwell, R. K. 2007 Species richness and evolutionary niche dynamics: a spatial pattern-oriented simulation experiment. *Am. Nat.* **170**, 602–616. (doi:10.1086/521315)
- Rangel, T. F., Diniz-Filho, J. A. F. & Bini, L. M. 2010 SAM: a comprehensive application for spatial analysis in macroecology. *Ecography* **33**, 46–50. (doi:10.1111/j.1600-0587.2009.06299.x)
- Richardson, J. E., Pennington, R. T., Pennington, T. D. & Hollingsworth, P. M. 2001 Rapid diversification of a species-rich genus of Neotropical rain forest trees. *Science* **293**, 2242. (doi:10.1126/science.1061421)
- Rutherford, S. & D'Hondt, S. 2000 Early onset and tropical forcing of 100 000-year Pleistocene glacial cycles. *Nature* **408**, 72–75. (doi:10.1038/35040533)
- Schluter, D. & Rambaut, A. 1996 Ecological speciation in postglacial fishes. *Phil. Trans. R. Soc. Lond. B* **351**, 807–814. (doi:10.1098/rstb.1996.0075)
- Sinervo, B. *et al.* 2010 Erosion of lizard diversity by climate change and altered thermal niches. *Science* **328**, 894. (doi:10.1126/science.1184695)
- Smith, F. A., Betancourt, J. L. & Brown, J. H. 1995 Evolution of body size in the woodrat over the past 25 000 years of climate change. *Science* **270**, 2012–2014. (doi:10.1126/science.270.5244.2012)
- Sokal, R. & Rohlf, F. 1995 *Biometry: the principles and practice of statistics in biological research*. New York, NY: WH Freeman.
- Voelker, G. 2010 Repeated vicariance of Eurasian songbird lineages since the Late Miocene. *J. Biogeogr.* **31**, 1251–1261. (doi:10.1111/j.1365-2699.2010.02313.x)
- Warwick, R. M. & Clarke, K. R. 1998 Taxonomic distinctness and environmental assessment. *J. Appl. Ecol.* **35**, 532–543. (doi:10.1046/j.1365-2664.1998.3540532.x)
- Wille, M., Hooghiemstra, H., Behling, H., van der Borg, K. & Negret, A. J. 2001 Environmental change in the Colombian subandean forest belt from 8 pollen records: the last 50 kyr. *Veg. Hist. Archaeobot.* **10**, 61–77. (doi:10.1007/PL00006921)
- Williams, J. W., Jackson, S. T. & Kutzbach, J. E. 2007 Projected distributions of novel and disappearing climates by 2100 AD. *Proc. Natl Acad. Sci. USA* **104**, 5738–5742. (doi:10.1073/pnas.0606292104)