Published in final edited form as:

Toxicol Appl Pharmacol. 2010 August 1; 246(3): 188. doi:10.1016/j.taap.2010.05.001.

Corrigendum to “Immunotoxicity and biodistribution analysis of arsenic trioxide in C57B1/6 mice following a 2-week inhalation exposure”:

[*Toxicol. Appl. Pharmacol.* 241 (2009) 253–259]

Scott W. Burchiel^{a,*}, Leah A. Mitchell^a, Fredine T. Lauer^a, Xi Sun^a, Jacob D. McDonald^b, Laurie G. Hudson^a, and Ke Jian Liu^a

^a Toxicology and Pharmaceutical Sciences Program, The University of New Mexico College of Pharmacy, Albuquerque, NM 87131, USA

^b The Lovelace Respiratory Research Institute, Albuquerque, NM 87109, USA

On page 255, Fig. 1, the concentration of As detected in tissues and blood should reflect ng/g. The correct figure is shown here.

Acknowledgments

The authors wish to include the following acknowledgment: This work was supported in part by supplements to the University of New Mexico NIEHS P30-ES012093 and to the University of Arizona P30-ES006694 Core Research Centers.

---

*Corresponding author. Sburchiel@salud.unm.edu (S.W. Burchiel).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Figure 1.

Tissue analysis of total arsenic in tissues by ICP/MS immediately following the last nose-only inhalation exposure. There were five mice per group that received air only (Control), 50 µg/m³ ATO for 14 days (3 h per day). Results are expressed as the average level of arsenic (ng/g tissue) ± standard deviation.