Published in final edited form as:

Lancet. 1973 May 19; 1(7812): 1082–1085.

SURVIVAL OF A HOMOLOGOUS PARATHYROID IMPLANT IN AN IMMUNOSUPPRESSED PATIENT

Carl G. Groth, William S. Hammond, Shunzaburo Iwatsuki, Mordecai Popovtzer, Sergio Cascardo, Charles G. Halgrimson, and Thomas E. Starzl

Departments of Surgery, Medicine, and Pathology, University of Colorado School of Medicine, and Veterans Administration Hospital, Denver, Colorado, U.S.A

Summary

A patient who had undergone subtotal parathyroidectomy while on chronic haemodialysis became severely hypocalcaemic after receiving a well-functioning cadaveric renal graft. After a homologous parathyroid implant, there was a biochemical improvement in the patient, and at biopsy 6 months later the parathyroid graft was histologically normal. 21 months after the implant, serum calcium and phosphorus remain normal without calcium supplementation.

Introduction

About a hundred attempts at homologous parathyroid transplantation in man have been reported,^{1–3} but with three possible exceptions there has been no histological proof of graft survival. Two of the three reports were published 50 years ago, and inconclusively described “remnants of three small glands which, although degenerated, had the histologic appearance of parathyroid tissues”⁴ and “a nodule containing an array of aligned cells rather unusual in appearance, with an architecture revealing epithelial cell-like structures resembling parathyroid”.⁵ In the third instance, histologically intact homologous parathyroid tissue was found in a millipore chamber that had been implanted for 5 months, but the patient had demonstrated no biochemical or clinical improvement.⁶

We report the morphological survival and apparent functioning of a homologous parathyroid implant in a patient who, unlike the earlier parathyroid-transplant recipients, was immunosuppressed.

Case-report

A 46-year-old male with polycystic kidney disease had been on chronic haemodialysis since 1966. In 1969, skeletal pain, spontaneous rib fractures, and raised serum-calcium levels (11.5–12.2 mg. per 100 ml.) developed. Exploration of the neck in February, 1970, revealed parathyroid hyperplasia, and three and a half glands were removed. Postoperatively, serum-calcium was within normal limits and the bone disease improved.

After cadaveric renal transplantation at the Denver Veterans Administration Hospital on May 28, 1971, renal function became normal and rejection was never diagnosed (fig. 1). Postoperatively the patient was immunosuppressed with cyclophosphamide or azathioprine, prednisone, and antilymphocyte globulin (fig. 1). Other medications included reserpine, hydralazine hydrochloride, and chlorothiazide. The first 2 postoperative months were

complicated by recurrent convulsive seizures, dysphasia, and dyskinesia. Neurological examination only showed arteriosclerotic vessels by cerebral angiography. Subsequently, the patient's neurological condition improved slowly and at present only moderate dysphasia remains.

Immediately after the renal transplantation, serum-calcium dropped to subnormal levels (6–7.5 mg. per 100 ml.). Oral supplementation with calcium did not bring the serum levels to normal and calcium had to be administered intravenously (fig. 1). During this time, tubular reabsorption of phosphorus was 75 to 95%.

52 days after renal transplantation, four separate pieces of homologous hyperplastic parathyroid tissue (approximately 150 mg.) were implanted into the patient's pectoralis muscle. The grafts were obtained from another renal-transplant recipient undergoing subtotal parathyroidectomy. The donor was blood-group A, while the recipient was of blood-group O.

Initially, the recipient's serum-calcium remained below normal, necessitating further supplementation with calcium and vitamin D₂ (fig. 1). But 2 months after implantation, all supplementation could be stopped. In January, 1972, a biopsy specimen was taken from the site of the parathyroid grafting. This was followed by a transient drop in serum-calcium, and supplementary treatment was reinstituted for a few days (fig. 1). In the ensuing 12 months, serum calcium and phosphorus have been within normal limits.

Results

The parathyroid tissue used for implantation consisted microscopically of sheets of enlarged chief cells with occasional acini, characteristic of secondary hyperplasia. On biopsy 6 months after implantation, cords and clusters of chief cells were set in a fine, highly vascularised stroma characteristic of normal parathyroid tissue (fig. 2). Occasional interstitial collections of lymphocytes and plasma-cells suggested slight homograft rejection. Electron microscopy was performed on tissue which had been embedded in a paraffin block and was thus poorly preserved ultra-structurally. However, large amounts of glycogen, occasional prominent arrays of rough endoplasmic reticulum, and secretion granules characteristic of functioning parathyroid cells⁷ were apparent (fig. 3).

Discussion

The early assumption that homologous endocrine implants enjoy a major immunological privilege has never been substantiated. On the contrary, several experimental studies have shown that rejection of thyroid-parathyroid grafts does occur.^{8–10}

Parathyroid grafting during immunosuppression has not previously been attempted in man. Our patient was receiving immunosuppressive drugs after renal transplantation. Immunosuppression was highly effective, as judged by the uninterrupted and excellent functioning of the kidney graft thereafter. Consequently, it was not surprising that there was only slight evidence of rejection in the biopsy specimen of the parathyroid graft obtained after 6 months.

There was strong biochemical evidence that hypoparathyroidism had developed after subtotal parathyroidectomy. When the artificial metabolic conditions of chronic haemodialysis were followed by the normal renal function after transplantation, serum-calcium fell to extremely low levels. At the same time, the fractional tubular reabsorption of phosphorus far exceeded that usually recorded in patients with good graft function who are being treated with steroids and thiazides.¹¹

Recovery from surgically caused hypoparathyroidism has been reported after temporary administration of vitamin D and may even occur “spontaneously”.¹² However, it seems more likely that in the present patient the restoration of normal calcium homeostasis was the consequence of hormone excretion from the graft. Indirect evidence for graft function was obtained from the electron-microscopic pictures of the biopsy specimen. The delay, in effect, could be attributed to the time required for revascularisation and healing of the free, grafted tissue. Efforts to measure parathormone levels in the stored sera from our patient gave results which were not reproducible and were consequently not as useful for following the postoperative course as were other serum levels which measured parathyroid function.

Better techniques for the measurement of parathormone will be invaluable for assessing parathyroid-homograft function. Parathyroid transplantation will probably be indicated in other recipients of renal homografts, since the removal of hyperplastic parathyroid glands to control bone disease is becoming more common. An accident to the residual glandular tissue could lead to the situation which developed in our recipient. Ironically, the best opportunities for obtaining a suitable homograft for such a patient will be provided by other members of the kidney-transplant population.

Acknowledgments

This work was supported by research grants from the Veterans Administration, by grants RR-00051 and RR-00069 from the National Institutes of Health, and by grants AI-AM-08898, AM-07772, and HE-09110 from the United States Public Health Service.

References

1. Jacob SW, Dunphy JE. Am J Surg 1963;105:196. [PubMed: 13964291]
2. Jordan GL Jr. Adv Surg 1966;2:199. [PubMed: 5928809]
3. Brooks, JR. Human Organ Support and Replacement: Transplantation and Artificial Prostheses. Hardy, JD., editor. Vol. 176. Springfield; Illinois: 1971.
4. Brown AE. Ann Surg 1922;75:418. [PubMed: 17864618]
5. Feugoni C, Scimone V, Comolli A. Presse méd 1926;34:355.
6. Brooks JR, Hill GJ. Am J Surg 1960;99:588. [PubMed: 13804785]
7. Caper CC. Am J Med 1971;50:598. [PubMed: 4930268]
8. Fisher B, Lee S, Fisher ER, Sakai A, Rabinovitz M. Archs Surg 1964;89:65.
9. Fisher B, Fisher ER, Feduska N, Sakai A. Surgery, St Louis 1967;62:1025.
10. Lance EM. Transplantation 1967;5:1471.
11. Popovtzer M. Unpublished.
12. Watkins E Jr, Dell GO, Snow JC, Adams HD. J Am med Ass 1962;182:140.

Fig. 1. Course and treatment after renal transplantation and subsequent parathyroid implantation. Vertical arrows indicate intravenous infusion of 540 mg. calcium gluconate.

Fig. 2. Parathyroid tissue obtained at biopsy 6 months after homologous implantation
The glandular tissue is surrounded by skeletal muscle. Area shown is focally infiltrated by lymphocytes and plasma-cells (arrows); most of graft was without such cells. (Haematoxylin and eosin. $\times 100$.)

Fig. 3. Parathyroid tissue obtained at biopsy 6 months after homologous implantation
Granular endoplasmic reticulum (ER), glycogen (G), and secretory granules (arrows) are shown. ($\times 21,000$.)