Published in final edited form as:

J Allergy Clin Immunol. 2010 December ; 126(6): 1131–1138. doi:10.1016/j.jaci.2010.08.002.

Factors predicting inhaled corticosteroid responsiveness in African American patients with asthma

Wendy Gould, MD^{(1),*}, Edward L. Peterson, PhD^{(2),*}, Gloria Karungi, MBChB, MS⁽³⁾, Amanda Zoratti, BA⁽³⁾, John Gaggin, BSN⁽³⁾, Ghazwan Toma, MD, MPH⁽³⁾, Shiqing Yan, MD⁽³⁾, Albert M. Levin, PhD⁽²⁾, James J. Yang, PhD⁽²⁾, Karen Wells, BS⁽²⁾, Mingqun Wang, MS⁽³⁾, Robert R. Burke, MD, MS⁽¹⁾, Kenneth Beckman, PhD⁽⁴⁾, Danijela Popadic, BS⁽⁵⁾, Susan J. Land, PhD⁽⁵⁾, Rajesh Kumar, MD⁽⁶⁾, Max A. Seibold, PhD⁽⁷⁾, David E. Lanfear, MD, MS^{(1),(3)}, Esteban G. Burchard, MD, MPH^{(8),(9)}, and L. Keoki Williams, MD, MPH^{(1),(2),(3)}

⁽¹⁾Department of Internal Medicine, Henry Ford Health System, 1 Ford Place, Detroit, Michigan 48202

⁽²⁾Department of Biostatistics and Research Epidemiology, Henry Ford Health System, 1 Ford Place, Detroit, Michigan 48202

⁽³⁾Center for Health Services Research, Henry Ford Health System, 1 Ford Place, Detroit, Michigan 48202

⁽⁴⁾Biomedical Genomics Center, University of Minnesota, 426 Church Street SE, Minneapolis, MN 55455

⁽⁵⁾Applied Genomics Technology Center, Wayne State University, 275 East Hancock, Detroit, MI 48201

⁽⁶⁾Children's Memorial Hospital, Department of Pediatrics, Northwestern University Feinberg School of Medicine, 2300 Childrens' Plaza, Chicago, Illinois, 60614

⁽⁷⁾Department of Medicine, National Jewish Medical and Research Center, 1400 Jackson St., Denver, Colorado 80206

⁽⁸⁾Department of Medicine, University of California San Francisco, San Francisco General Hospital, Box 2911, San Francisco, California 94143

⁽⁹⁾Department of Bioengineering and Therapeutic Sciences, University of California San Francisco, San Francisco, California 94143

Abstract

Background—African American patients suffer disproportionately from uncontrolled asthma. Treatment with an inhaled corticosteroid (ICS) is considered first-line therapy for persistent asthma.

© 2010 American Academy of Allergy, Asthma and Immunology. Published by Mosby, Inc. All rights reserved.

Corresponding Author L. Keoki Williams, MD, MPH Center for Health Services Research Henry Ford Health System 1 Ford Place, 3A Detroit, Michigan 48202 (313)874-6693 (office) (313)874-7137 (fax) kwillia5@hfhs.org.

*These individuals contributed equally to the manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Objective—To determine the degree to which African American patients respond to ICS medication and whether the level of response is influenced by other factors, including genetic ancestry.

Methods—Patients aged 12-56 years who received care from a large health system in southeast Michigan and who resided in Detroit were recruited to participate if they had a diagnosis of asthma. Patients were treated with 6 weeks of inhaled beclomethasone dipropionate, and pulmonary function was re-measured after treatment. Ancestry was determined by genotyping ancestry informative markers. The main outcome measure was ICS responsiveness defined as the change in pre-bronchodilator FEV₁ over the 6-week course of treatment.

Results—Among 147 participating African American patients with asthma, average improvement in FEV₁ following 6 weeks of ICS treatment was 11.6%. The mean proportion of African ancestry in this group was 78.4%. The degree of baseline bronchodilator reversibility was the only factor consistently associated ICS responsiveness as measured by both an improvement in FEV₁ and in patient reported asthma control (P=0.001 and P=0.021, respectively). The proportion of African ancestry was not significantly associated with ICS responsiveness.

Conclusions—While baseline pulmonary function parameters appear to be associated with the likelihood to respond to ICS treatment, the proportion of genetic African ancestry does not. This study suggests that genetic ancestry may not contribute to differences in ICS controller response among African American patients with asthma.

Clinical Implications—Although African American patients suffer disproportionately from asthma-related complications, response to ICS controller therapy does not appear to be dependent on an individual's proportion of African ancestry.

Capsule summary—Personalized medicine will be most beneficial to groups disproportionately affected by disease complications. Here we find baseline bronchodilator reversibility but not African ancestry to be associated with ICS responsiveness among African American patients with asthma.

Keywords

inhaled corticosteroids; asthma; race-ethnicity; continental population groups; ancestry; urban health

INTRODUCTION

Over 20 million individuals are estimated to have asthma currently in the U.S.(1) African American individuals appear to be particularly susceptible to asthma-related complications with rates of asthma-related emergency department visits, hospitalizations, and death approximately 2-3 times the rates found in white individuals.(1) There may be many reasons for these apparent disparities,(2) including differences in the following factors: allergic sensitization and environmental exposures,(3-6) access to care,(7) adherence to and utilization of asthma controller therapy,(8;9) and treatment response.(10;11)

Inhaled corticosteroids (ICS) are considered first-line therapy for the treatment of persistent asthma,(12) as their use has been consistently shown to reduce airway responsiveness, diminish symptoms, and prevent exacerbations.(13-15) However, at least one *in vitro* study has suggested that steroid responsiveness may differ between African American and white individuals, as evidenced by between group differences in the ability of dexamethasone to suppress T-cell proliferative responses.(16) Given the existing disparities in asthma morbidity by race-ethnicity and the importance of ICS therapy in asthma control, it is important to clarify the factors associated with ICS responsiveness among African American

patients with asthma as this knowledge may improve the tailoring of asthma controller therapy.

As we and others have previously shown, there tends to be considerable variation in the proportion of African and European ancestry among African American individuals.(3;17) Since genetic substructure tends to follow historic geographic boundaries between ancestral groups,(18) we reasoned that treatment response may also differ according to biogeographic ancestry among African American patients with asthma. In other words, were between population genetic differences the explanation for disparities in treatment response among patients with asthma, we would also expect this to be reflected in a significant relationship between ancestral proportion and ICS response within African American individuals.

Therefore, this study sought to identify the degree to which African American patients with asthma respond to ICS therapy and the factors associated with treatment response (e.g., age, sex, socioeconomic status, asthma duration, allergic sensitization, passive smoke exposure, and pulmonary function). Many of these factors have been associated with asthma control and severity, and thus may also influence treatment response.(6;19-21) We were also keenly interested in assessing for a potential association between genetic ancestry and ICS treatment response for the reasons mentioned. Treatment response was defined by both the change in pulmonary function (i.e., forced expiratory volume at one second [FEV₁]) and the change in patient reported asthma control following 6 weeks of inhaled beclomethasone dipropionate treatment. Study participants were self-identified African American patients with asthma recruited within the city of Detroit as part of the Study of Asthma Phenotypes and Pharmacogenomic Interactions by Race-Ethnicity (SAPPHIRE).

METHODS

Study population and setting

The Institutional Review Board of Henry Ford Health System approved the study as described. All study patients received care through a large health system serving southeast Michigan. Those patients residing within the outer city boundaries of Detroit (including the cities of Hamtramck and Highland Park) were invited for screening if they met the following criteria: age 12-56 years, at least one coded clinical diagnosis of asthma (i.e., from the electronic record), and no recorded diagnosis of chronic obstructive pulmonary disease or congestive heart failure. The age criterion was selected to be consistent with current U.S. asthma guidelines which recommend similar controller treatment for individuals who are 12 years of age and older.(12) Informed written consent was obtained for study participants or their guardians prior to the first screening examination, and written assent was obtained from all participants who were minors.

Screening assessment

On the initial screening visit, patients answered a staff-administered structured questionnaire, performed spirometry with an assessment of bronchodilator response, and provided a blood sample for DNA analysis and serologic testing. Where available, the survey instrument included validated questions adapted from multiple sources. These questions included asthma onset, disease triggers, asthma control as assessed by the Asthma Control Test (QualityMetric Inc., Lincoln, RI),(22) detailed household socioeconomic questions and self-identified race-ethnicity (available at <http://www.census.gov/dmd/www/2000quest.html>), health-related missed work and school, (23) tobacco exposure,(24;25) environmental smoke exposure (available at http://www.cdc.gov/nchs/data/nhanes/nhanes_05_06/fi_smq_d.pdf),(25) and environmental

exposures.(26) A venous blood sample was collected for subsequent DNA isolation and serologic testing.

Spirometry was performed in accordance with current 2005 ATS/ERS recommendations(27;28) using a Fleisch-type pneumotachometer (KoKo PFT Spirometer®, nSpire Health Inc., Louisville, CO). Predicted values for pulmonary function parameters, such as the forced expiratory volume at one second (FEV₁), were obtained using standard equations derived from the U.S. population.(29) If possible, patients were asked to withhold their bronchodilator medications 12 hours before lung function tests. We determined bronchodilator response by administering increasing doses of inhaled albuterol sulfate hydrofluoroalkane (HFA) (GlaxoSmithKline, Research Triangle Park, NC). A 360µg dose of albuterol was delivered from a standard metered dose inhaler (MDI) using an AeroChamber Plus® Z STAT spacer (Monahan Medical Corp., Plattsburgh, NY). Albuterol was administered a second time (180µg for children <16 years, and 360µg for persons ≥16 years) if a >12% improvement in FEV₁ was not achieved following the first dose. Baseline bronchodilator reversibility was calculated as the percent change in FEV₁ pre- and post-albuterol administration on the initial screening visit.(30;31)

Inhaled corticosteroid treatment trial

Individuals who confirmed that they had a prior clinical diagnosis of asthma were eligible for the 6-week ICS treatment trial to assess corticosteroid responsiveness if they also met the following (i.e., in addition to the criteria for screening): a baseline FEV₁ between 40-90% predicted, >12% baseline bronchodilator reversibility, no smoking in the preceding year or <10 pack-year smoking history total, not pregnant at the time of enrollment and not intending to get pregnant during the 6-week treatment period, and were steroid naïve (i.e., not treated with an oral or inhaled corticosteroid in the 4 weeks preceding enrollment). Individuals who met these criteria and agreed to treatment received 160µg beclomethasone dipropionate HFA (Teva Specialty Pharmaceuticals LLC, Horsham, PA) twice daily per MDI for a total of 6 weeks. Participants were instructed in proper inhaler technique, including use of the provided spacer (AeroChamber Plus® Z STAT, Monahan Medical Corp., Plattsburgh, NY). Medication adherence was assessed electronically with the use of Doser CT® (Meditrack products, Hudson, MA), a monitoring device which attaches to the inhaler.(32;33) This was done to account for the level of use and its potential effect on treatment response over the study period. To estimate adherence we divided the number of canister actuations registered by the prescribed number of doses (i.e., 168) over the 6-week treatment period. In situations where the follow-up was not precisely at 6-weeks we adjusted the denominator accordingly for the number of days between the initial and follow-up visit.

DNA isolation, genotyping, and assessment of ancestry

Genomic DNA was extracted from whole blood using a Gentra AutoPure LS (Gentra Systems, Inc., Minneapolis, MN). Genotyping was performed with iPLEX® gold (Sequenom, Inc., San Diego, CA). Each individual was genotyped for 107 single nucleotide polymorphisms (SNPs), which were informative for determining individuals' ancestry due to their differing allele frequencies in continental population groups. This set of ancestry informative markers (AIMs) has been described in detail elsewhere,(34) but in short was developed from the following ancestral populations: West African individuals living in London, U.K. and South Carolina, U.S.; individuals of European ancestry from Coriell's North American Caucasian panel (Coriell Institute for Medical Research, Camden, New Jersey); and Native American individuals (i.e., Mayan and Nahua) from villages in Tlapa in the state of Guerrero, Mexico.

We used the software package PSMIX to determine individual ancestry by maximum likelihood estimation.(35;36) The procedure stopped when the change in the parameter estimate between consecutive iterations was less than 10^{-6} . As an independent check, we estimated ancestral proportions using the program STRUCTURE, which employs a Bayesian approach.(37) The correlation between ancestry estimates using both approaches was 0.99, but in the final analysis we used the estimates derived from PSMIX.

Allergen-specific and total serum immunoglobulin E (IgE) levels

Individual IgE levels were measured from stored serum samples. Assays for allergen-specific immunoglobulin E (IgE) were performed using a commercial assay (ImmunoCAP, Phadia AB, Portage, MI). The samples were analyzed for IgE levels to the following aeroallergens: dog (*Canis domesticus*), cat (*Felis domesticus*), cockroach (*Blattella germanica*), ragweed (short, *Ambrosia artemisiifolia*), grass (timothy, *Phleum pratense*), *Alternaria* (*Alternaria alternata*), and dust mite (*Dermatophagoides farinae*). Individual allergen-specific IgE levels greater than or equal to 0.35 international units per milliliter (IU/ml) were considered positive. Allergic sensitization was defined as having one or more positive allergen-specific IgE to one of the seven aeroallergens tested. We also measured total serum IgE using a commercial assay from the same manufacturer.

Statistical analysis

Our analysis was restricted to SAPPHERE participants who by self-report were African American. We assessed steroid responsiveness as the change in FEV₁ and Asthma Control Test (ACT) score pre- and post-ICS therapy. Linear regression was used to model these continuous outcome variables. We assessed the univariable and multivariable relationship between potential predictors of steroid responsiveness and outcomes. These predictor variables included age, sex, body mass index (in kilograms per meter squared), household income (log transformed to normalize its distribution), educational level (dichotomously defined as ≤high school degree vs. >high school), percent African ancestry, baseline ACT score (dichotomously defined as ≤19 [poor control] and >19 [good control], and included in only the model with change in FEV₁ as the dependent variable), asthma duration, individual allergen-specific sensitivity (dichotomously defined as positive or negative for each allergen - dog, cat, cockroach, ragweed, timothy grass, *Alternaria* species, and dust mite), total serum IgE level (natural log transformed to normalize the distribution and reduce the influence of outliers), passive smoke exposure (as all participants were reported non-smokers), baseline percent predicted FEV₁, baseline bronchodilator reversibility, and level of adherence to study medication. Since some study individuals were minors, we used the income and level of educational attainment for the head of the household as proxies for socioeconomic status. Backward stepwise regression was used to identify those factors related to steroid responsiveness. At each step the variable with the largest P-value was excluded in the model; the procedure stops when all remaining variables have P-values <0.05. As a secondary analysis, we assessed for factors associated with changes in patient-reported asthma control following 6-weeks of ICS therapy (i.e., the change in the ACT score pre- and post-therapy). Again we assessed the univariable and multivariable relationship between change in asthma control (continuously measured) and the aforementioned potential predictor variables (i.e., with the exception of baseline Asthma Control Test score). Backward stepwise regression was used to identify those factors related to steroid responsiveness as measured by improvement in patient-reported asthma control. All analyses were performed using SAS v9.1 (SAS Institute Inc., Cary, NC.)(38)

RESULTS

Of the 1,985 patients from Detroit screened, 1,777 (89.5%) identified themselves as African American. Of these patients, 214 met the criteria for enrollment in the 6-week ICS treatment trial and 185 agreed to enroll. Of those enrolled, 147 (79.5%) completed the 6-week assessment. The characteristics of those patients who did and did not complete the 6-weeks of treatment are shown in Table 1. None of the characteristics examined differed between those who finished treatment and those who did not. Among those who completed the study, average age was 32.7 years (standard deviation [SD], 13.6 years) and average duration of asthma was 21.1 years (SD, 12.8 years). The average proportion of African ancestry was 78.4%, and the distribution of African and European ancestry is shown in Figure 1. Over half, 87 (59.2%), had poorly controlled asthma at baseline as defined by an Asthma Control Test score ≤ 19 , and most, 126 (85.7%), had at least one positive allergen-specific IgE result. Adherence to treatment was good on average with patients taking 79.3% of their prescribed dose during the study period.

The average percentage improvement in pre-bronchodilator FEV₁ following 6 weeks of ICS treatment was 11.6% (Table 1) and the distribution of FEV₁ improvement across all study participants is shown in Figure 2. After accounting for all of the factors assessed, only the degree of baseline bronchodilator reversibility and the baseline FEV₁ percent of predicted were associated with ICS treatment response (Table 2 – adjusted model). Every 1% increase in baseline bronchodilator reversibility was associated with an ICS treatment-related improvement in FEV₁ of 0.48% (standard error [SE], 0.14%). Conversely, percent of predicted FEV₁ at baseline was inversely associated the change in FEV₁ in that individuals with lower percent predicted FEV₁ at baseline were more likely to improve following 6 weeks of treatment (0.41% improvement in FEV₁ per 1% lower baseline percent of predicted FEV₁). The backward stepwise regression model also demonstrated that baseline percent of predicted FEV₁ and baseline bronchodilator reversibility were associated with ICS treatment response ($P=0.001$ for both). In none of these models was the proportion of African ancestry significantly associated with the change in FEV₁ following 6 weeks of ICS treatment.

We also examined the factors associated with ICS treatment response as assessed by changes in patient reported asthma control (i.e., a change in ACT score over the course of therapy). In the unadjusted analysis, the degree of baseline bronchodilator reversibility (a 0.13 point improvement in ACT score per 1% increase in baseline bronchodilator reversibility, $P=0.006$) and the baseline FEV₁ percent of predicted (a 0.12 point improvement in ACT score per 1% lower baseline percent of predicted FEV₁, $P=0.013$) were associated with subsequent patient-reported treatment response (Table 3). Household income was also negatively associated with the change in the ACT score ($P=0.002$). In the stepwise regression model, household income was again negatively associated with treatment-related improvement in asthma control ($P=0.001$), whereas baseline bronchodilator reversibility was positively associated with treatment-related improvement in asthma control (a 0.11 point improvement in ACT score per 1% increase in baseline bronchodilator reversibility, $P=0.021$).

Post-hoc analysis using a 15% improvement in FEV₁ as a cut point for defining ICS responsiveness did not substantively change our conclusions with the exception that baseline ACT score was also significantly associated with this outcome in the stepwise model (data not shown). In addition, when categorizing the results of allergen-specific IgE values to just atopic and non-atopic (i.e., patients with ≥ 1 allergen-specific IgE result vs. none), atopy was still not associated with ICS treatment response. Finally, adding and removing socioeconomic variables (i.e., income and educational-level) from the regression models did

not affect the relationship (i.e., the lack of relationship) between ancestry and ICS treatment response.

DISCUSSION

This study is one of the largest to date to assess ICS responsiveness in African American patients with asthma. As a result, we have made a number of observations which may influence future approaches and opinions regarding asthma controller treatment in this population group.

First, we observed that the average improvement in FEV₁ following 6-week course of ICS therapy was 11.6% in our study population. This is comparable to other studies assessing ICS responsiveness using similar entry criteria but having a much lower proportion of African American patients. For example, in the Predicting Response to Inhaled Corticosteroid Efficacy (PRICE) trial, Martin et al. treated 72 individuals with 6 weeks of inhaled beclomethasone.(39) Patients were age 18-55 years, had a baseline percent of predicted FEV₁ between 55-85% with documented methacholine responsiveness, denied corticosteroid exposure for the 4 weeks preceding enrollment, and had <10 pack-years of smoke exposure with no smoking in the preceding year. Most of these individuals (55.6%) were considered non-Hispanic white. The mean improvement in FEV₁ was 8.3% after 6-weeks of therapy.

Similarly, Tantisira and colleagues examined ICS responsiveness after 6-8 weeks of ICS therapy in 3 clinical trials involving both children and adults.(40) The mean percent change in FEV₁ was 7.0%, 8.3%, and 6.7%, among studies comprised of 7.0%, 14.1%, and 18.8% African American patients, respectively. In short, improvement in FEV₁ following 6 weeks of ICS therapy was not lower in our SAPHIRE study population despite including only self-reported African American patients with asthma.

Differential response to asthma medications by race-ethnicity has been previously reported by us and others.(41-43) However, this differential response may be medication class specific, such as to inhaled beta-agonists.(44;45) In a recent study assessing step-up treatments for children with poorly controlled asthma, Hispanic and non-Hispanic white children were more likely to show improvement with an added long-acting beta-agonist medication (LABA) when compared with an additional leukotriene receptor antagonist medication or an ICS medication.(46) In contrast, African American children appear to respond equally well to LABA or ICS step-up therapy.

To our knowledge, this is the first study to evaluate the relationship between genetic ancestry and ICS responsiveness. We and others have previously demonstrated a wide range of African and European admixture among African American patients.(3;17;47) This is relevant to the current study in that genetic variation tends to follow historic geographic boundaries between populations.(24) Therefore, were there a strong genetic underpinning for disparate ICS response between persons of African and European ancestry, we would expect that association would be evident even when assessing the relationship with ancestry within African American individuals. Given our sample size and the variation in African ancestry within our study population, we estimate that we had 80% power to detect a correlation as small as 0.24 between ancestry and ICS responsiveness (as defined by a change in FEV₁). The proportion of African ancestry was not associated with ICS treatment response in any of our analyses.

This study also confirms the association between baseline bronchodilator responsiveness and steroid-related improvement in pulmonary function seen by other groups.(39;48) In fact, our data suggest that the degree of baseline bronchodilator responsiveness explains much

variation in treatment response, such as 32.9% of the variation in FEV₁ improvement and 5.2% of the variation in Asthma Control Test improvement (data not shown). The negative association that we observed between income level and the change in self-reported asthma control over the course of ICS treatment may relate to environmental exposures which mitigate treatment response or perhaps to differences in other stressors which affect asthma management.(49)

It is important to note this study's limitations. First, this study was performed in only one city and therefore may not be representative of African American patients in other U.S. locations. Moreover, it also remains to be seen if these findings hold for other admixed groups, such as Puerto Rican patients, who suffer from even higher rates of asthma.(2) Nevertheless, our study does constitute one of the largest evaluations of ICS responsiveness in African American patients to date, and our estimates of admixture are very similar to that reported for African American individuals in other areas of the United States.(34;50) In estimating African ancestry, we used markers informative for West African ancestry. While this assumption may have resulted in some incorrect estimates of ancestry, recent studies suggest that West African origin accounts for the majority of genetic variance in African American individuals, and that on average other African populations account for ~8% of individual ancestry.(51;52)

There are undoubtedly other factors associated with ICS responsiveness which may have been overlooked by us or await discovery (and therefore are not included in our models). Medication adherence, which we have previously shown to be associated with poor asthma outcomes,(53) was not found to be associated with treatment response. However, this is most likely the consequence of the high levels of medication adherence in this study overall (i.e., ~81% of study individuals had >50% adherence – data not shown), which allowed us to assess other factors associated with drug response.

Although our study population was limited to those patients who met our restrictive entry criteria, this was done so as to be comparable to existing studies of ICS responsiveness.(39;48) However, these criteria were not perfectly congruent with severity categories used in current U.S. guidelines.(12) Lastly, we assessed the change in FEV₁ as continuous measure of ICS response. We thought that this was a reasonable approach given the normal distribution of the outcome variable and the greater power inherent in this analytic design. Nevertheless, other studies have used various thresholds for FEV₁ improvement to define a therapeutic response.(39;48) As a post hoc analysis, we show that our study conclusions are not altered by using a 15% improvement in FEV₁ as a cut point.

In summary, our study found that the proportion of African ancestry was not significantly associated with response to inhaled steroids, the quintessential asthma controller medication. This does not rule out the possibility of genetic determinants specific to African and European ancestry, but rather suggests that such *local* genetic effects may be offsetting when assessed as an overall (or *global*) ancestral proportion. Moreover, although we observed a consistent relationship between the degree of baseline bronchodilator reversibility and ICS responsiveness, the proportion of variance explained suggests that there are other important determinants. Some genetic determinates of ICS response have already been described;(54-56) however, there are undoubtedly other important genetic, environmental, and gene-environment interactions which play a role in response to controller therapy. As these determinants become better known, we may be able to better tailor or *personalize* our selection of medications to those most likely to respond. This has particular relevance to groups disproportionately affected by disease complications,(57) such as African Americans patients with asthma. This study importantly suggests that *global* genetic ancestry may not be such a factor when considering ICS therapy for African

American patients with asthma; however, replication in other African American study populations with asthma are needed to confirm these findings.

Acknowledgments

This work was supported by grants from the Fund for Henry Ford Hospital; the American Asthma Foundation; and the National Institute of Allergy and Infectious Diseases (AI79139, AI61774), the National Heart Lung and Blood Institute (HL79055), and the National Institute of Diabetes and Digestive and Kidney Diseases (DK64695), National Institutes of Health (Dr. Williams). Dr. Burchard receives support from the National Institutes of Health (HL078885, HL088133, ES015794), the Robert Wood Johnson Foundation Amos Medical Faculty Development Program, and the Flight Attendant Medical Research Institute (FAMRI). Dr. Kumar receives support from the National Institutes of Health (K23HL093023-01).

Abbreviations

ICS	inhaled corticosteroid
FEV ₁	forced expiratory volume at one second
AIM	Ancestry informative marker

REFERENCES

- (1). Moorman JE, Rudd RA, Johnson CA, King M, Minor P, Bailey C, et al. National surveillance for asthma—United States, 1980–2004. *MMWR Surveill Summ* 2007;56(8):1–54. [PubMed: 17947969]
- (2). Joseph CL, Williams LK, Ownby DR, Saltzgaber J, Johnson CC. Applying epidemiologic concepts of primary, secondary, and tertiary prevention to the elimination of racial disparities in asthma. *J Allergy Clin Immunol* 2006;117(2):233–40. [PubMed: 16461121]
- (3). Yang JJ, Burchard EG, Choudhry S, Johnson CC, Ownby DR, Favro D, et al. Differences in allergic sensitization by self-reported race and genetic ancestry. *J Allergy Clin Immunol* 2008;122(4):820–7. [PubMed: 19014772]
- (4). Joseph CL, Ownby DR, Peterson EL, Johnson CC. Racial differences in physiologic parameters related to asthma among middle-class children. *Chest* 2000;117(5):1336–44. [PubMed: 10807820]
- (5). Gelber LE, Seltzer LH, Bouzoukis JK, Pollart SM, Chapman MD, Platts-Mills TA. Sensitization and exposure to indoor allergens as risk factors for asthma among patients presenting to hospital. *Am Rev Respir Dis* 1993;147(3):573–8. [PubMed: 8442589]
- (6). Rosenstreich DL, Eggleston P, Kattan M, Baker D, Slavin RG, Gergen P, et al. The role of cockroach allergy and exposure to cockroach allergen in causing morbidity among inner-city children with asthma. *N Engl J Med* 1997;336(19):1356–63. [PubMed: 9134876]
- (7). Wilper AP, Woolhandler S, Lasser KE, McCormick D, Bor DH, Himmelstein DU. A national study of chronic disease prevalence and access to care in uninsured U.S. adults. *Ann Intern Med* 2008;149(3):170–6. [PubMed: 18678844]
- (8). Williams LK, Joseph CL, Peterson EL, Moon C, Xi H, Krajenta R, et al. Race-ethnicity, crime, and other factors associated with adherence to inhaled corticosteroids. *J Allergy Clin Immunol* 2007;119(1):168–75. [PubMed: 17208598]
- (9). Williams LK, Joseph CL, Peterson EL, Wells K, Wang M, Chowdhry VK, et al. Patients with asthma who do not fill their inhaled corticosteroids: a study of primary nonadherence. *J Allergy Clin Immunol* 2007;120(5):1153–9. [PubMed: 17936894]
- (10). Chan MT, Leung DY, Szeffler SJ, Spahn JD. Difficult-to-control asthma: clinical characteristics of steroid-insensitive asthma. *J Allergy Clin Immunol* 1998;101(5):594–601. [PubMed: 9600494]
- (11). Naqvi M, Tcheurekdjian H, DeBoard JA, Williams LK, Navarro D, Castro RA, et al. Inhaled corticosteroids augment bronchodilator responsiveness in Latino but not African American asthmatics. *Ann Allergy Asthma Immunol* 2008;100(6):551–7. [PubMed: 18592818]

- (12). Expert Panel Report 3 (EPR-3): Guidelines for the Diagnosis and Management of Asthma-Summary Report 2007. *J Allergy Clin Immunol* 2007;120(5 Suppl):S94–138. [PubMed: 17983880]
- (13). Haahtela T, Jarvinen M, Kava T, Kiviranta K, Koskinen S, Lehtonen K, et al. Comparison of a beta 2-agonist, terbutaline, with an inhaled corticosteroid, budesonide, in newly detected asthma. *N Engl J Med* 1991;325(6):388–92. [PubMed: 2062329]
- (14). Kerstjens HA, Brand PL, Hughes MD, Robinson NJ, Postma DS, Sluiter HJ, et al. A comparison of bronchodilator therapy with or without inhaled corticosteroid therapy for obstructive airways disease. Dutch Chronic Non-Specific Lung Disease Study Group. *N Engl J Med* 1992;327(20):1413–9. [PubMed: 1357553]
- (15). Pauwels RA, Pedersen S, Busse WW, Tan WC, Chen YZ, Ohlsson SV, et al. Early intervention with budesonide in mild persistent asthma: a randomised, double-blind trial. *Lancet* 2003;361(9363):1071–6. [PubMed: 12672309]
- (16). Federico MJ, Covar RA, Brown EE, Leung DY, Spahn JD. Racial differences in T-lymphocyte response to glucocorticoids. *Chest* 2005;127(2):571–8. [PubMed: 15705998]
- (17). Parra EJ, Marcini A, Akey J, Martinson J, Batzer MA, Cooper R, et al. Estimating African American admixture proportions by use of population-specific alleles. *Am J Hum Genet* 1998;63(6):1839–51. [PubMed: 9837836]
- (18). Rosenberg NA, Pritchard JK, Weber JL, Cann HM, Kidd KK, Zhivotovsky LA, et al. Genetic structure of human populations. *Science* 2002;298(5602):2381–5. [PubMed: 12493913]
- (19). Eisner MD, Klein J, Hammond SK, Koren G, Lactao G, Iribarren C. Directly measured second hand smoke exposure and asthma health outcomes. *Thorax* 2005;60(10):814–21. [PubMed: 16192366]
- (20). Trupin L, Balmes JR, Chen H, Eisner MD, Hammond SK, Katz PP, et al. An integrated model of environmental factors in adult asthma lung function and disease severity: a cross-sectional study. *Environ Health* 2010;9(1):24. [PubMed: 20487557]
- (21). Bacon SL, Bouchard A, Loucks EB, Lavoie KL. Individual-level socioeconomic status is associated with worse asthma morbidity in patients with asthma. *Respir Res* 2009;10:125. [PubMed: 20017907]
- (22). Nathan RA, Sorkness CA, Kosinski M, Schatz M, Li JT, Marcus P, et al. Development of the asthma control test: a survey for assessing asthma control. *J Allergy Clin Immunol* 2004;113(1):59–65. [PubMed: 14713908]
- (23). Kessler RC, Barber C, Beck A, Berglund P, Cleary PD, McKenas D, et al. The World Health Organization Health and Work Performance Questionnaire (HPQ). *J Occup Environ Med* 2003;45(2):156–74. [PubMed: 12625231]
- (24). Leffondre K, Abrahamowicz M, Siemiatycki J, Rachet B. Modeling smoking history: a comparison of different approaches. *Am J Epidemiol* 2002;156(9):813–23. [PubMed: 12396999]
- (25). Plan and operation of the Third National Health and Nutrition Examination Survey, 1988–94. Series 1: programs and collection procedures. *Vital Health Stat* 1994;(32):1–407.
- (26). Burney PG, Luczynska C, Chinn S, Jarvis D. The European Community Respiratory Health Survey. *Eur Respir J* 1994;7(5):954–60. [PubMed: 8050554]
- (27). Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J* 2005;26(2):319–38. [PubMed: 16055882]
- (28). Miller MR, Crapo R, Hankinson J, Brusasco V, Burgos F, Casaburi R, et al. General considerations for lung function testing. *Eur Respir J* 2005;26(1):153–61. [PubMed: 15994402]
- (29). Hankinson JL, Odencrantz JR, Fedan KB. Spirometric reference values from a sample of the general U.S. population. *Am J Respir Crit Care Med* 1999;159(1):179–87. [PubMed: 9872837]
- (30). Lung function testing: selection of reference values and interpretative strategies. American Thoracic Society. *Am Rev Respir Dis* 1991;144(5):1202–18. [PubMed: 1952453]
- (31). National Heart LaBIN. Global Initiative for Asthma: Global Strategy for Asthma Management and Prevention. 2002. 02-3659
- (32). O'Connor SL, Bender BG, Gavin-Devitt LA, Wamboldt MZ, Milgrom H, Szeffler S, et al. Measuring adherence with the Doser CT in children with asthma. *J Asthma* 2004;41(6):663–70. [PubMed: 15584316]

- (33). Julius SM, Sherman JM, Hendeles L. Accuracy of three electronic monitors for metered-dose inhalers. *Chest* 2002;121(3):871–6. [PubMed: 11888975]
- (34). Yaeger R, vila-Bront A, Abdul K, Nolan PC, Grann VR, Birchette MG, et al. Comparing genetic ancestry and self-described race in african americans born in the United States and in Africa. *Cancer Epidemiol Biomarkers Prev* 2008;17(6):1329–38. [PubMed: 18559547]
- (35). Wu B, Liu N, Zhao H. PSMIX: an R package for population structure inference via maximum likelihood method. *BMC Bioinformatics* 2006;7(317)
- (36). Tang H, Peng J, Wang P, Risch NJ. Estimation of individual admixture: analytical and study design considerations. *Genet Epidemiol* 2005;28(4):289–301. [PubMed: 15712363]
- (37). Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus genotype data. *Genetics* 2000;155(2):945–59. [PubMed: 10835412]
- (38). SAS Institute Inc.. SAS/STAT Users Guide. Version 9.1 ed. SAS Institute Inc.; Cary, NC: 2004.
- (39). Martin RJ, Szeffler SJ, King TS, Kraft M, Boushey HA, Chinchilli VM, et al. The Predicting Response to Inhaled Corticosteroid Efficacy (PRICE) trial. *J Allergy Clin Immunol* 2007;119(1):73–80. [PubMed: 17208587]
- (40). Tantisira KG, Lake S, Silverman ES, Palmer LJ, Lazarus R, Silverman EK, et al. Corticosteroid pharmacogenetics: association of sequence variants in CRHR1 with improved lung function in asthmatics treated with inhaled corticosteroids. *Hum Mol Genet* 2004;13(13):1353–9. [PubMed: 15128701]
- (41). Naqvi M, Tcheurekdjian H, DeBoard JA, Williams LK, Navarro D, Castro RA, et al. Inhaled corticosteroids and augmented bronchodilator responsiveness in Latino and African American asthmatic patients. *Ann Allergy Asthma Immunol* 2008;100(6):551–7. [PubMed: 18592818]
- (42). Naqvi M, Thyne S, Choudhry S, Tsai HJ, Navarro D, Castro RA, et al. Ethnic-specific differences in bronchodilator responsiveness among African Americans, Puerto Ricans, and Mexicans with asthma. *J Asthma* 2007;44(8):639–48. [PubMed: 17943575]
- (43). Corvol H, De GA, Eng C, Seibold M, Ziv E, Chapela R, et al. Genetic ancestry modifies pharmacogenetic gene-gene interaction for asthma. *Pharmacogenet Genomics* 2009;19(7):489–96. [PubMed: 19503017]
- (44). Choudhry S, Ung N, Avila PC, Ziv E, Nazario S, Casal J, et al. Pharmacogenetic differences in response to albuterol between Puerto Ricans and Mexicans with asthma. *Am J Respir Crit Care Med* 2005;171(6):563–70. [PubMed: 15557128]
- (45). Corvol H, Burchard EG. Pharmacogenetic response to albuterol among asthmatics. *Pharmacogenomics* 2008;9(5):505–10. [PubMed: 18466098]
- (46). Lemanske RF Jr. Mauger DT, Sorkness CA, Jackson DJ, Boehmer SJ, Martinez FD, et al. Step-up Therapy for Children with Uncontrolled Asthma Receiving Inhaled Corticosteroids. *N Engl J Med*. 2010
- (47). Tsai HJ, Kho JY, Shaikh N, Choudhry S, Naqvi M, Navarro D, et al. Admixture-matched case-control study: a practical approach for genetic association studies in admixed populations. *Hum Genet* 2006;118(5):626–39. [PubMed: 16273390]
- (48). Szeffler SJ, Martin RJ, King TS, Boushey HA, Cherniack RM, Chinchilli VM, et al. Significant variability in response to inhaled corticosteroids for persistent asthma. *J Allergy Clin Immunol* 2002;109(3):410–8. [PubMed: 11897984]
- (49). Bryant-Stephens T. Asthma disparities in urban environments. *J Allergy Clin Immunol* 2009;123(6):1199–206. [PubMed: 19501229]
- (50). McKeigue PM, Carpenter JR, Parra EJ, Shriver MD. Estimation of admixture and detection of linkage in admixed populations by a Bayesian approach: application to African-American populations. *Ann Hum Genet* 2000;64(Pt 2):171–86. [PubMed: 11246470]
- (51). Tishkoff SA, Reed FA, Friedlaender FR, Ehret C, Ranciaro A, Froment A, et al. The genetic structure and history of Africans and African Americans. *Science* 2009;324(5930):1035–44. [PubMed: 19407144]
- (52). Bryc K, Auton A, Nelson MR, Oksenberg JR, Hauser SL, Williams S, et al. Genome-wide patterns of population structure and admixture in West Africans and African Americans. *Proc Natl Acad Sci U S A* 2010;107(2):786–91. [PubMed: 20080753]

- (53). Williams LK, Pladevall M, Xi H, Peterson EL, Joseph C, Lafata JE, et al. Relationship between adherence to inhaled corticosteroids and poor outcomes among adults with asthma. *The Journal of Allergy and Clinical Immunology* 2004;114(6):1288–93. [PubMed: 15577825]
- (54). Kazani S, Wechsler ME, Israel E. The role of pharmacogenomics in improving the management of asthma. *J Allergy Clin Immunol* 2010;125(2):295–302. [PubMed: 20159237]
- (55). Tantisira KG, Silverman ES, Mariani TJ, Xu J, Richter BG, Klanderman BJ, et al. FCER2: a pharmacogenetic basis for severe exacerbations in children with asthma. *J Allergy Clin Immunol* 2007;120(6):1285–91. [PubMed: 17980418]
- (56). Tantisira KG, Small KM, Litonjua AA, Weiss ST, Liggett SB. Molecular properties and pharmacogenetics of a polymorphism of adenylyl cyclase type 9 in asthma: interaction between beta-agonist and corticosteroid pathways. *Hum Mol Genet* 2005;14(12):1671–7. [PubMed: 15879435]
- (57). Daar AS, Singer PA. Pharmacogenetics and geographical ancestry: implications for drug development and global health. *Nat Rev Genet* 2005;6(3):241–6. [PubMed: 15738965]

Figure 1. Distribution of the African ancestry (blue) and European ancestry (gray) for African American patients with asthma who completed 6-weeks of inhaled corticosteroid therapy. Genetic ancestry was estimated in 135 of the 147 participants who completed therapy.

Figure 2. Distribution of the change in forced expiratory volume at one second (FEV₁) between baseline and following 6 weeks of inhaled corticosteroid treatment among study participants (n=147).

Table 1

Characteristics of African American participants with asthma who completed 6-weeks of inhaled corticosteroid therapy and those who did not

Characteristic	Individuals who completed the study (n=147)	Individuals who did not complete the study (n=38)	P-value *
Age (years) – mean $\pm$ SD [†]	32.7 $\pm$ 13.6	31.3 $\pm$ 13.0	0.574
Female – no. (%)	86 (58.5)	25 (65.8)	0.414
Body mass index (kg/m ²) – mean $\pm$ SD	33.3 $\pm$ 10.1	32.5 $\pm$ 11.2	0.650
Household income – median (range)	\$32,500 (\$1,000 – \$225,000)	\$27,500 (\$1,000 – \$162,000)	0.320
>High school education – no. (%)	89 (60.5)	18 (47.4)	0.143
African ancestry – mean proportion $\pm$ SD [‡]	78.4 $\pm$ 10.6	78.1 $\pm$ 12.4	0.888
Asthma Control Test score at baseline – mean $\pm$ SD [†]	17.5 $\pm$ 5.2	17.4 $\pm$ 4.9	0.935
Asthma Control Test score $\leq$ 19 (i.e., poor control) at baseline – no. (%)	87 (59.2)	26 (68.4)	0.298
Duration of asthma (years) – mean $\pm$ SD [§]	21.1 $\pm$ 12.8	22.9 $\pm$ 13.2	0.460
Positive allergen-specific IgE result (i.e., $\geq$ 0.35 kUA/l) – no. (%)			
Cat	72 (52.2)	23 (62.2)	0.279
Dog	79 (57.3)	20 (54.1)	0.728
Dust mite	69 (50.0)	17 (46.0)	0.661
Cockroach	60 (43.5)	16 (43.2)	0.980
Timothy Grass	67 (48.6)	19 (51.4)	0.762
Ragweed	80 (58.0)	17 (46.0)	0.191
Alternaria	75 (54.4)	21 (56.8)	0.794
$\geq$ 1 positive allergen-specific IgE	126 (85.7)	32 (84.2)	0.815
Total serum IgE level (kU/L) – geometric mean $\pm$ geometric SD	170.8 $\pm$ 4.1	185.1 $\pm$ 4.9	0.766
Passive smoke exposure – no. (%) [¶]	68 (46.3)	16 (42.1)	0.647
FEV ₁ percent predicted at baseline – mean $\pm$ SD [†]	74.6 $\pm$ 13.3	72.9 $\pm$ 12.6	0.470
FEV ₁ percent reversibility at baseline – mean $\pm$ SD [†]	22.0 $\pm$ 12.8	23.1 $\pm$ 13.9	0.629
Inhaled corticosteroid adherence during treatment period (%) – mean $\pm$ SD ^{**}	79.3 $\pm$ 22.1	--	--

Characteristic	Individuals who completed the study (n=147)	Individuals who did not complete the study (n=38)	P-value [*]
Average improvement in FEV ₁ after 6-weeks of ICS medication – mean $\pm$ SD [†]	11.6 $\pm$ 16.5	--	--

SD denotes standard deviation; IgE, immunoglobulin E; kAU/L, allergen-specific kilounits per liter; kU/L, kilounits per liter; FEV₁, forced expiratory volume in one second; ICS, inhaled corticosteroid; and NA, not assessed.

^{*} For the comparison of individuals who completed therapy with those who did not.

[†] Data available for 147 and 38 patients, respectively

[‡] African ancestry was estimated using the 107 ancestry informative markers and the software program PSMIX. Data available for 135 and 35 patients, respectively

[§] Data available for 146 and 36 patients, respectively

^{||} Data available for 138 and 37 patients

[¶] Passive smoke exposure was assessed by survey using questions derived elsewhere.(25)

^{**} Assessed as a ratio of actual canister actuations to prescribed dose over the 6-week treatment period. Data available for 135 patients

Characteristics related to the change in the forced expiratory volume in one second (FEV₁) following 6 weeks of inhaled corticosteroid treatment among African American patient with asthma*

Table 2

Characteristic	Unadjusted models		Adjusted model†		Backward stepwise model‡	
	% change in FEV ₁ ± SE	P-value	% change in FEV ₁ ± SE	P-value	% change in FEV ₁ ± SE	P-value
Age (per 10-year intervals)	1.52 ± 1.00	0.132	0.35 ± 1.37	0.800		
Female	-1.75 ± 2.77	0.528	-2.59 ± 2.78	0.354		
Body mass index (per 1 unit increase in kg/m ²)	0.19 ± 0.13	0.162	0.06 ± 0.14	0.698		
Household income	-0.77 ± 1.03	0.456	1.13 ± 1.09	0.303		
>High school education	-2.80 ± 2.78	0.316	0.19 ± 2.93	0.949		
African ancestry (per 10% increase in African ancestry)§	0.64 ± 1.36	0.637	-3.01 ± 13.30	0.821		
Asthma control test – poor control¶	6.44 ± 2.73	0.020	2.57 ± 2.79	0.359		
Duration of asthma (per 5-year intervals)	0.74 ± 0.53	0.167	-0.10 ± 0.64	0.879		
Positive allergen-specific IgE result¶						
Cat	5.57 ± 2.77	0.047	2.90 ± 3.95	0.465		
Dog	2.11 ± 2.84	0.459	-2.61 ± 4.03	0.519		
Dust mite	0.24 ± 2.81	0.933	-1.94 ± 3.65	0.596		

Characteristic	Unadjusted models		Adjusted model [†]		Backward stepwise model [‡]	
	% change in FEV ₁ ± SE	P-value	% change in FEV ₁ ± SE	P-value	% change in FEV ₁ ± SE	P-value
Cockroach	-1.89 ± 2.83	0.506	-2.37 ± 3.17	0.455		
Timothy Grass	4.07 ± 2.79	0.147	3.58 ± 3.25	0.272		
Ragweed	3.95 ± 2.83	0.165	3.65 ± 3.44	0.291		
Alternaria	0.59 ± 2.82	0.834	-2.62 ± 3.15	0.408		
Total serum IgE level ^{**}	0.96 ± 0.99	0.336	0.51 ± 1.35	0.707		
Passive smoke exposure ^{††}	-4.56 ± 2.71	0.095	-3.10 ± 2.61	0.237		
FEV ₁ percent predicted at baseline (per 1% decrease) ^{†‡}	0.68 ± 0.09	0.001	0.41 ± 0.14	0.005	0.42 ± 0.12	0.001
FEV ₁ percent bronchodilator reversibility at baseline (per 1% increase) ^{§§}	0.74 ± 0.09	0.001	0.48 ± 0.14	0.001	0.52 ± 0.13	0.001
Inhaled corticosteroid adherence during treatment period (per 10% increase in adherence)	0.04 ± 0.64	0.955	0.04 ± 0.58	0.949		

FEV₁ denotes the forced expiratory volume at one second; SE, standard error; and IgE, immunoglobulin E.

* Outcome was a continuous measure of the percentage increase in FEV₁ between the initial FEV₁ measured at the baseline visit and the initial FEV₁ measured at the follow-up visit following 6 weeks of inhaled corticosteroid therapy. The relationship between the predictor variables and outcome was assessed using linear regression.

[†] Adjusted for all other variables shown.

[‡] All variables shown are included in the model initially. The variables with the highest P-values are sequentially dropped and the model is refit after each exclusion until all the variables left are significant at the P<0.05 level. The variables presented are those that remain in the final model.

§ African ancestry was estimated using the 107 ancestry informative markers and the software program PSMIX.

|| The Asthma Control Test has a score range of 5-25 with higher scores signifying better control. The results were dichotomized with patient scores ≤ 19 considered to be poor control.

¶ Individual allergen-specific IgE results ≥ 0.35 kUA/l were considered to be positive.

** Total IgE levels were natural log transformed to normalize their distribution.

†† All participants were current non-smokers. Passive smoke exposure was assessed by survey using questions derived elsewhere.(25)

‡‡ Based on the predictive equation of Hankinson et al.(29)

§§ The percent change in the measured FEV₁ from baseline following the administration of albuterol. All participating patients had at least 12% reversibility at baseline.

||| Assessed as a ratio of actual canister actuations to prescribed dose over the 6-week treatment period.

Table 3

Characteristics related to the change in the Asthma Control Test following 6 weeks of inhaled corticosteroid treatment among African American patient with asthma*

Characteristic	Unadjusted models		Adjusted model [†]		Backward stepwise model [‡]	
	Change in ACT score $\pm$ SE	P-value	Change in ACT score $\pm$ SE	P-value	Change in ACT score $\pm$ SE	P-value
Age (per 10-year intervals)	-0.31 $\pm$ 0.44	0.478	-0.81 $\pm$ 0.63	0.202		
Female	0.71 $\pm$ 1.21	0.559	2.58 $\pm$ 1.26	0.043		
Body mass index (per 1 unit increase in kg/m ²)	0.04 $\pm$ 0.06	0.533	-0.03 $\pm$ 0.07	0.619		
Household income	-1.41 $\pm$ 0.44	0.002	-1.63 $\pm$ 0.48	0.001	-1.51 $\pm$ 0.42	0.001
>High school education	-1.09 $\pm$ 1.21	0.370	0.36 $\pm$ 1.34	0.790		
African ancestry (per 10% increase in African ancestry) [§]	-2.97 $\pm$ 5.94	0.618	-3.31 $\pm$ 6.09	0.588		
Duration of asthma (per 5-year intervals)	-0.06 $\pm$ 0.23	0.810	0.16 $\pm$ 0.17	0.577		
Positive allergen-specific IgE result						
Cat	1.05 $\pm$ 1.22	0.390	0.55 $\pm$ 1.79	0.760		
Dog	0.20 $\pm$ 1.24	0.870	-0.69 $\pm$ 1.83	0.707		
Dust mite	-0.88 $\pm$ 1.22	0.471	-1.75 $\pm$ 1.66	0.294		
Cockroach	-0.44 $\pm$ 1.24	0.722	-1.04 $\pm$ 1.45	0.475		
Timothy Grass	1.00 $\pm$ 1.22	0.413	1.94 $\pm$ 1.49	0.195		

Characteristic	Unadjusted models		Adjusted model [†]		Backward stepwise model [‡]	
	Change in ACT score $\pm$ SE	P-value	Change in ACT score $\pm$ SE	P-value	Change in ACT score $\pm$ SE	P-value
Ragweed	1.30 $\pm$ 1.24	0.294	2.59 $\pm$ 1.55	0.097		
Alternaria	-0.81 $\pm$ 1.23	0.508	-1.16 $\pm$ 1.43	0.420		
Total serum IgE level [¶]	0.09 $\pm$ 0.43	0.837	-0.32 $\pm$ 0.62	0.607		
Passive smoke exposure ^{**}	-1.50 $\pm$ 1.19	0.210	-1.89 $\pm$ 1.19	0.116		
FEV ₁ percent predicted at baseline (per 1% decrease) ^{††}	0.12 $\pm$ 0.05	0.013	0.03 $\pm$ 0.06	0.618		
FEV ₁ percent bronchodilator reversibility at baseline (per 1% increase) ^{‡‡}	0.13 $\pm$ 0.05	0.006	0.08 $\pm$ 0.06	0.200	0.11 $\pm$ 0.05	0.021
Inhaled corticosteroid adherence during treatment period (per 10% increase in adherence) ^{§§}	0.34 $\pm$ 0.26	0.190	0.27 $\pm$ 0.27	0.309		

FEV₁ denotes the forced expiratory volume at one second; SE, standard error; and IgE, immunoglobulin E.

* Outcome was a continuous measure of the difference in the Asthma Control Test between the baseline assessment and after 6-weeks of inhaled corticosteroid therapy. Positive changes reflect an improvement in patient reported asthma control over the treatment period. The relationship between the predictor variables and outcome was assessed using linear regression.

[†] Adjusted for all other variables shown.

[‡] All variables shown are included in the model initially. The variables with the highest P-values are sequentially dropped and the model is refit after each exclusion until all the variables left are significant at the P<0.05 level. The variables presented are those that remain in the final model.

[§] African ancestry was estimated using the 107 ancestry informative markers and the software program PSMIX.

[¶] Individual allergen-specific IgE results ≥ 0.35 kUA/l were considered to be positive.

^{¶¶} Total IgE levels were natural log transformed to normalize their distribution.

†† All participants were current non-smokers. Passive smoke exposure was assessed by survey using questions derived elsewhere.(25)

†† Based on the predictive equation of Hankinson et al.(29)

‡‡ The percent change in the measured FEV₁ from baseline following the administration of albuterol. All participating patients had at least 12% reversibility at baseline.

§§ Assessed as a ratio of actual canister actuations to prescribed dose over the 6-week treatment period.