CORRESPONDENCE

The Diagnosis of Urinary Tract Infection: A Systematic Review

by Dr. med. Guido Schmiemann MPH, Dr. med. Eberhard Kniehl, Klaus Gebhardt, Dr. med. Martha M. Matejczyk, Prof. Dr. med. Eva Hummers-Pradier in volume 21/2010

Methods of Investigation

The clinical diagnosis of urinary tract infection is beset by a high rate of errors, if the examination methods are faulty. Even in a primary care practice, microscopic urinalysis should be possible in order to make a diagnosis. So called asymptomatic urinary tract infections may require treatment in chronic urological pathologies. Primary attempts to treat a presumed urinary tract infection with fluoroquinolones often lead to antimicrobial resistance. Cotrimoxazole, amoxicillin, or nitrofurantoin may be used successfully.

The gold standard for a diagnosis of urinary tract infection, as the authors write, lies in identifying the causative agent by means of urine culture from midstream urine. Experiences from urological practice have shown that immersion culture media are often colonized by three or more pathogens, as a result of secondary contamination. After cleaning the external genitalia, bacteria should be identified in midstream urine by using a dipstick test, and antibiotic treatment should be given if significant bacteriuria (more than 100 000 pathogen germs/ml urine) is present. Pathogen concentrations of 10 000 require control examinations. And even a number of 1000 pathogens can indicate a urinary tract infection in isostenuria. In case of doubt, catheter examination will clarify the cause and enable a diagnosis.

“A burning sensation while urinating” in the patient's medical history is not pathognomonic for cystitis and also applies to bladder cancer.

Performing solely urinalysis by dipstick testing is certainly not sufficient to make a diagnosis. An absence of nitrites does not exclude a urinary tract infection.

The urine finding enables an exact diagnosis by conducting a dipstick test of midstream urine or catheter urine and by determining the causative strain and resistance patterns and microscopic findings.

DOI: 10.3238/arztebl.2010.0824a

REFERENCES

- Schmiemann G, Kniehl E, Gebhardt K, Matejczyk M, Hummers-Pradier E: The diagnosis of urinary tract infection: A systematic review. Dtsch Arztebl Int 2010; 107(21): 361–7.

Dr. med. Hans-Bernd Roleff
Peter-Nonnenmühlen-Allee 77,
41063 Mönchengladbach, Germany

Bacteriological finding

I read with great interest the critical review article on diagnosing urinary tract infections (UTI), which was based on publications. It showed beautifully the dilemma posed by the diagnostic evaluation of UTI. Even in publications this is not optimal. Too often, the sole focus is on laboratory findings (pathogen numbers that are regarded as the gold standard for UTI) or test strips. The clinical picture with findings from the vulva and the urethral area, and microscopic urinalysis are often neglected. I can understand that subjective findings count less because clinical aspects are often ignored during medical training and their implication, especially in studies, is usually less accepted. Unfortunately, this often leads to a situation whereby it is bacteriological findings that are treated, not infections.

Dysuria may indicate more than a UTI—for example, vulvitis, dermatitis of the urethral exit, skin damage owing to exaggerated cleaning routines, or recurrent genital herpes. Infections are acute events and are accompanied by an inflammatory reaction—that is, a raised number of leukocytes in the urine. This can be identified immediately by means of a wet mount enlarged 400 times. More than 3 leukocytes in the visual field are suspicious. Unfortunately, the publications mention only the esterase test for detecting leukocytes, and to detect bacteria, only microscopy. Both are highly unspecific. Especially in women, the main group of patients with UTI, the vulva, the vaginal flora, and the number of leukocytes in the vaginal discharge have to be considered because they all contaminate urine.

Skin improvements in the anal area, the source of intestinal bacteria that cause UTI, is important.

DOI: 10.3238/arztebl.2010.0824b

REFERENCES

- Schmiemann G, Kniehl E, Gebhardt K, Matejczyk M, Hummers-Pradier E: The diagnosis of urinary tract infection: A systematic review. Dtsch Arztebl Int 2010; 107(21): 361–7.

Prof. Dr. med. Eiko E. Petersen
Eichbergstraße 18
79117 Freiburg, Germany
eiko.petersen@web.de

Bladder puncture is required

The authors write: “The gold standard for the diagnosis of a urinary tract infection is the detection of the pathogen in the presence of clinical symptoms. The pathogen is detected and identified by urine culture (using midstream urine).” We wish to contradict this statement.

Women's urethras almost always contain pathogens and those of men often do so. Bladder puncture is therefore the only method that will deliver urine that provides a definite result. In 1972 we examined 1013 consecutive patients (192 male, 812 female) in a

nephrology outpatient department. UTI were suspected or needed to be ruled out in all of these patients (1). Bladder puncture was undertaken first and then mid-stream urine was collected. In 192 patients we found a rate of pathogens in excess of 100 000/mL. 30% of those 192 patients had a sterile bladder puncture specimen and would have received unnecessary antibiotic treatment if this limit had been adhered to. For a number of pathogens between 10 000 and 100 000, 61% of bladder puncture specimens were sterile, and for less than 10 000, 87%. In patients with sterile mid-stream urine (319 patients) the bladder puncture specimen was also sterile. Strictly speaking it is only sterile midstream urine that enables any safe assumptions. In many other cases, varying numbers of healthy patients will be treated and varying numbers of sick patients will not be treated. midstream urine can therefore not be the “gold standard.”

Thanks to the availability of ultrasonography, bladder puncture is an easy and safe procedure. It has been described as less painful than venepuncture. The patient has to put up with a certain bladder pressure before the puncture. Considering the amount of work involved for bacteriologists, clinicians should do their best to provide usable specimens.

To our knowledge until today the results have neither been confirmed nor falsified

DOI: 10.3238/arztebl.2010.0824c

REFERENCES

1. Fiegel P, Höffler D, Ringelmann R: Die Bedeutung der Blasenpunktion in der nephrologischen Diagnostik. *Die Medizinische Welt* 1972; 23: 1705–6.
2. Schmiemann G, Kniehl E, Gebhardt K, Matejczyk M, Hummers-Pradier E: The diagnosis of urinary tract infection: A systematic review. *Dtsch Arztebl Int* 2010; 107(21): 361–7.

Dr. med. Peter Fiegel
Am Keltenlager 6
55126 Mainz, Germany

Prof. Dr. med. Dieter Höffler
Rappmühlstraße 10
54331 Weiterstadt, Germany
dhoeffler@t-online.de

In Reply:

In our article we explained the need to distinguish between urinary tract infections and asymptomatic bacteriuria. The term asymptomatic urinary tract infection should be avoided. Clinical practice requires the differentiation between a disease and a condition – the latter requires treatment only in rare, individual cases (for example, eradication of asymptomatic bacteriuria in pregnant women). We refer our readers to the current German guidelines (1, 3) on diagnosis and therapy, where this terminology was consented. This differentiation should ensure that only infections, not bacteriological findings are treated as suggested by Fiegel and Höffler.

With regard to the relevance of leukocytes in urine, which Dr Roleff and Professor Petersen mention, we wish to add that the lack of a microscopic finding of

leukocytes has a negative predictive value of 95%, making it suitable for ruling out an infection. However, according to a meta-analysis, the sensitivity is only 37–96% and therefore less suitable for detecting the pathology (2). On this background, we question Dr Roleff's postulate that microscopic urine examination ought to be possible in the primary care setting. Dysuria (or the patient's reporting of “a burning sensation when urinating”) is not pathognomonic—and our article did not say it was. Reports of typical problems in the medical history (“burning sensation when urinating”) do, however, increase the probability that a UTI is present. Dysuria, though very occasionally present in (individual) cases of bladder cancer, is of very limited value with regard to diagnosing malignancies, due to their low prevalence, as well as low sensitivity and specificity of the symptom. “The gold standard to diagnose UTI in presence of typical symptoms is a bacteriological urine culture, with identification and quantification of the pathogen and sensitivity testing.” This definition was agreed upon by the author group of the S3 guideline for UTI in the Association for Scientific Medical Societies in Germany (AWMF, <http://www.uni-duesseldorf.de/AWMF/>). The method of collecting urine, as criticized by Fiegel and Höffler, was not part of this definition. Depending on how the urine was collected, different pathogen numbers are indicative of a diagnosis; several articles have compared the bacteriological findings from bladder puncture with other forms of collecting urine. Bladder puncture undoubtedly yields high quality specimens, but for various reasons (risk of injury, cost) it is not appropriate for routine diagnostic evaluation in adults in primary care. The situation may well be different for neonates/infants. The study mentioned by Fiegel and Höffler examined the (specialized) clientele of a nephrology outpatient ward and therefore cannot be translated to the situation in primary care.

DOI: 10.3238/arztebl.2010.0825

REFERENCES

1. Schmiemann G, Gebhardt K, Matejczyk M, Hummers-Pradier E: Brennen beim Wasserlassen – Anwenderversion der S3-Leitlinie Harnwegsinfekte. Düsseldorf: Omicron publishing 2009.
2. Whiting P, Westwood M, Bojke L, Palmer S, Richardson G, Cooper J, et al.: Clinical effectiveness and cost-effectiveness of tests for the diagnosis and investigation of urinary tract infection in children: a systematic review and economic model. *Health technology assessment* 2006;10(36): 1–154.
3. Epidemiologie, Diagnostik, antimikrobielle Therapie und Management von erwachsenen Patienten mit Harnwegsinfektionen Nr. 043/044 (in press) <http://www.uni-duesseldorf.de/WWW/AWMF/II/043-044-m.htm>
4. Schmiemann G, Kniehl E, Gebhardt K, Matejczyk M, Hummers-Pradier E: The diagnosis of urinary tract infection: A systematic review. *Dtsch Arztebl Int* 2010; 107(21): 361–7.

Dr. med. Eberhard Kniehl
Prof. Dr. med. Eva Hummers-Pradier

Dr. med. Guido Schmiemann MPH
Institut für Allgemeinmedizin
Medizinische Hochschule Hannover, 30625 Hannover, Germany
schmiemann.guido@mh-hannover.de

Conflict of interest statement

The authors declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.