Published in final edited form as:

Tetrahedron. 2010 August 14; 66(33): 6383–6390. doi:10.1016/j.tet.2010.04.002.

Dihydro-3-(triphenylphosphoranylidene)-2,5-thiophendione: A Convenient Synthron for the Preparation of Substituted 1,4-Thiazepin-5-ones and Piperidinones via the Intermediacy of Thioacids

David Crich^{a,b,*} and Md. Yeajur Rahaman^b

^aCentre de Recherche de Gif, Institut de Chimie des Substances Naturelles, CNRS, Avenue de la Terrasse, 91198 Gif-sur-Yvette, France

^bDepartment of Chemistry, Wayne State University, 5101 Cass Avenue, Detroit, MI 48202

Abstract

Reaction of thiomaleic anhydride with triphenylphosphine gives the title compound which undergoes reaction with a variety of aldehydes to give a range of alkylidene thiomaleic anhydrides (substituted monothio itaconic anhydrides). Subsequent treatment with *tert*-butoxycarbonylamino-substituted thiols, or under radical conditions with *tert*-butoxycarbonylamino-substituted alkyl halides results in a series of substituted monothiomaleic anhydrides, that on exposure to trifluoroacetic acid and then base lead to thiocarboxyl substituted 1,4-thiazepin-5-ones and piperidinones, respectively, that are ultimately trapped by reaction with 2,4-dinitrobenzenesulfonamides to give the corresponding amides.

1. Introduction

Monothioacids¹⁻⁶ present numerous advantages over the simple carboxylic acids by virtue of their acidity and the enhanced nucleophilicity of the thiocarboxylate function. Thus, for example, thioacids may be alkylated selectively in the presence of acids,⁷ and react with electrophiles such as electron-deficient aromatic and heteroaromatic systems,⁸⁻¹¹ isothiocyanates and isocyanates,^{12,13} isonitriles,¹⁴⁻¹⁷ a variety of coupling reagents,¹⁸⁻²¹ azides,²²⁻³⁰ and alkenes³¹⁻³⁴ more efficiently and under milder conditions than the corresponding acids. In this context we^{35,36} and others³⁷ have been exploring new methods for the preparation of a variety of thioacids for subsequent application in amide and peptide bond formation. In particular we have been interested in methods based on the nucleophilic ring opening of monothio thiosuccinic and thiomaleic anhydrides,³⁸⁻⁴² and of monothio- β -lactones⁴³ and on the application of the ensuing thioacids in multicomponent coupling processes. Here, we report on the preparation of dihydro-3-(triphenylphosphoranylidene)-2,5-thiophendione, its elaboration to a variety of monothio

© 2010 Elsevier Ltd. All rights reserved.

*dcrich@icsn.cnrs-gif.fr; fax: (33) 1 69 07 77 52.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

itaconic anhydride derivatives, and the use of these latter products in multicomponent coupling reactions leading to the formation of 1,4-thiazepin-5-ones.

2. Results and Discussion

By analogy with the chemistry of 2-(triphenylphosphanylidene)succinic anhydride,⁴⁴⁻⁵¹ we reasoned that treatment of monothiomaleic anhydride^{39,40,52,53} with triphenylphosphine would result in the formation of dihydro-3-(triphenylphosphoranylidene)-2,5-thiophendione **1**, and that this ylide would be stable, isolable, and a suitable partner for reaction with aldehydes to give substituted monothioitaconic anhydride derivatives. In the event, reaction of monothio maleic anhydride with triphenylphosphine in glacial acetic acid at room temperature provided the crystalline **1** in 86% yield (Scheme 1).

Subsequent reaction with a selection of aromatic and aliphatic aldehydes at room temperature gave the corresponding substituted monothioitaconic anhydrides in excellent yields as the pure *E*-isomers (Table 1), as ascertained by nOe studies, in agreement with the comparable reactions⁴⁶ of 2-(triphenylphosphanylidene)succinic anhydride.

With a series of alkylidene monothiosuccinic anhydrides in hand, we turned our attention to tandem Michael addition – nucleophilic ring opening processes with suitably constituted mercapto amines, of the kind previously practiced on monothiomaleic anhydride,^{39,40} with subsequent trapping of the released thioacid by reaction with 2,4-dinitrobenzenesulfonamides.^{9-11,36} A series of reactions were conducted in which either 2-mercaptoaniline or 2-mercaptoethylamine were stirred in DMF at 0 °C overnight before the addition of cesium carbonate and a 2,4-dinitrobenzenesulfonamide leading, ultimately, to the isolation of a series of substituted 1,4-thiazepin-5-ones, or their benzo-fused analogues, as presented in Table 2.

For each of the tandem sequences reported in Table 2 only the thiazepinone regioisomer, arising from ring closure of the amine onto the proximal carbonyl group following conjugate addition, was isolated.⁵⁴ In the case of the ethylidene succinic anhydride (Table 2, entries 5-8), two diastereomeric products were formed in an approximate equimolar ratio and no attempt was made to separate the two isomers. With moderately more bulky octylidene and cyclopropylidene systems **4** and **5** (Table 2, entries 9-12), low to moderate selectivity was observed in favor of the *trans*-isomer, whereas in the case of the somewhat more hindered benzylidene system **2** the *trans*-isomer was typically favored by a factor of 4-5:1 over the *cis*-isomer (Table 2, entries 1-4). This preferential formation of the *trans*-isomers, with selectivity increasing with the size of the olefinic substituent, is best rationalized in terms of the standard models for stereoselective protonation of enolates bearing a stereogenic center β-to the incipient carbonyl group.⁵⁵⁻⁶⁴ Thus, following conjugate addition of the thiolate anion, the enolate adopts a conformation that minimizes A^{1,3} strain^{65,66} leading to protonation through either of transitions states **A** or **B** (Scheme 2). As the size of the alkene substituent (*R*) increases transition state **B** is increasingly favored, resulting in overall increased *trans*-selectivity.⁶⁷

Having demonstrated the 1,4-addition sequence with aminothiols, we turned our attention to the conjugate addition of functionalized carbon radicals and the preparation of piperidinone-based systems. Thus, radical addition of *N*-(*tert*-butoxycarbonyl) 2-iodoethylamine **22** to the ethylidene succinic anhydride **3** in toluene at 80 °C under the aegis of tris(trimethylsilyl)silane (TTMS)⁶⁸ and azobisisobutyronitrile (AIBN), gave the adduct **23** in 52% yields as a 3:1 *anti*–*syn*-mixture of diastereoisomers (Scheme 3), whose stereochemistry was assigned following subsequent conversion to the *cis*- and *trans*-piperidinones, respectively (*vide infra*). The stereoselectivity of this radical addition is

rationalized by a model directly analogous to that advanced for enolate protonation in Scheme 2. Thus, hydrogen atom transfer to the α -carbonyl radical occurs from the face opposite the more bulky alkyl chain in a conformation that minimizes $A^{1,3}$ strain, as is the case for other related diastereoselective radical reactions.⁶⁹⁻⁷⁴

Subsequent treatment of this mixture of diastereomers with trifluoroacetic acid removed the *tert*-butoxycarbonyl protecting groups resulting, after removal of the acid under vacuum and treatment with base, in the cyclization of the liberated amine onto the proximal carbonyl carbon and the formation of the piperidinone skeleton bearing a thioacid substituent. Without isolation, this last species was captured by reaction with electron-deficient sulfonamides resulting overall in the piperidinonyl acetamides **24** and **25** (Scheme 4).^{54,75}

3. Conclusion

Dihydro-3-(triphenylphosphoranylidene)-2,5-thiophendione is readily accessed by reaction of monothiomaleic anhydride with triphenylphosphine. It reacts with a range of aldehydes in highly *E*-selective Wittig reactions to give substituted monothio itaconic anhydride derivatives. These latter substances undergo conjugate addition reactions with *tert*-butoxycarbonylamino-derived thiols and alkyl radicals to give a series of adducts that, on treatment with trifluoroacetic anhydride followed by base, lead to thiocarboxyl-substituted heterocycles. Finally, the thiocarboxylates can be trapped in situ by reaction with 2,4-dinitrobenzenesulfonamides to give the corresponding amides.

4. Experimental Section

4.1. Dihydro-3-(triphenylphosphoranylidene)-2,5-thiophendione (**1**)

Triphenylphosphine (2.62 g, 10 mmol) was added to monothiomaleic anhydride (1.14 g, 10 mmol) in glacial acetic acid (5 mL) at room temperature and stirred for 3 h. Removal of the solvent under vacuum at room temperature and trituration of the residue with diethyl ether (3 $\times$ 5 mL) afforded an off-white solid in 86 % yield. Mp: 137-137.5 °C (decomp.); IR (film) 1700 and 1611 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.69-7.67 (m, 9H), 7.56 (dt, J = 2.5, 7.5 Hz, 6H), 3.34 (d, J = 1 Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 205.6, 178.8, 133.6, 129.6, 124.8, 124.07, 50.0; ^{31}P NMR (400 MHz, CDCl_3) δ 12.78; ESIHRMS, m/z calcd. for $\text{C}_{22}\text{H}_{17}\text{O}_2\text{PSNa}$ ($M + \text{Na}$) $^+$: 399.0585, found: 399.0588.

4.2. General Procedure for Wittig Reactions

Reagent **1** (376 mg, 1 mmol) was stirred with excess aldehyde (1 mL) at room temperature for 10 h. Excess aldehyde was removed under vacuum at room temperature and the residue was purified by chromatographic over silica gel, that had been prewashed with acetone followed by hexanes, to give the alkylidene monothiosuccinic anhydrides.

4.2.1. *E*-2-(Benzylidene)thiosuccinic Anhydride (2**)**—Rapid chromatographic purification over silica gel eluting with 10 % EtOAc in hexanes afforded a pale yellow oil in 78 % yield. IR (film) 1731 and 1692 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.68 (t, J = 2.5 Hz, 1H), 7.51-7.46 (m, 5H), 3.98 (d, J = 2.5 Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 197.7, 192.7, 135.3, 133.8, 131.2, 131.1, 130.8, 130.1, 129.5, 128.4, 45.3; ESIHRMS m/z calcd for $\text{C}_{11}\text{H}_7\text{O}_2\text{S}$ (M 1) $^-$: 203.0172, found: 203.0180.

4.2.2. *E*-2-(Ethylidene)thiosuccinic Anhydride (3**)**—Rapid chromatographic purification over silica gel eluting with 10 % EtOAc in hexanes afforded a colorless oil in 76 % yield. IR (film) 1742 and 1706 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 6.99-6.94 (m, 1H), 3.66-3.65 (m, 2H), 1.91 (dt J = 2.0, 7.5 Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 197.6,

191.3, 135.2, 134.0, 43.2, 16.5; ESIHRMS m/z calcd for $C_6H_5O_2S$ (M-1)⁻: 141.0016, found: 141.0022.

4.2.3. E-2-(Octylidene)thiosuccinic Anhydride (4)—Rapid chromatographic purification over silica gel eluting with 10 % EtOAc in hexanes afforded a colorless oil in 72 % yield. IR (film) 1742 and 1706 cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 6.91-6.87 (m, 1H), 3.64 (dd, J = 1.0, 2.5 Hz, 2H), 2.19 (q, J = 7.5 Hz, 2H), 1.56-1.50 (m, 2H), 1.38-1.24 (m, 8H), 0.89 (t, J = 7.0 Hz, 3H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 197.7, 191.5, 140.5, 132.8, 43.3, 31.9, 31.0, 29.6, 29.2, 28.2, 22.8, 14.3; ESIHRMS m/z calcd for $C_{12}H_{17}O_2S$ (M-1)⁻: 225.0955, found: 225.0941.

4.2.4. E-2-(Cyclopropylmethylidene)thiosuccinic Anhydride (5)—Rapid chromatographic purification over silica gel eluting with 10 % EtOAc in hexanes afforded a colorless oil in 84 % yield. IR (film) 1737 and 1699 cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 6.27 (dt, J = 11.0, 2.5 Hz, 1H), 3.77 (d, J = 2.0 Hz, 2H), 1.45-1.38 (m, 1H), 1.16-1.12 (m, 2H), 0.85-0.81 (m, 2H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 198.2, 190.9, 145.6, 130.0, 43.6, 14.4, 10.3; ESIHRMS m/z calcd for $C_8H_7O_2S$ (M-1)⁻: 167.0172, found: 167.0166.

4.3. General Procedure for the Three Component Coupling of Alkylidene Thiosuccinic Anhydrides with Amino thiols and 2,4-Dinitrobenzenesulfonamides

The amino thiol (0.5 mmol) was added to a stirred solution of the alkylidene thiosuccinic anhydride (0.5 mmol) in DMF (5 mL) at 0 °C. The reaction mixture was stirred for overnight at 0 °C before Cs_2CO_3 (0.6 mmol) was added, followed immediately by the sulfonamide (0.5 mmol). Stirring was continued while the reaction mixture was allowed to warm to room temperature and the subsequently for a further 1 h. The solvent was then removed under vacuum and the residue was dissolved in ethyl acetate, washed with water followed by brine, and dried over Na_2SO_4 . Purification was then realized by silica gel column chromatography.

4.3.1. *cis*- and *trans*-N-[(Benzo[*b*]-5-oxo-7-phenyl-1,4-thiazepin-6yl)acetyl]piperidine (10)—Chromatographic purification by eluting with 70% ethyl acetate in hexanes afforded the *trans* and *cis* isomers with yields of 60% and 12% respectively.

***trans*-Isomer:** pale yellow solid, Mp 226-227 °C; IR (film) 1678 and 1631 cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 7.53 (d, J = 7.5 Hz, 1H), 7.44 (t, J = 7.7 Hz, 1H), 7.35 (d, J = 7.5 Hz, 1H), 7.30-7.24 (m, 3H), 7.20 (t, J = 7.2 Hz, 2H), 7.11 (d, J = 7.5 Hz, 2H), 4.32 (d, J = 12.5 Hz, 1H), 3.61 (dt, J = 2.0, 11.5 Hz, 1H), 3.46-3.39 (m, 1H), 3.36-3.24 (m, 2H), 3.18-3.12 (m, 1H), 3.08 (dd, J = 11.0, 16.0 Hz, 1H), 1.80 (d, J = 16.0 Hz, 1H), 1.56-1.32 (m, 6H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 174.5, 169.0, 143.4, 141.7, 136.0, 130.8, 129.2, 128.1, 126.7, 126.6, 126.3, 124.0, 56.9, 46.5, 44.6, 43.0, 33.9, 26.3, 25.6, 24.6; ESIHRMS m/z calcd for $C_{22}H_{24}N_2O_2SNa$ (M + Na)⁺: 403.1456, found: 403.1473.

***cis*-Isomer:** yellow solid, Mp. 163-164 °C; IR (film) 1676 and 1637 cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 7.66 (dd, J = 1.5, 8.0 Hz, 1H), 7.54 (br s, 1H), 7.51 (d, J = 7.0 Hz, 2H), 7.42-7.34 (m, 4H), 7.21 (t, J = 7.0 Hz, 2H), 5.02 (d, J = 7.0 Hz, 1H), 3.66-3.61 (m, 1H), 3.59-3.54 (m, 1H), 3.34-3.29 (m, 1H), 3.17-3.07 (m, 2H), 3.53 (dd, J = 8.0, 16.5 Hz, 1H), 2.07 (dd, J = 5.5, 16.5 Hz, 1H), 1.55-1.37 (m, 6H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 173.8, 168.6, 141.9, 137.2, 135.3, 130.6, 129.7, 128.8, 128.6, 126.7, 123.4, 57.8, 46.5, 43.0, 42.1, 32.9, 26.3, 25.7, 24.7; ESIHRMS m/z calcd for $C_{22}H_{24}N_2O_2SNa$ (M + Na)⁺: 403.1456, found: 403.1469.

4.3.2. *cis*- and *trans*-*N*-(2-Phenylethyl) (Benzo[*b*]-5-oxo-7-phenyl-1,4-thiazepin-6yl)acetamide (11)—Chromatographic purification eluting with 70% ethyl acetate in hexanes afforded the *trans* and *cis* isomers with yields of 63% and 13%, respectively.

***trans*-Isomer:** pale yellow solid, Mp 214-215 °C; IR (film) 1665 and 1642 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.56 (d, *J* = 8 Hz, 1H), 7.50-7.45 (m, 2H), 7.34-7.26 (m, 5H), 7.24-7.20 (m, 2H), 7.14 (d, *J* = 7.5 Hz, 3H), 7.09 (d, *J* = 6.5 Hz, 2H), 5.56 (br s, 1H), 4.27 (d, *J* = 12 Hz, 1H), 3.55 (dt, *J* = 2.5, 11.5 Hz, 1H), 3.39 (q, *J* = 6.5 Hz, 2H), 2.73-2.65 (m, 3H), 1.82 (dd, *J* = 2.5, 15 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 173.9, 170.7, 142.9, 141.1, 139.0, 136.2, 130.8, 129.3, 129.0, 128.8, 128.2, 127.1, 126.7, 126.4, 123.9, 57.1, 44.9, 40.8, 36.6, 35.8; ESIHRMS *m/z* calcd for C₂₅H₂₄N₂O₂SNa (M + Na)⁺: 439.1456, found: 439.1457.

***cis*-Isomer:** pale yellow solid, Mp 151-151.5 °C; IR (film) 1671 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.66 (d, *J* = 6.5 Hz, 1H), 7.60 (br s, 1H), 7.47-7.38 (m, 3H), 7.35-7.28 (m, 4H), 7.24-7.12 (m, 6H), 5.45 (br s, 1H), 4.94 (d, *J* = 7.0 Hz, 1H), 3.57-3.52 (m, 1H), 3.47-3.34 (m, 2H), 2.72 (t, *J* = 7.0 Hz, 2H), 2.28 (dd, *J* = 9.0, 15.0 Hz, 1H), 1.89 (dd, *J* = 5.0, 15.0 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 173.4, 170.7, 141.5, 138.9, 136.8, 135.4, 130.6, 129.7, 128.9, 128.8, 128.4, 127.0, 126.7, 123.4, 58.2, 42.5, 40.1, 36.2, 35.7; ESIHRMS *m/z* calcd for C₂₅H₂₄N₂O₂SNa (M + Na)⁺: 439.1456, found: 439.1462.

4.3.3. *cis*- and *trans*-*N*-[(5-Oxo-7-phenyl-1,4-thiazepin-6yl)acetyl]piperidine (12)—Chromatographic purification eluting with 4% methanol in dichloromethane afforded the *trans* and *cis* isomers with yields of 52% and 13%, respectively.

***trans*-Isomer:** light yellow gum, IR (film) 1675 and 1614 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.28-7.25 (m, 2H), 7.22-7.19 (m, 3H), 5.85 (br s, 1H), 4.15 (d, *J* = 9.5 Hz, 1H), 3.66-3.54 (m, 2H), 3.52-3.43 (m, 2H), 3.33-3.28 (m, 1H), 3.14-3.05 (m, 3H), 2.96-2.85 (m, 3H), 1.50-1.18 (m, 6H); ¹³C NMR (125 MHz, CDCl₃) δ 171.7, 171.6, 139.4, 129.7, 128.5, 126.7, 47.0, 43.8, 43.4, 43.0, 42.0, 37.3, 28.5, 25.5, 24.6; ESIHRMS *m/z* calcd for C₁₈H₂₄N₂O₂SNa (M + Na)⁺: 355.1456, found: 355.1445.

***cis*-Isomer:** light yellow gum, IR (film) 1638 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.37-7.27 (m, 5H), 5.95 (t, *J* = 6.5 Hz, 1H), 4.10-4.00 (m, 2H), 3.89 (d, *J* = 10.0 Hz, 1H), 3.68-3.61 (m, 1H), 3.52-3.46 (m, 1H), 3.44-3.36 (m, 1H), 3.28-3.22 (m, 1H), 3.16-3.10 (m, 1H), 2.97-2.89 (m, 1H), 2.87-2.81 (m, 2H), 1.88 (dd, *J* = 3.0, 16.0 Hz, 1H), 1.58-1.30 (m, 6H); ¹³C NMR (125 MHz, CDCl₃) δ 177.0, 169.5, 140.5, 131.6, 129.3, 129.1, 128.1, 128.0, 47.3, 46.5, 46.0, 43.2, 43.0, 34.0, 32.1, 26.3, 25.6, 24.7; ESIHRMS *m/z* calcd for C₁₈H₂₄N₂O₂SNa (M + Na)⁺: 355.1456, found: 355.1454.

4.3.4. *cis*- and *trans*-*N*-(2-Phenylethyl) (5-Oxo-7-phenyl-1,4-thiazepin-6yl)acetamide (13)—Chromatographic purification eluting with 4% methanol in dichloromethane afforded a pale yellow gum in 67 % yield as 1:4 *cis*:*trans*-mixture.

***trans*-Isomer:** IR (film) 1666 and 1640 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.35-7.28 (m, 7H), 7.24-7.19 (m, 2H), 7.16 (d, *J* = 7.0 Hz, 2H), 6.01 (br s, 1H), 5.52 (br s, 1H), 4.02-3.84 (m, 3H), 3.66-3.60 (m, 1H), 3.49-3.35 (m, 2H), 2.96-2.87 (m, 1H), 2.82-2.71 (m, 2H), 2.43 (dd, *J* = 11.0, 14.7 Hz, 1H), 1.91 (dd, *J* = 3.2, 14.7 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 176.8, 171.5, 139.9, 139.0, 129.1, 129.0, 128.8, 128.3, 128.1, 126.6, 46.3, 43.6, 40.9, 37.5, 35.8, 32.5; ESIHRMS *m/z* calcd for C₂₁H₂₄N₂O₂SNa (M + Na)⁺: 391.1456, found: 391.1449.

***cis*-Isomer:** the minor *cis* isomer was identified in the mixture by characteristic signals at δ 6.09 (br s, 1H), 5.63 (br s, 1H), 3.06-3.01 (m, 1H), 2.53 (dd, J = 8.2, 14.7 Hz, 1H), 2.09 (dd, J = 4.7, 14.7 Hz, 1H) in the ^1H -NMR spectrum, and by δ 175.2, 171.2, 139.3, 128.7, 128.2, 126.7, 48.0, 45.9, 40.7, 38.1, 29.8 in the ^{13}C -NMR spectrum.

4.3.5. *cis*- and *trans*-*N*-[(Benzo[*b*]-5-oxo-7-methyl-1,4-thiazepin-6yl)acetyl]piperidine (14)—Chromatographic purification eluting with 70 % ethyl acetate in hexanes afforded a pale yellow solid in 74 % yield as 1:1 *cis*:*trans*-mixture.

***trans*-Isomer:** IR (film) 1674 and 1633 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.61 (dd, J = 1.5, 7.5 Hz, 1 H), 7.42 (br s, 1H), 7.40-7.32 (m, 1 H), 7.24 (d, J = 7.0 Hz, 1H), 7.19-7.13 (m, 1H), 3.57-3.51 (m, 1H), 3.47-3.38 (m, 2H), 3.33-3.27 (m, 1H), 3.16 (dd, J = 10.5, 16 Hz, 1H), 2.92 (dt, J = 2.5, 11 Hz, 1H), 2.24 (dd, J = 2.7, 16.2 Hz, 1H), 1.65-1.42 (m, 6H), 1.39 (d, J = 6.5 Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 175.0, 169.1, 141.5, 136.6, 135.0, 130.7, 128.3, 126.5, 125.8, 123.7, 48.3, 47.7, 46.7, 45.9, 43.2, 33.6, 26.5, 23.1, 16.9; ESIHRMS m/z calculated for $\text{C}_{17}\text{H}_{22}\text{N}_2\text{O}_2\text{SNa}$ ($M + \text{Na}$) $^+$: 341.1300, found: 341.1291.

***cis*-Isomer:** the *cis* isomer was identified in the mixture by characteristic signals at δ 7.57 (dd, J = 12.0, 7.7 Hz, 1H), 3.95-3.87 (m, 1H), 2.17 (dd, J = 3.7, 16.2 Hz, 1H), 1.37 (d, J = 6.5 Hz, 3H) in the ^1H -NMR spectrum, and by δ 173.3, 168.7, 141.8, 123.4, 126.4, 130.2, 47.7, 43.1, 41.5, 32.5, 25.7, 24.7 in the ^{13}C -NMR spectrum.

4.3.6. *cis*- and *trans*-*N*-(2-Phenylethyl) (Benzo[*b*]-5-oxo-7-methyl -1,4-thiazepin-6yl)acetamide (15)—Chromatographic purification eluting with 70 % ethyl acetate in hexanes afforded a pale yellow oil in 72 % yield as 1:1 *cis*:*trans*-mixture.

***trans*-Isomer:** IR (film) 1670 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.62 (d, J = 7.5 Hz, 1H), 7.57-7.30 (m, 4H), 7.25-7.14 (m, 5H), 6.15 (br s, 1H), 3.50-3.40 (m, 2H), 3.26-3.20 (m, 1H), 2.82-2.76 (m, 3H), 2.70 (dd, J = 10.5, 14.5 Hz, 1H), 2.30 (dd, J = 2.2, 14.2 Hz, 1H), 1.39 (d, J = 7.0 Hz, 3H), ^{13}C NMR (125 MHz, CDCl_3) δ 173.1, 170.8, 141.3, 139.0, 136.7, 130.7, 129.0, 128.8, 126.8, 125.8, 123.6, 48.0, 46.6, , 41.0, 35.9, 35.6, 23.0, 16.5; ESIHRMS m/z calculated for $\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_2\text{SNa}$ ($M + \text{Na}$) $^+$: 377.1300, found: 377.1284.

***cis*-Isomer:** the *cis* isomer was identified in the mixture by characteristic signals at δ 7.57 (d, J = 7.5 Hz, 1H), 7.08 (d, J = 7.5 Hz, 1H), 5.72 (br s, 1H), 3.87 (q, J = 6.5 Hz, 1H), 3.36-3.32 (m, 2H), 2.06 (dd, J = 3.5, 14.7 Hz, 1H), 1.31 (d, J = 6.5 Hz, 3H) in the ^1H -NMR spectrum, and by δ 175.1, 170.9, 141.1, 139.2, 135.1, 130.2, 128.1, 126.7, 123.4, 48.8, 42.3, 40.8, 36.5, 35.87 in the ^{13}C -NMR spectrum.

4.3.7. *cis*- and *trans*-*N*-[(5-Oxo-7-methyl-1,4-thiazepin-6yl)acetyl]piperidine (16)—Chromatographic purification eluting with 4% methanol in dichloromethane afforded a pale yellow oil in 58 % yield as 1:1 *cis*:*trans*-mixture.

***trans*-Isomer:** IR (film) 1663 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 6.00 (br s, 1H), 3.93-3.97(m, 1H), 3.60-3.49 (m, 5H), 3.40 (dt, J = 3.5, 9.7 Hz, 1H), 3.11-3.05 (m, 1H), 2.98 (dd, J = 11.7, 14.2 Hz, 1H), 2.54 (dd, J = 11.5, 14.5 Hz, 1H), 2.37 (dd, J = 3.2, 16.2 Hz, 1H), 1.70-1.51 (m, 6H), 1.38 (d, J = 7.0 Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 176.2, 169.2, 48.5, 46.7, 45.7, 43.1, 36.0, 33.2, 26.6, 25.8, 24.8, 20.3, 16.7; ESIHRMS m/z calculated for $\text{C}_{13}\text{H}_{22}\text{N}_2\text{O}_2\text{SNa}$ ($M + \text{Na}$) $^+$: 293.1300, found: 293.1284.

***cis*-Isomer:** the *cis* isomer was identified in the mixture by characteristic signals at δ 5.87 (br s, 1H), 4.03 (dd, J = 5.5, 8.0 Hz, 1H), 3.80-3.73 (m, 1H), 3.48-3.41 (m, 5H), 2.88 (dt, J =

4.5, 14.5 Hz, 1H), 2.83-2.74 (m, 3H), 2.24 (dd, $J = 5.5, 16.0$ Hz, 1H), 1.37 (d, $J = 7.0$ Hz, 3H) in the ^1H -NMR spectrum, and by δ 176.9, 169.4, 48.3, 46.8, 43.2, 42.2, 35.9, 34.7, 27.3, 26.5, 25.7 in the ^{13}C -NMR spectrum.

4.3.8. *cis*- and *trans*-*N*-(2-Phenylethyl) (5-oxo-7-methyl-1,4-thiazepin-6yl)acetamide (17)—Chromatographic purification eluting with 4% methanol in dichloromethane afforded a pale yellow oil in 62 % yield as 1:1 *cis*:*trans*-mixture.

***trans*-Isomer:** IR (film) 1649 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.31 (t, $J = 7.5$ Hz, 2H), 7.24-7.19 (m, 3H), 6.09 (br s, 1H), 5.96 (br s, 1H), 3.83-3.77 (m, 1H), 3.66-3.59 (m, 1H), 3.57-3.44 (m, 3H), 2.87 (dt, $J = 4.2, 19.7$ Hz, 1H), 2.82-2.77 (m, 4H), 2.52 (dd, $J = 5.5, 15.0$ Hz, 1H), 2.16 (dd, $J = 4.7, 14.2$ Hz, 1H), 1.33 (d, $J = 7.0$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 176.0, 171.2, 139.2, 129.0, 126.7, 49.2, 45.8, 42.2, 40.9, 38.6, 37.0, 36.2, 26.9, 20.3, 16.5; ESIHRMS m/z calculated for $\text{C}_{16}\text{H}_{22}\text{N}_2\text{O}_2\text{SNa}$ ($M + \text{Na}$) $^+$: 329.1300, found: 329.1302.

***cis*-Isomer:** the *cis* isomer was identified in the mixture by characteristic signals at δ 6.07 (br s, 1H), 5.94-5.90 (m, 1H), 3.89 (dd, $J = 4.5, 9.5$ Hz, 1H), 3.27-3.21 (m, 1H), 2.94 (dd, $J = 10.7, 15.0$ Hz, 1H), 2.75-2.66 (m, 2H), 2.38 (dd, $J = 3.7, 14.2$ Hz, 1H), 1.38 (d, $J = 7.0$ Hz, 3H) in the ^1H -NMR spectrum, and by δ 176.5, 171.3, 139.1, 128.8, 126.6, 49.1, 40.8, 36.3, 35.9, 28.9 in the ^{13}C -NMR spectrum.

4.3.9. *cis*- and *trans*-*N*-[(Benzo[*b*]5-oxo-7-octanyl-1,4-thiazepin-6yl)acetyl]piperidine (18)—Chromatographic purification eluting with 70 % ethyl acetate in hexanes afforded a pale yellow oil in 67% yield as 2:5 *cis*:*trans*-mixture.

***trans*-Isomer:** IR (film) 1677 and 1641 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.57 (t, $J = 8.0$ Hz, 1H), 7.51 (br s, 1H), 7.34-7.31 (m, 1H), 7.23 (d, $J = 8.0$ Hz, 1H), 7.17-7.12 (m, 1H), 3.53-3.36 (m, 4H), 3.19-3.09 (m, 1H), 3.00 (dt, $J = 2.0, 11.0$ Hz, 1H), 2.26 (d, $J = 15.5$ Hz, 1H), 1.84-1.70 (m, 1 H), 1.50-1.56 (m, 4H), 1.56-1.40 (m, 5H), 1.40-1.14 (m, 9H), 0.92-0.83 (m, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 175.3, 169.2, 141.8, 136.6, 130.6, 126.4, 126.3, 123.6, 54.2, 46.7, 44.6, 43.2, 43.1, 34.9, 33.5, 32.0, 29.8, 29.4, , 26.5, 25.7, 24.8, 22.9, 14.3; ESIHRMS m/z calculated for $\text{C}_{23}\text{H}_{34}\text{N}_2\text{O}_2\text{SNa}$ ($M + \text{Na}$) $^+$:425.2239, found: 425.2249.

***cis*-Isomer:** the *cis* isomer was identified in the mixture by characteristic signals at δ 7.65 (br s, 1H), 7.32 (t, $J = 7.5$ Hz, 3H), 7.05 (d, $J = 7.5$ Hz, 1H), 3.78-3.74 (m, 1H), 2.16 (dd, $J = 3.2, 16.2$ Hz, 1H) in the ^1H -NMR spectrum, and by δ 173.6, 168.8, 135.1, 130.0, 128.1, 123.4, 53.6, 41.4, 32.7, 27.7, 26.4, 24.7 in the ^{13}C -NMR spectrum.

4.3.10. *cis*- and *trans*-*N*-(2-Phenylethyl) (Benzo[*b*]5-oxo-7-octanyl -1,4-thiazepin-6yl)acetamide (19)—Chromatographic purification eluting with 70 % ethyl acetate in hexanes afforded a pale yellow oil in 61% yield as 2:5 *cis*:*trans*-mixture.

***trans*-Isomer:** IR (film) 1672 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.61-7.54 (m, 2H), 7.38 (dt, $J = 1.5, 7.5$ Hz, 1H), 7.35-7.30 (m, 3H), 7.25-7.14 (m, 4H), 6.18 (br s, 1H), 3.54-3.41 (m, 2H), 3.06 (t, $J = 10.0$ Hz, 1H), 2.88-2.70 (m, 4H), 2.32 (dd, $J = 2.5, 14.0$ Hz, 1H), 1.67-1.52 (m, 2H), 1.45-1.20 (m, 10H), 0.92-0.88 (m, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 175.2, 171.1, 141.4, 139.2, 136.7, 135.3, 130.6, 129.0, 128.5, 126.8, 126.7, 123.6, 54.6, 45.3, 40.9, 36.6, 35.8, 34.7, 32.0, 29.7, 29.4, 26.4, 22.9, 14.3; ESIHRMS m/z calculated for $\text{C}_{26}\text{H}_{34}\text{N}_2\text{O}_2\text{SNa}$ ($M + \text{Na}$) $^+$: 461.2239, found: 461.2261.

***cis*-Isomer:** the *cis* isomer was identified in the mixture by characteristic signals at δ 7.66 (br s, 1H), 7.07 (d, J = 8.0 Hz, 1H), 5.76 (br s, 1H), 3.77-3.72 (m, 1H), 3.40-3.35 (m, 1H), 2.06 (dd, J = 3.7, 14.2 Hz, 1H), 1.78-1.72 (m, 1H) in the ^1H -NMR spectrum, and by δ 173.2, 170.8, 141.2, 139.0, 130.1, 126.4, 123.3, 53.8, 42.2, 31.7, 27.5 in the ^{13}C -NMR spectrum.

4.3.11. *cis*- and *trans*-*N*-[(Benzo[*b*]-5-oxo-7-cyclopropyl-1,4-thiazepin-6yl)acetyl]piperidine (20)—Chromatographic purification eluting with 70 % ethyl acetate in hexanes afforded a pale yellow oil in 57% yield as 2:3 *cis*:*trans*-mixture.

***trans*-Isomer:** IR (film) 1677 and 1638 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.55 (dd, J = 1.0, 7.5 Hz, 1H), 7.38-7.31 (m, 2H), 7.24-7.12 (m, 2H), 3.58-3.35 (m, 5 H), 3.26 (dd, J = 3.0, 11.5 Hz, 1H), 3.20-3.15 (m, 1H), 3.09 (dd, J = 11.0, 16.0 Hz, 1H), 2.46 (dd, J = 5.0, 16.5 Hz, 1H), 1.52-1.40 (m, 6H), 0.76-0.60 (m, 2H), 0.35-0.25 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 173.9, 169.0, 141.6, 135.1, 130.1, 126.6, 123.5, 59.0, 45.3, 42.4, 32.8, 25.7, 24.7, 18.4, 12.5, 7.1, 3.9, ESIHRMS m/z calculated for $\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}_2\text{SNa}$ ($M + \text{Na}$) $^+$: 367.1456, found: 367.1450.

***cis*-Isomer:** the minor *cis* isomer was identified in the mixture by characteristic signals at δ 7.63 (dd, J = 1.0, 4.5 Hz, 1H), 2.73-2.67 (m, 1H), 0.53-0.48 (m, 2H) in the ^1H -NMR spectrum, and by δ 174.8, 169.4, 141.3, 136.1, 130.4, 128.7, 126.4, 123.4, 59.4, 46.7, 43.1, 33.5, 26.5, 9.0, 3.1 in the ^{13}C -NMR spectrum.

4.3.12. *cis*- and *trans*-*N*-(2-Phenylethyl) (Benzo[*b*]-5-oxo-7-cyclopropyl-1,4-thiazepin-6yl)acetamide (21)—Chromatographic purification eluting with 70% ethyl acetate in hexanes afforded a pale yellow oil in 60% yield as 2:3 *cis*:*trans*-mixture.

***trans*-Isomer,** IR (film) 1671 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.65 (dd, J = 1.0, 7.5 Hz, 1H), 7.40-7.30 (m, 4H), 7.25-7.15 (m, 5H), 5.93 (br s, 1H), 3.53-3.45 (m, 2H), 3.22-3.17 (m, 1H), 2.79 (q, J = 7.0 Hz, 2H), 2.70 (d, J = 7.0 Hz, 2H), 1.04-0.97 (m, 1H), 0.77-0.59 (m, 3H), 0.32-0.22 (m, 2H). ^{13}C NMR (125 MHz, CDCl_3) δ 174.6, 171.3, 140.8, 139.2, 136.2, 130.4, 129.0, 128.9, 128.8, 126.4, 123.4, 59.5, 45.9, 40.9, 36.7, 35.9, 31.5, 18.2, 9.0, 3.8; ESIHRMS m/z calculated for $\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_2\text{SNa}$ ($M + \text{Na}$) $^+$: 403.1456, found: 403.1472.

***cis*-Isomer:** the minor *cis* isomer was identified in the mixture by characteristic signals at δ 7.56 (dd, J = 1.2, 7.7 Hz, 1H), 7.09 (d, J = 7.5 Hz, 1H), 5.72 (br s, 1H), 3.42-3.38 (m, 1H), 3.11 (dd, J = 5.5, 10.5 Hz, 1H), 3.00 (dd, J = 9.7, 14.7 Hz, 1H), 2.64 (dd, J = 9.5, 11.5 Hz, 1H), 2.26 (dd, J = 4.5, 14.5 Hz, 1H), 1.19-1.14 (m, 1H), 0.92-0.86 (m, 1H), 0.51-0.46 (m, 1H) in the ^1H -NMR spectrum, and by δ 173.5, 171.1, 141.2, 135.1, 130.2, 128.5, 126.8, 126.7, 123.5, 59.4, 43.1, 35.9, 25.9, 12.4, 7.2, 3.0 in the ^{13}C -NMR spectrum.

4.4. 2-[3-(*tert*-Butyloxycarbonylamino)-2-methylpropyl]succinic thioanhydride (23)

Tris(trimethylsilyl)silane (373 mg, 1.5 mmol) and AIBN (33 mg, 0.2 mmol) in dry degassed toluene (5 mL) were added drop-wise to a stirred mixture of **22** (407 mg, 1.5 mmol) and **3** (142 mg, 1 mmol) in dry degassed toluene (10 mL) at 90 °C over 5 h by syringe pump under a N_2 atmosphere. When the addition was complete, the reaction mixture was allowed to stir for an additional 1 h at 90 °C before it was cooled to room temperature and the solvent was removed under vacuum. Rapid chromatographic purification over silica gel, pre-washed with acetone followed by hexanes, eluting with 30% ethyl acetate in hexanes afforded a colorless oil in 52% yield as a 3:1 inseparable *anti*:*syn* diastereomeric mixture.

***anti*-Isomer**—IR (film) 1708 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 4.58 (br s, 1H), 3.55-3.02 (m, 4H), 2.88 (dd, J = 4.0, 6.5 Hz, 1H), 2.26-2.18 (m, 1H), 1.66-1.58 (m, 1H),

1.54-1.41 (m, 10H), 1.04 (d, $J = 7.0$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 203.0, 199.9, 156.2, 79.6, 57.4, 44.4, 38.6, 35.7, 33.3, 28.6, 17.2; ESIHRMS m/z calculated for $\text{C}_{13}\text{H}_{21}\text{NO}_4\text{SNa}$ ($M + \text{Na}$) $^+$: 310.1089, found: 310.1082.

syn-Isomer—the minor *syn* isomer was identified in the mixture by characteristic signals at δ 2.84 (dd, $J = 4.0, 6.5$ Hz, 1H), 2.44-2.36 (m, 1H), 0.96 (d, $J = 7.0$ Hz, 3H) in the ^1H -NMR spectrum, and by δ 199.8, 56.9, 42.3, 38.3, 32.6, 31.5, 14.7 in the ^{13}C -NMR spectrum.

4.5. General Procedure for Piperidinone Synthesis from the Radical Adduct 23

To a stirred solution of **23** (145 mg, 0.5 mmol) in dichloromethane (20 mL) at 0 °C, trifluoroacetic acid (5 mL) was added drop-wise. Stirring was maintained for 1 h before the acid was removed by azeotropic distillation with toluene (5 mL $\times$ 3), after which the residue was dried under vacuum. The residue was dissolved in DMF (20 mL) and cooled down to 0 °C before 2,4,6-collidine (0.090 mg, 0.75 mmol) was added drop-wise and the reaction mixture was stirred for 1 h at 0 °C. The resulting reaction mixture was maintained at 0 °C, and Cs_2CO_3 (0.195 mg, 0.6 mmol) followed by the sulfonamide (0.6 mmol) were added. The reaction mixture was warmed to room temperature and stirred for 1.5 h before the solvent was removed under vacuum and the residue was dissolved in EtOAc (30 mL). The organic layer was washed successively with water and brine, dried over Na_2SO_4 , concentrated and purified by silica gel column chromatography.

4.5.1. *N*-(2-Phenylethyl) (4-Methyl-2-oxopiperidin-3-yl)acetamide (**24**)—

Chromatographic purification eluting with 4% methanol in dichloromethane afforded a pale yellow solid in 78% yield as a 3:1 *cis*:*trans* mixture.

cis-Isomer: IR (film) 1650 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.30 (t, $J = 7.5$ Hz, 2H), 7.24-7.19 (m, 3H), 6.86 (br s, 1H), 5.78 (br s, 1H), 3.54-3.45 (m, 2H), 3.38 (dt, $J = 5.5, 11.5$ Hz, 1H), 3.32-3.23 (m, 1H), 2.84-2.79 (m, 2H), 2.77-2.70 (m, 1H), 2.28-2.21 (m, 1H), 2.09 (dd, $J = 4.5, 14.5$ Hz, 1H), 2.01-1.94 (m, 1H), 1.88-1.78 (m, 2H), 0.93 (d, $J = 7.0$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 174.7, 172.6, 139.4, 129.0, 126.5, 43.4, 40.9, 39.1, 36.4, 35.9, 30.6, 28.9, 13.9; ESIHRMS m/z calculated for $\text{C}_{16}\text{H}_{22}\text{N}_2\text{O}_2\text{Na}$ ($M + \text{Na}$) $^+$: 297.1579, found: 297.1591.

trans-Isomer: the minor *trans* isomer was identified in the mixture by characteristic signals at δ 6.50 (br s, 1H), 5.85 (br s, 1H), 2.50 (dd, $J = 6.0, 14.5$ Hz, 1H), 2.18-2.14 (m, 1H), 1.72-1.66 (m, 1H), 1.1 (d, $J = 6.5$ Hz, 3H) in the ^1H -NMR spectrum, and by δ 174.4, 171.9, 139.3, 129.0, 128.7, 46.6, 41.3, 40.7, 36.0, 35.8, 31.9, 30.7, 20.4 in the ^{13}C -NMR spectrum.

4.5.2. *N*-[(4-Methyl-2-oxopiperidin-3-yl)acetyl]piperidine (25**)**—Chromatographic purification eluting with 4% methanol in dichloromethane afforded the *cis* and *trans* isomers with yields of 56% and 19%, respectively.

cis-Isomer: light yellow gum, IR (film) 1635 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 5.63 (br s, 1H), 3.64-3.59 (m, 1H), 3.56-3.39 (m, 4H), 3.34-3.29 (m, 1H), 3.14-3.09 (m, 2H), 2.43-2.37 (m, 1H), 2.27 (dd, $J = 9.7, 17.2$ Hz, 1H), 2.09-2.03 (m, 1H), 1.75-1.52 (m, 7H), 0.96 (d, $J = 7.0$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 174.1, 170.0, 46.8, 43.1, 42.9, 38.9, 30.6, 29.3, 28.5, 26.7, 25.8, 24.9, 14.0; ESIHRMS m/z calculated for $\text{C}_{13}\text{H}_{22}\text{N}_2\text{O}_2\text{Na}$ ($M + \text{Na}$) $^+$: 261.1579, found: 261.1561.

trans-Isomer: light yellow gum, IR (film) 1646 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 5.72 (br s, 1H), 3.59-3.40 (m, 5H), 3.32-3.26 (m, 1H), 2.94 (dd, $J = 5.5, 16.5$ Hz, 1H), 2.70 (dd, $J = 3.2, 16.2$ Hz, 1H), 2.26-2.20 (m, 1H), 2.17-2.12 (m, 1H), 1.85 (dd, $J = 3.2, 13.2$ Hz, 1H),

1.68–1.50 (m, 7H), 1.02 (d, $J = 7.0$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 174.6, 169.6, 46.7, 45.6, 43.0, 41.2, 37.9, 32.1, 31.5, 31.1, 29.9, 26.6, 25.8, 24.8, 20.6; ESIHRMS m/z calcd for $\text{C}_{13}\text{H}_{22}\text{N}_2\text{O}_2\text{Na}$ ($\text{M} + \text{Na}$) $^+$: 261.1579, found: 261.1564.

Acknowledgments

We thank the NIH (GM57335 and GM62160) for support of this work.

4.7. References

1. Niyomura O, Kato S. *Top. Curr. Chem* 2005;251:1–12.
2. Kato S, Kawahara Y, Kageyama H, Yamada R, Niyomura O, Murai T, Kanda T. *J. Am. Chem. Soc* 1996;118:1262.
3. Hadad CM, Rablen PR, Wiberg KB. *J. Org. Chem* 1998;63:8668–8681.
4. Bauer, W.; Kühlein, K. *Methoden der Organischen Chemie*, 4th Ed; Carbonsäure und Carbonsäure Derivate. Falbe, J., editor. Stuttgart; Thieme: 1985. p. 832–890.
5. Kato, S.; Murai, T. *The Chemistry of Acid Derivatives*. Patai, S., editor. Wiley; Chichester: 1992. p. 803–847.
6. Scheithauer S, Mayer R. *Topics in Sulfur Chemistry* 1979;4:1–373.
7. Lu W, Qasim MA, Kent SBH. *J. Am. Chem. Soc* 1996;118:8518–8523.
8. Crich D, Sharma I. *Angew. Chem. Int. Ed* 2009;48:2355–2358.
9. Talan RS, Sanki AK, Sucheck SJ. *Carbohydr. Res* 2009;344:2048–2050. [PubMed: 19643396]
10. Messeri T, Sternbach DD, Tomkinson NCO. *Tetrahedron Lett* 1998;39:1673–1676.
11. Messeri T, Sternbach DD, Tomkinson NCO. *Tetrahedron Lett* 1998;39:1669–1672.
12. Crich D, Sasaki K. *Org. Lett* 2009;11:3514–3517. [PubMed: 19719195]
13. Kricheldorf HR, Leppert E. *Makromol. Chem* 1973;167:47–68.
14. Stockdill JL, Wua X, Danishefsky SJ. *Tetrahedron Lett* 2009;50:5152–5155. [PubMed: 20160954]
15. Wua X, Li X, Danishefsky SJ. *Tetrahedron Lett* 2009;50:1523–1525. [PubMed: 20161329]
16. Yuan Y, Zhu J, Li X, Wua X, Danishefsky SJ. *Tetrahedron Lett* 2009;50:2329–2333. [PubMed: 20161291]
17. Bao Y, Li X, Danishefsky SJ. *J. Am. Chem. Soc* 2009;131:12924–12926. [PubMed: 19697932]
18. Blake J. *Int. Pept. Prot. Res* 1981;17:273–274.
19. Yamashiro D, Blake JF. *Int. J. Pept. Prot. Chem* 1981;18:383–392.
20. Mitin YV, Zapevalova NP. *Int. J. Pept. Prot. Chem* 1990;35:352–356.
21. Høeg-Jensen T, Olsen CE, Holm A. *J. Org. Chem* 1994;59:1257–1263.
22. Kolakowski RV, Shangguan N, Sauers RR, Williams LJ. *J. Am. Chem. Soc* 2006;128:5695–5702. [PubMed: 16637636]
23. Kolakowski RV, Shangguan N, Williams LJ. *Tetrahedron Lett* 2006;47:1163–1166.
24. Shangguan N, Katukojvala S, Greenberg R, Williams LJ. *J. Am. Chem. Soc* 2003;125:7754–7755. [PubMed: 12822965]
25. Barlett KN, Kolakowski RV, Katukojvala S, Williams L. *J. Org. Lett* 2006;8:823–826.
26. Merkx R, Brouwer AR, Rijkers DTS, Liskamp RM. *J. Org. Lett* 2005;7:1125–1128.
27. Merkx R, van Haren MJ, Rijkers DTS, Liskamp RMJ. *J. Org. Chem* 2007;72:4574–4577. [PubMed: 17497928]
28. Rakotomanomana N, Lacombe J-M, Pavia A. *Carbohydr. Res* 1990;197:318–323.
29. Rosen T, Lico IM, Chu DTW. *J. Org. Chem* 1988;53:1580–1582.
30. McKervey MA, O'Sullivan MB, Myers PL, Green RH. *Chem. Comm* 1993:94–96.
31. Crich, D. *Organosulfur Chemistry: Synthetic Aspects*. Page, P., editor. Academic Press; London: 1995. p. 49–88.
32. Stacey FW, Harris JF. *React* 1963;13:150–376.
33. Weiewer M, Chaminade X, Bayon JC, Dunach E. *Eur. J. Org. Chem* 2007:2464–2469.

34. Weiwer M, Dunach E. *Tetrahedron Lett* 2005;47:287–289.
35. Crich D, Sana K. *J. Org. Chem* 2009;74:7383–7388. [PubMed: 19725529]
36. Crich D, Sana K, Guo S. *Org. Lett* 2007;9:4423–4426. [PubMed: 17900128]
37. Rao Y, Li X, Nagorny P, Hayashida J, Danishefsky SJ. *Tetrahedron Lett* 2009;50:6684–6686. [PubMed: 20160926]
38. Crich D, Rahaman MY. *J. Org. Chem* 2009;74:6792–6796. [PubMed: 19670833]
39. Crich D, Sasaki K, Rahaman MY, Bowers AA. *J. Org. Chem* 2009;74:3886–3893. [PubMed: 19385609]
40. Crich D, Bowers AA. *Org. Lett* 2007;9:5323–5325. [PubMed: 17979281]
41. Mapp AK, Ansari AZ, Ptashne M, Dervan PB. *Proc. Natl. Acad. Sci., USA* 2000;97:3930–3935. [PubMed: 10760265]
42. Mapp AK, Dervan PB. *Tetrahedron Lett* 2000;41:9451–9454.
43. Crich D, Sana K. *J. Org. Chem* 2009;74:3389–3393. [PubMed: 19388715]
44. Aksnes G. *Acta Chem. Scand* 1961;15:692–694.
45. Bacaloglu R, Blasko A, Bunton CA, Cerichelli G, Castaneda F, Rivera E. *J. Chem. Soc., Perkin 2* 1995:965–972.
46. Balasubramaniyan V, Tongare DB, Gosavi SS, Babar SM. *Proc. Ind. Acad. Sci., Chem. Sci* 1993;105:265–271.
47. Bourdel E, Doulut S, Jarretou G, Labbe-Jullie C, Fehrentz J-A, Doumbia O, Kitabgi P, Martinez J. *Int. J. Pept. Prot. Res* 1996;48:148–155.
48. Cunha S, Kascheres A. *J. Brazilian Chem. Soc* 2000;11:525–529.
49. Doulut S, Dubuc I, Rodriguez M, Vecchini F, Fulcrand H, Barelli H, Checler F, Bourdel E, Aumelas A, Lallement JC, Kitabgi P, Costentin J, Martinez J. *J. Med. Chem* 1993;36:1369–1379. [PubMed: 8496905]
50. Hudson RF, Chopard PA. *Helv. Chim. Acta* 1963;46:2178–2185.
51. Osuch C, Franz JE, Zienty FB. *J. Org. Chem* 1964;29:3721–3722.
52. Kates MJ, Schauble JH. *J. Org. Chem* 1995;60:6676–6677.
53. Kates MJ, Schauble JH. *J. Heterocycl. Chem* 1995;32:971–978.
54. Structures and stereochemistries were assigned based on a combination of one and two dimensional NMR experiments.
55. McGarvey GJ, Williams JM. *J. Am. Chem. Soc* 1985;107:1435–1438.
56. Yamamoto Y, Yamada J-i, Uyehara T. *J. Am. Chem. Soc* 1987;109:5820–5822.
57. Fleming I, Hill JHM, Parker D, Waterson D. *J. Chem. Soc., Chem. Commun* 1985:318–319.
58. Crump RANC, Fleming I, Hill JHM, Parker D, Reddy NL, Waterson D. *J. Chem. Soc., Perkin Trans. 1* 1992:3277–3294.
59. Paddon-Row MN, Rondan NG, Houk KN. *J. Am. Chem. Soc* 1982;104:7162–7166.
60. Zimmerman HE. *Acc. Chem. Res* 1987;20:263–268.
61. Davies HML, Hodges LM, Gregg TM. *J. Org. Chem* 2001;66:7898–7902. [PubMed: 11701056]
62. Mohrig JR, Rosenberg RE, Apostol JW, Bastienanssen M, Evans JW, Franklin SJ, Frisbie CD, Fu SS, Hamm ML, Hirose CB, Hunstad DA, James TL, King RW, Larson CJ, Latham HA, Owen DA, Stein KA, Warnet R. *J. Am. Chem. Soc* 1997;119:479–486.
63. Banfi L, Guanti G. *Tetrahedron: Asymmetry* 1999;10:439–447.
64. Davies SG, Garner AC, Long MJC, Morrison RM, Roberts PM, Savory ED, Smith AD, Sweet MJ, Withey JM. *Org. Biomol. Chem* 2005;3:2762–2775. [PubMed: 16032355]
65. Karabatsos GJ, Fenoglio DJ. *Top. Stereochem* 1970;5:167–203.
66. Hoffmann RW. *Chem. Rev* 1989;89:1841–1860.
67. This model assumes kinetic selectivity. We were unable to isolate the intermediate adducts before rearrangement to the thiazepinone system and so could not test for equilibration at that level. The cis- and trans-thiazepinones were somewhat resistant to equilibration under mild acid and basic conditions suggesting that at least equilibration did not occur after rearrangement.
68. Chatgililoglu C. *Acc. Chem. Res* 1992;25:188–194.

69. Giese B, Damm W, Wetterlich F, Zeitz H-G. *Tetrahedron Lett* 1992;1863–1866.
70. Hart DJ, Huang H-C. *Tetrahedron Lett* 1985;26:3749–3752.
71. Kopping B, Chatgililoglu C, Zehnder M, Giese B. *J. Org. Chem* 1992;57:3994–4000.
72. Durkin KA, Liotta DC, Rancourt J, Lavellée J-F, Boisvert L, Guindon Y. *J. Am. Chem. Soc* 1992;114:4912–4914.
73. Erdmann P, Schafer P, Springer R, Zeitz H-G, Giese B. *Helv. Chim. Acta* 1992;75:638–644.
74. Curran, DP.; Porter, NA.; Giese, B. *Stereochemistry of Radical Reactions*. Weinheim; VCH: 1996.
75. Equilibration of the isomers of 25 with sodium methoxide in hot methanol resulted in an equilibrium mixture favoring the trans-isomer indicating the kinetic nature of the products observed.

Scheme 1.

Formation of dihydro-3-(triphenylphosphoranylidene)-2,5-thiophendione

Scheme 2.

Model for Stereoselective Protonation en Route to the 1,4-Thiazepin-5-ones.

Scheme 3.
Radical Addition to Ethylidene Monothiosuccinic Anhydride

Scheme 4.
Synthesis of Piperidinonyl acetamides

Table 1
Olefination Reactions of Reagent 1

Entry	RCHO	Product	% Yield
1	PhCHO	 2	78
2	MeCHO	 3	76
3	C ₇ H ₁₅ CHO	 4	72
4	c-C ₃ H ₅ CHO	 5	84

Table 2
Reaction of Alkylidene Monothiosuccinic Anhydrides with Mercaptoamines and 2,4-Dinitrobenzenesulfonamides

Entry	Substrate	Aminothiols	Sulfonamide (Ar = C ₆ H ₃ (NO ₂) ₂)	Product	% Yield (cis:trans) ratio
1					72 (1:5)
2					76 (1:5)

Entry	Substrate	Aminothioliol	Sulfonamide (Ar = C ₆ H ₅ (NO ₂) ₂)	Product	% Yield (cis:trans) ratio
3					65 (1:4)
4					67 (1:4)

Entry	Substrate	Aminothiols	Sulfonamide (Ar = C ₆ H ₅ (NO ₂) ₂)	Product	% Yield (cis:trans) ratio
5					74 (1:1)
6					72 (1:1)
7					58 (1:1)

Entry	Substrate	Aminothiols	Sulfonamide (Ar = C ₆ H ₅ (NO ₂) ₂)	Product	% Yield (cis:trans) ratio
8					62 (1:1)
9					67 (2:5)
10					61 (2:5)

Entry	Substrate	Aminothiols	Sulfonamide (Ar = C ₆ H ₃ (NO ₂) ₂)	Product	% Yield (cis:trans) ratio
11					57 (2:3)
12					60 (2:3)