Published in final edited form as:

Tetrahedron Lett. 2010 December 15; 51(50): 6649–6650. doi:10.1016/j.tetlet.2010.10.075.

Convenient one-pot synthesis of *trans*-1,2-diaryloxiranes from the direct coupling of benzyl halides

Freeman M. Wong, Yee Man Chan, David X. Chen, and Weiming Wu*

Department of Chemistry and Biochemistry, San Francisco State University, San Francisco, CA 94132

Abstract

trans-1,2-Diaryloxiranes were conveniently prepared in an one-pot reaction by the direct coupling of benzyl halides in the presence of silver oxide and DMSO under mild conditions.

Epoxides or oxiranes are an important class of chemicals and their reactions and synthesis have been extensively discussed.^{1–3} In this report, we present an unusual direct coupling reaction of benzyl halides to form symmetrically substituted *trans*-1,2-diaryloxiranes in the presence of silver oxide (Ag₂O) and DMSO under mild conditions.

Previously we reported a convenient method for the conversion of benzyl halides to aldehydes and ketones in the presence of Ag₂O and pyridine N-oxide.⁴ When DMSO was employed instead of pyridine N-oxide as the source of oxygen, it was noted that reaction with benzyl bromide gave a mixture of products.⁴ An epoxide, *trans*-stilbene oxide, was isolated from the product mixture. Further experiments with DMSO as the solvent have led to the development of the current method. The overall reaction involves the unusual coupling of the benzylic carbon to form the *trans*-1,2-diaryloxirane as the main product.

The reactions were carried out in DMSO. One equivalent of silver oxide was utilized to facilitate the reaction. Silver oxide presumably assists the heterolysis of the carbon-halogen bond and also functions as a base. The reaction mechanism is yet to be illustrated and awaits further investigation. However, a small amount of cross-coupling product was isolated when *p*-chlorobenzaldehyde was added to the reaction of benzyl bromide with silver oxide and DMSO, indicating possible involvement of aldehyde as a reaction intermediate. The reaction does not work with benzyl bromide derivatives with strong electron-withdrawing groups such as nitro or cyano groups.

*Corresponding author. Tel.: 1-415-338-1436; fax: 1-415-338-2384; wuw@sfsu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The reactions were conveniently carried out by stirring the reaction mixture overnight at room temperature. Upon completion of the reaction, the resulting mixture was centrifuged, decanted, and concentrated to give the epoxides as reasonably pure crude products. The epoxides were further purified through flash chromatography using hexane (or a hexane/methylene chloride mixture) as solvent. As shown in Table 1, the epoxides were obtained in moderate yield. All epoxides were identified as the *trans*-isomer by their NMR spectra.⁵

The above results have demonstrated that benzyl bromide derivatives can be directly coupled to form *trans*-epoxide. The one-pot reaction is conveniently carried out and can be of interest if symmetrically substituted 1,2-diaryloxiranes are desired.

Experimental Details

All reagents were obtained from commercial sources and used without further purification. *Typical experimental procedure:* Silver oxide (0.68 g, 2.9 mmol) was added to a solution of benzyl bromide (0.50 g, 2.9 mmol) in anhydrous DMSO (5 mL) in a round-bottom flask. The resulting mixture was stirred overnight under nitrogen. The reaction mixture was centrifuged, decanted, and concentrated to give the crude product. The product was purified by flash chromatography using hexane as the solvent (0.14 g, 48%). The product was identified to be *trans*-stilbene oxide by comparison of its NMR spectrum and melting point to that of an authentic sample.⁵

Acknowledgments

This investigation was supported by the National Institutes of Health, MBRS SCORE Program – Grant #5 S06 GM52588. The NMR facility was funded by the National Science Foundation (DUE-9451624 and DBI 0521342). Initial experiments were carried out by Chi M. Ho, Q. Y. Rudy Wu, Peter R. Wu, and Lili Wong in our laboratory. We are indebted to Professor Ihsan Erden (SFSU) for helpful discussions.

References

1. Smith, MB.; March, J. *March's Advanced Organic Chemistry. Reactions, Mechanisms, and Structure*. 5. Wiley; New York: 2001.
2. Erden, I. Monocyclic Oxiranes and Oxirenes. In: Katritzky, AR.; Rees, CW.; Scriven, EFV., editors. *Comprehensive Heterocyclic Chemistry II*. Vol. 1A. Pergamon; New York: 1997.
3. Erden, I. Oxiranes and Oxirenes: Fused Ring Derivatives. In: Katritzky, AR.; Rees, CW.; Scriven, EFV., editors. *Comprehensive Heterocyclic Chemistry II*. Vol. 1A. Pergamon; New York: 1997.
4. Chen DX, Ho CM, Wu QYR, Wu PR, Wong FM, Wu W. *Tetrahedron Lett* 2008;49:4147–4148. [PubMed: 19756261]
5. Pouchert, CJ.; Behnke, J. *The Aldrich Library of ¹³C and ¹H FT-NMR Spectra*. Aldrich; Milwaukee: 1992.

Table 1

Coupling of benzyl bromide derivatives to epoxides

Entry	Benzyl bromide derivatives	Yield (isolated)
1	R = H	48%
2	R = <i>p</i> -Br	43%
3	R = <i>o</i> -Cl	48%
7	R = 2-Cl-4-F	52%
4	R = <i>p</i> -CO ₂ CH ₃	58%
5	R = <i>m</i> -CO ₂ CH ₃	57%
6	R = <i>p</i> -CF ₃	48%