Assembly and Function of Integrin Receptors Is Dependent on Opposing α and β Cytoplasmic Domains

Roger Briesewitz, Andreas Kern, and Eugene E. Marcantonio*

Departments of Pathology and Anatomy & Cell Biology, College of Physicians and Surgeons, Columbia University, New York, New York, 10032

Submitted February 27, 1995; Accepted May 26, 1995
Monitoring Editor: Erkki Ruoslahti

The membrane proximal regions of integrin α and β subunits are highly conserved in evolution. In particular, all integrin α subunits share the KXGFFKR sequence at the beginning of their cytoplasmic domains. Previous work has shown that this domain is important in integrin receptor assembly. Using chimeric integrin α and β subunits, we show that the native cytoplasmic domains of both subunits must be present for efficient assembly. Most strikingly, chimeric $\alpha 1$ and $\beta 1$ subunits with reciprocally swapped intracellular domains dimerize selectively into collagen IV receptors expressed at high levels on the surface. However, these receptors, which bind ligand efficiently, are deficient in a variety of post-ligand binding events, including cytoskeletal association and induction of tyrosine phosphorylation. Furthermore, deletion of the distal α cytoplasmic domain in the swapped heterodimers leads to ligand-independent focal contact localization, which also occurs in wild-type subunits when the distal cytoplasmic domain is deleted. These results show that proper integrin assembly requires opposed α and β cytoplasmic domains, and this opposition prevents ligand-independent focal contact localization. Our working hypothesis is that these two domains may associate during receptor assembly and provide the mechanism for integrin receptor latency.

INTRODUCTION

Integrin receptors each consist of an α and a β subunit, which associate to mediate interactions between cells and the extracellular matrix (ECM). Many integrin receptors involved in adhesion concentrate in specific regions within the plasma membrane after binding their ECM substrates. These specialized domains of the plasma membrane are recognized in interference reflection microscopy as dark areas and are called focal contact sites (Burridge *et al.*, 1988). The actin microfilament network is anchored at focal contact sites, which also contain other components of the cytoskeleton, such as vinculin, α -actinin, and talin (Chen *et al.*, 1985; Damsky *et al.*, 1985; Dejana *et al.*, 1988; Singer *et al.*, 1988). The latter two proteins have been shown to interact with the cytoplasmic domain of the

$\beta 1$ subunit in vitro, suggesting a direct interaction in vivo (Horwitz *et al.*, 1986; Otey *et al.*, 1990). Thus, integrins can provide a mechanical linkage between the ECM and the cytoskeleton.

In addition to integrin-dependent cytoskeletal organization, there are a number of signals generated after cells attach to a matrix via integrin receptors. All of these signals require the clustering of integrins through the binding to either antibody or multivalent ligands (e.g., Kornberg *et al.*, 1991; Schwartz *et al.*, 1991a). In several cell types, the Na/H antiport system is activated, leading to an increase in intracellular pH (Schwartz *et al.*, 1991b). A second signal that can occur is an increase in intracellular Ca^{2+} (Jaconi *et al.*, 1991; Schwartz *et al.*, 1993).

It has also become clear that the binding of integrins to ligands is associated with tyrosine phosphorylation of specific cytoplasmic proteins (Guan *et al.*, 1991; Burridge *et al.*, 1992). A phosphoprotein has been localized to focal adhesions in fibroblasts and has been

* Corresponding author: Department of Pathology, 630 W. 168th St., Black Bldg. 1427, New York, NY 10032.

termed focal adhesion kinase, or pp125^{FAK} (Schaller *et al.*, 1992). This intracellular kinase is phosphorylated rapidly upon integrin binding to ligand (Guan *et al.*, 1991), and is a major phosphoprotein in most adherent cells. Cobb *et al.* (1994) have shown that two Src-related kinases, pp60^{src} and pp59^{lyn}, associate in a stable manner with pp125^{FAK}. This interaction is dependent on the SH2 domain of Src (Cobb *et al.*, 1994; Xing *et al.*, 1994), and on the major autophosphorylation site of pp125^{FAK} (Schaller *et al.*, 1994). These results suggest that pp125^{FAK}, which is phosphorylated in response to integrin binding to ligand, may lead to activation of Src-related kinases. Phosphorylated FAK can interact with the Ras pathway via the binding of the adaptor protein Grb2, thus promoting growth (Schlaepfer *et al.*, 1994). In addition, Vuori and Ruoslahti (1994) have shown that insulin receptor substrate-1 associates with $\alpha v \beta 3$, and may also play a role in attachment-dependent growth.

All integrin subunits contain a large extracellular domain, a single transmembrane segment, and a carboxyl terminal cytoplasmic domain of varying length. In general, the cytoplasmic domains of both the α and β subunits are quite short, from 13 ($\alpha 1$) to 58 amino acids ($\beta 7$). Overall, the β cytoplasmic domains are more highly conserved among classes and between species than the α cytoplasmic domains. In particular, the $\beta 1$ cytoplasmic domain is highly conserved in both vertebrates and invertebrates (DeSimone and Hynes, 1988; Marcantonio and Hynes, 1988). Previous work has shown that this domain is required for integrin focal contact localization. There are three regions that seem to be crucial as identified by site-directed mutagenesis studies (Solowska *et al.*, 1989; Hayashi *et al.*, 1990; Marcantonio *et al.*, 1990; Reszka *et al.*, 1992). Furthermore, it has been shown using reporter constructs that the $\beta 1$ cytoplasmic domain itself is able to direct chimeric proteins to focal contacts (Geiger *et al.*, 1992; LaFlamme *et al.*, 1992). These findings suggest that the $\beta 1$ cytoplasmic domain is not only necessary, but may be sufficient to direct integrins to focal contact sites; however, integrin localization to focal contact sites *in vivo* is entirely ligand dependent (Dejana *et al.*, 1988; Singer *et al.*, 1988). In addition, LaFlamme *et al.* (1992) showed that addition of a soluble ligand was effective in promoting receptor movement to focal contact sites, even when this ligand is not incorporated into the matrix. One explanation for these results would be that occupation of the integrin by ligand leads to a change in the receptor, by which the β cytoplasmic domain becomes "active" or "unmasked."

It has been shown that the α cytoplasmic domain may play a role in the control of localization of integrin to focal contacts (Briesewitz *et al.*, 1993b; Ylänné *et al.*, 1993). Deletion of most of the $\alpha 1$ cytoplasmic domain (preserving the conserved GFFKR sequence)

leads to the localization of this integrin to focal contacts even on substrates that $\alpha 1 \beta 1$ does not bind (Briesewitz *et al.*, 1993b). Thus, loss of the α cytoplasmic domain leads to ligand-independent focal contact localization. These experiments suggest that in native integrins, the α cytoplasmic domain may mask the $\beta 1$ cytoplasmic domain, preventing ligand-independent focal contact localization.

The simplest mechanism by which an α cytoplasmic domain could influence the β cytoplasmic domain would be an association between the two cytoplasmic domains. Most models of integrin heterodimeric structures, however, show that only the amino terminal segments of the extracellular domains of α and β subunits are associated (e.g. Nermut *et al.*, 1988). We have designed experiments to test whether the specific α and β transmembrane and cytoplasmic segments are required for the normal assembly and function of integrins. These experiments involve replacement of the native transmembrane and cytoplasmic domains of $\alpha 1$ and $\beta 1$ subunits with their respective partners in a reciprocal swap. Chimeric α and β subunits were introduced into cell lines, which led to selective dimerization of these chimera and a lack of dimerization with endogenous integrins. Analysis of the function of the chimeras showed that the swapped integrins were defective in post-ligand binding events, despite the ability to effectively bind to ligand. Interestingly, removal of a portion of the α cytoplasmic domain sequence led to ligand-independent focal contact localization, whereas the intact domains did not have this property. This last finding exactly mimics the behavior of native integrins. These results show that the α and β cytoplasmic domains are involved in heterodimer formation, a result that implies that they may be associated during receptor assembly and function.

MATERIALS AND METHODS

Plasmid Construction

Restriction enzymes and T4 polynucleotide kinase were purchased from New England Biolabs (Beverly, MA), T4 DNA ligase was obtained from Life Technologies (Gaithersburg, MD), and *Taq* polymerase was obtained from Boehringer-Mannheim (Indianapolis, IN). Oligonucleotides were purchased from Operon Technologies (Alameda, CA) and were used directly without further purification.

For the construction of the chimeric cDNAs $\beta \times \alpha c$ and $\alpha \times \beta c$, fragments of the 3' end of the human $\alpha 1$ and chicken $\beta 1$ cDNA containing the transmembrane domains were subcloned into pBlue-script (Stratagene, La Jolla, CA). Using polymerase chain reaction (PCR), fragments were produced that included the transmembrane and cytoplasmic domains of $\alpha 1$ and $\beta 1$. In addition, fragments of the extracellular domains of $\alpha 1$ and $\beta 1$ were prepared by PCR. Extracellular fragments were connected to intracellular ones by blunt ligation, and the chimeric cDNA fragments were then exchanged with the corresponding 3' sequences of the full length $\alpha 1$ and chicken $\beta 1$ sequences in the expression vector pLEN (Marcantonio *et al.*, 1990). The oligonucleotides that define the junction of the chimeric sequences are listed below. The numbers assigned to the oligonucleotides correspond to the sequence position of the 5'

nucleotides according to Briesewitz *et al.* (1993a) for $\alpha 1$ and Tamkun *et al.* (1986) for chicken $\beta 1$. The letters "A" and "S" denote "anti-sense" and "sense" orientation of the primer.

$\alpha\beta c$: $\alpha 1$ 3429A 5' TGG CAC TCT GCC CGG TA 3' $\beta 1$ 2296S 5' ATC ATA CCC ATT GTA GCT 3'

βxac : $\beta 1$ 2295A 5' GTC AGG GCC ACT AGG GCA 3' $\alpha 1$ 3430S 5' TTA TGG CTC ATC CTG CT 3'

The corresponding protein sequences for the junctions of these two chimeric subunits are:

$\alpha\beta c$: NH₂:... P-G-R-V-P-[underline]I-I-P-I-V-A... .COOH $\alpha 1$ (extracellular) ($\beta 1$ transmembrane)

βxac : NH₂:... C-P-S-G-P-D-L-W-V-I-L COOH $\beta 1$ (extracellular) ($\alpha 1$ transmembrane)

The chimeric cDNA βxac was made via PCR using as a template the βxac fragment cloned into pBluescript. An oligonucleotide was designed that contained a termination codon at its 5' end and primes in the anti-sense direction at the GFFKR sequence. Using the T7 primer of the pBluescript vector, a PCR fragment was made that encompassed the chimeric sequence βxac , but lacking the $\alpha 1$ cytoplasmic sequences 3' of the GFFKR. The PCR fragment was subsequently subcloned into pLEN $\beta 1$.

For the construction of the chimeric cDNA $\alpha T\beta$, the method of overlapping PCR was used (Horton *et al.*, 1990). This fragment was cloned into pBluescript first, and then subcloned into pLEN $\alpha 1$.

The oligonucleotides used were:

for $\alpha 1$ -Tac: $\alpha 1$ 3429A 5' TGG CAC TCT GCC CGG TA 3' Tac 892S 5' GGC AGA GTG CCA CAG GTA GCA GTG GCC GG 3'

for $\beta 1$ -Tac: $\beta 1$ 2365S 5' AAA CTA CTG ATG ATC AT 3' Tac 960A 5' CAT CAG TAG TTT CTG CCA GGT GAG CCC ACT 3'

The corresponding protein sequences for the junctions of these chimeric subunits are:

$\alpha T\beta$: NH₂ P-G-R-V-P-Q-V-A-V-A L-T-W-Q-K-L-L-M-I COOH $\alpha 1$ (extracellular) IL2R (transmembrane) $\beta 1$ cytoplasmic

The full length cDNA for the interleukin-2 receptor α subunit (Giordano *et al.*, 1991) was kindly provided by Dr. T. Giordano (Abbott Labs, Abbott Park, IL).

The fidelity of all regions amplified by PCR was verified by sequencing.

DNA Transfections

NIH 3T3 cells were maintained in DME with 10% calf serum (CS). Cells (5×10^5), plated the previous day, were co-transfected with 10–20 μ g of the various chimeric cDNAs in the pLEN expression vector system (Marcantonio *et al.*, 1990) and 2 μ g pSV2neo (Southern and Berg, 1982) for the primary transgene or pSV HIS-D (Hartman and Mulligan, 1988) for the second transgene in cells already neo-resistant. The calcium phosphate transfection method was used as previously described (Solowska *et al.*, 1989). After 3 days, the cells were split into DME with 10% CS containing 1 mg/ml G418 (Geneticin; Life Technologies) in the case of the first transgene or 3 mM histidinol (Sigma Chemical, St. Louis, MO)/0.5 mg/ml G418 for the second transgene. After 10–14 days, resistant clones were isolated and screened by immunoprecipitation. For each construct, multiple clones were isolated and characterized. Although the data shown are usually from one clone, similar results have been seen with the other clones as well.

Antibodies

Rabbit anti- $\beta 1$ cytoplasmic domain antibodies and rabbit anti- $\alpha 5$ cytoplasmic domain antibodies were prepared as described (Marcantonio and Hynes, 1988; Hynes *et al.*, 1989). CSAT monoclonal antibody was prepared from hybridomas. Polyclonal rabbit anti-chicken $\beta 1$ was prepared as described previously (Marcantonio *et al.*, 1990). Monoclonal anti-human $\alpha 1$ (TS2/7) was a gift of T. Springer (Harvard Medical School, Cambridge, MA). The rabbit anti-talin serum was a kind gift of K. Burrige (University of North Carolina, Chapel Hill, NC). Rabbit anti- $\alpha 1$ cytoplasmic domain

antibody was a gift of G. Tarone (University of Torino, Torino, Italy). Rabbit anti-phosphotyrosine was a gift of J. Sap (New York University, NY). Monoclonal anti-human $\alpha 1$ 1B3.1 was prepared as described (Bank *et al.*, 1989). Rabbit inhibitory anti-rat $\beta 1$ (Lenny) antibodies were provided by C. Buck (Wistar Institute, Philadelphia, PA) (Albelda *et al.*, 1989).

Radiolabeling and Immunoprecipitation

Cells were labeled with Na¹²⁵I (Dupont NEN, Boston, MA) and lactoperoxidase as a monolayer as described (Hynes, 1973). 10^7 cells and 1–2 mCi/ml were used per experiment. The cells were extracted with 0.5% NP-40 and immunoprecipitated as described (Marcantonio and Hynes, 1988).

For pulse-chase analysis, 3T3 cells were incubated for 1 h in methionine-free DME plus 10% FCS, followed by incubation in the same medium plus 10% FCS containing 500 μ Ci/ml of [³⁵S]methionine (Translabel; ICN Radiochemicals, Irvine, CA). After 1 h, the radioactivity was removed, an equal volume of growth media was added, and the cells were incubated for the chase times listed in the figure legends. After the chase, the cells were washed three times with cold phosphate-buffered saline (PBS) followed by extraction with 0.5% NP-40 and immunoprecipitation as described (Marcantonio and Hynes, 1988). Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was performed by the method of Laemmli (1970). Separation gels were 7% acrylamide with a 3% stacking gel. Samples were prepared in sample buffer (5% SDS, 100 mM Tris-HCl, pH 6.8, 10 mM EDTA, 10% glycerol, and bromophenol blue) and boiled for 3 min.

Affinity Chromatography

The CB3 fragment of human collagen type IV was prepared and coupled to Sepharose CL4B (Pharmacia Biotech, Piscataway, NJ) as described (Vandenberg *et al.*, 1991). 3T3 cell lines were surface labeled with ¹²⁵I as described above. Cell extracts were prepared using 50 mM β octyl-glucoside (BOG) in either 50 mM Tris, pH 7.5, 150 mM NaCl, 1 mM MgCl₂, 1 mM MnCl₂ (TMM), or 50 mM Tris, pH 7.5, 150 mM NaCl, 1 mM MgCl₂ (TM). After a 10-min incubation on ice, detergent insoluble material was removed by centrifugation at $10,000 \times g$ for 20 min. Four-fifths of the resulting supernatant was incubated batch-wise with 100 μ l of packed beads (CB3-Sepharose) for 1 h at 4°C with end-over-end mixing. The beads were recovered by sedimentation, and washed four times with 50 mM BOG in TMM or TM, followed by elution with 1 ml of 50 mM BOG in TMM containing 20 mM EDTA. This eluate was concentrated by immunoprecipitation with rabbit anti- $\beta 1$ cytoplasmic peptide serum and compared with an immunoprecipitate of one-tenth of the cleared extract, followed by SDS-PAGE analysis.

Immunofluorescence

Solutions of 0.5 μ g of either fibronectin (FN) or the CB3 fragment of human collagen type IV in 70 μ l of water were air dried onto coverslips. After drying, the proteins were rehydrated in PBS for 30 min. Cells were treated with 25 μ g/ml cycloheximide for 2 h before trypsinization. The trypsinized cells were washed twice in PBS and plated onto the coated glass coverslips in serum-free media plus cycloheximide. The cells were then placed in the tissue culture incubator at 37°C for 3 h before fixing them in 4% paraformaldehyde in PBS for 5 min and permeabilizing them with 0.2% NP-40 in PBS for 5 min afterward. The antibody against talin was diluted 1:400 in 10% normal goat serum in PBS. TS2/7 hybridoma supernatant was supplemented with normal goat serum (10% final), and then used directly. The coverslips were overlaid with 20 μ l of primary antibody solution and incubated at 37°C for 30 min. After washing with PBS, the coverslips were incubated with 20 μ l of secondary antibody for 30 min at 37°C. The secondary antibodies were F(ab')₂ fragments of rhodamine-conjugated rabbit anti-mouse

IgG and fluorescein-conjugated goat anti-rabbit IgG (Cappel, Durham, NC), diluted 1:250 in 10% normal goat serum in PBS. Rhodamine-labeled phalloidin was diluted 1:500 in 10% normal goat serum in PBS and incubated with fixed and permeabilized cells for 30 min at 37°C.

Cell Adhesion Assays

Adhesion assays were done essentially as described (Vandenberg *et al.*, 1991). Adhesive proteins were diluted in 0.1 M acetic acid (Col IV) or distilled water (FN) and allowed to dry in microtiter wells at room temperature. After washing with PBS, remaining protein binding sites were blocked with bovine serum albumin (BSA; 1% in PBS). Cells were resuspended in serum-free DME ($4-5 \times 10^5$ /ml) and incubated in the wells for 30 min at 37°C. Nonadherent cells were removed by washing with PBS, and bound cells were fixed with 70% ethanol and stained with crystal violet (0.1% in distilled water). Excess stain was removed with water. Bound stain was dissolved with Triton X-100 (0.2% in distilled water) and the optical density was read at 595 nm. All experiments were done in triplicate.

For inhibition assays, cells were pre-incubated for 10 min at room temperature in DMEM with dilutions of inhibitory antibodies as indicated in the legend of Figure 4 and then transferred to wells coated with collagen IV (0.1 μ g/well). After 30 min at 37°C, non-adherent cells were removed and bound cells were fixed and detected as described above. All experiments were done in triplicate.

For adhesion-dependent phosphorylation experiments, 3T3 cells were treated for 2 h with 25 μ g/ml of cycloheximide. After trypsinization, cells were washed in PBS with 1 mg/ml of Soybean trypsin inhibitor (Sigma Chemical), followed by two washes in DMEM. The cells were then plated in the absence of serum in 1 mg/ml of heat-treated BSA at 37°C for 1 h on dishes previously coated with 10 μ g/ml of FN or 1 μ g/ml of collagen IV. Suspension controls were performed by incubating the cells in the presence of BSA but on uncoated plastic, to which the cells did not attach.

Western Blot Analysis

Cells were washed with PBS, lysed in 300 μ l of sample buffer containing 1 mM sodium orthovanadate, and then boiled for 15 min. Samples were normalized by the OD²⁶⁰, then electrophoresed on 7% gels and transferred to nitrocellulose (Life Technologies) (Towbin *et al.*, 1979). After blocking with Superblock (Pierce Chemical, Rockford, IL), phosphotyrosine-containing bands were stained with a 1:2000 dilution of rabbit anti-phosphotyrosine, followed by a 1:10000 dilution of protein A-peroxidase (Boehringer Mannheim, Indianapolis, IN). Bands were visualized using enhanced chemiluminescence (ECL; Amersham, Arlington Heights, IL) followed by exposure to film (1 min).

Flow Cytometry

Fluorescence flow cytometric analysis was performed on a FACScan cytofluorometer (Becton Dickinson, Mountain View, CA). The cells were trypsinized and washed with growth media. The cells were resuspended in PBS with 0.5% CS and 0.5% NaN₃ using 5×10^5 cells in 50 μ l. Staining was performed in primary antibody on ice for 30 min, followed by three washes with cold PBS. Cells were then resuspended and stained with FITC-conjugated goat anti-mouse antibody (Tago, Burlingame, CA) 1:100 dilution for 30 min. After three washes, cells were analyzed.

RESULTS

Requirement of Integrin α and β Intracellular Domains for Subunit Assembly

We have previously compared the assembly of soluble, truncated versions of the integrin $\alpha_1\beta_1$ with native

heterodimers in 3T3 cells (Briesewitz *et al.*, 1993a). Although the intact human α_1 subunit can associate with native mouse β_1 subunit (Briesewitz *et al.*, 1993a), and the intact avian β_1 subunit can interact with the native mouse α subunits (Solowska *et al.*, 1989; Hayashi *et al.*, 1990), the truncated forms of the human α_1 and the avian β_1 do not interact with the endogenous subunits, but instead preferentially interact with each other. Because the dimerization of truncated α and β is independent intracellularly from the native α and β , one possibility is that the transmembrane and cytoplasmic regions of α and β may play a role in dimerization of intact integrin subunits. Consistent with this hypothesis, the α and β cytoplasmic domains are highly conserved. In particular, all integrin α subunit cytoplasmic domains contain the sequence KXGFFKR in the membrane proximal region, which suggests that this sequence has an important general function.

We have designed experiments to test the role of these domains in the normal assembly and function of integrin receptors. Previous work has shown that the sequences distal to the GFFKR in α cytoplasmic domains are not required for assembly with endogenous native β subunits, because deletion of those sequences within α subunits has no effect on assembly (Briesewitz *et al.*, 1993b; Kawaguchi and Hemler, 1993; Ylännä *et al.*, 1993). To test the role of the α intracellular domain in the normal assembly and surface expression of heterodimers, we connected the transmembrane and cytoplasmic domains of the chicken β_1 subunit with the extracellular domain of the human α_1 subunit ($\alpha\beta_1$) and introduced it into NIH 3T3 cells. When the resultant lines were analyzed by pulse-chase, the chimeric subunit was synthesized but degraded, and did not assemble efficiently with endogenous subunits (Figure 1A). Furthermore, this individual subunit is not detectable on the surface in ¹²⁵I surface-labeling experiments (our unpublished results) and only small amounts may be seen with flow cytometry (Table 1). Thus the α intracellular domain is required for assembly of $\alpha_1\beta_1$ heterodimers.

To test the role of the β intracellular domain in the normal assembly and surface expression of heterodimers, we connected the transmembrane and cytoplasmic domains of the human α_1 subunit with the extracellular domain of the chicken β_1 subunit ($\beta\alpha_1$) and introduced it into NIH 3T3 cells. A similar chicken β_1 subunit contained a C-terminal deletion in the α_1 cytoplasmic domain distal to the KXGFFKR region ($\beta\alpha_{1\text{cyt}}$) (see also Figure 2). These chimeric cDNAs were introduced separately into NIH 3T3 cells and analyzed by pulse chase. Cells were labeled metabolically and chased for the times indicated. The immunoprecipitation of $\beta\alpha_1$ with an antibody against chicken β_1 showed that the chimeric β subunit does not dimerize with any α subunits (Figure 1B). Moreover, this protein does not mature, but appears to be

Figure 1. (A) The chimeric integrin subunit $\alpha_x\beta_c$ does not dimerize with native integrin β_1 subunits. NIH 3T3 cells expressing the chimeric subunit $\alpha_x\beta_c$ were labeled with [^{35}S]methionine for 1 h then lysed immediately (lane 1 and 4) or chased with fresh media for 2 h (lane 2 and 5) or 6 h (lane 3 and 6) before lysis. The cell lysates were immunoprecipitated with a rabbit anti- β_1 cytoplasmic peptide serum (lanes 1–3) or the mouse monoclonal anti-human α_1 antibody

Table 1. Surface expression levels of exogenous integrin subunits

Cell line (exogenous subunits)	Mean fluorescence (TS2/7)	Mean fluorescence (CSAT)
HA1 (α_1)	206	8
HA1 β_{av} (α_1/β_{av})	192	512
$\alpha_x\beta_c$ #3 ($\alpha_x\beta_c$)	19	7
$\beta_{x\alpha c}$ #13 ($\beta_{x\alpha c}$)	11	9
SW1 ($\alpha_x\beta_c/\beta_{x\alpha c}$)	271	204
SWC ($\alpha_x\beta_c/\beta_{x\alpha c}$)	213	243
ATB ($\alpha_T\beta/\beta_{x\alpha c}$)	308	217
3T3 (none)	13	11

Fluorescence flow cytometric analysis of cell lines stained with mouse anti-human α_1 antibody (TS2/7) and mouse anti-chicken β_1 (CSAT) followed by FITC-conjugated goat anti-mouse antibody.

retained in the endoplasmic reticulum in a stable manner. Furthermore, no $\beta_{x\alpha c}$ is detected in flow cytometry experiments (Table 1). Hence, the β_1 intracellular domain is required for the assembly with α subunits. Based on these results, we hypothesize that normal integrin heterodimer formation and function may require associations between both extracellular and intracellular domains.

Expression of Chimeric Integrin Heterodimers in Mouse NIH 3T3 Cells

The involvement of integrin intracellular domains in assembly suggests that chimeric α/β subunits containing "swapped" intracellular domains may dimerize selectively. If the wild-type human α_1 subunit is co-expressed together with the wild-type chicken β_1 subunit in NIH 3T3 cells (HA1 β_{av}), the resulting cell lines have a complex integrin profile that requires the use of species-specific antibodies to identify individual sets of heterodimers. Figure 2A shows the results of an immunoprecipitation of an extract derived from ^{125}I surface-labeled HA1 β_{av} cells. Immunoprecipitation with an anti-chicken β_1 antibody shows chicken β_1 in complex with both the endogenous α_5 and exogenous α_1 subunits, with no evidence of a selective

Figure 1 cont. TS2/7 (lanes 4–6). The diagram below the autoradiographs represents the exogenous α subunit expressed in the cells. (B) Biosynthesis and assembly of the native (mouse) β_1 subunit and the chimeric (chicken) β_1 subunit $\beta_{x\alpha c}$. NIH 3T3 cells expressing the chimeric chicken β_1 subunit $\beta_{x\alpha c}$, were labeled with [^{35}S]methionine for 1 h, and then lysed directly (0 h) or chased for the times (h) indicated. The lysates were immunoprecipitated with either a rabbit anti- β_1 cytoplasmic peptide serum, specific for the endogenous mouse β_1 subunit ($\beta_1 m$), or a rabbit anti-chicken β_1 serum, specific for the chimeric β_1 subunit $\beta_{x\alpha c}$ ($\beta_1 ch$). The immunoprecipitates were recovered with protein A-Sepharose. The chimeric β_1 subunit $\beta_{x\alpha c}$ is retained and does not assemble with endogenous α subunits.

Figure 2. The chimeric integrin subunit $\alpha\beta_c$ selectively dimerizes with the chimeric subunit $\beta_x\alpha_c$. In contrast, human α_1 and chicken β_1 assemble with each other and with endogenous subunits. (A) HA1 β_{av} cells coexpressing the human α_1 subunit and the chicken β_1 subunit were ^{125}I surface labeled, lysed, and immunoprecipitated with a rabbit anti- β_1 cytoplasmic peptide serum (lane 1), the mouse monoclonal anti-chicken β_1 antibody CSAT (lane 2), or the mouse monoclonal anti-human α_1 antibody TS2/7 (lane 3). (B) SW1 cells expressing the chimeric subunits $\alpha_x\beta_c$ and $\beta_x\alpha_c$ were ^{125}I surface labeled, lysed, and immunoprecipitated with a rabbit anti- β_1 cytoplasmic peptide serum (lane 4), a rabbit anti-chicken β_1 serum (lane 5), the mouse monoclonal anti-chicken β_1 antibody CSAT (lane 6), the mouse monoclonal anti-human α_1 antibody TS2/7 (lane 7), or a rabbit anti- α_1 cytoplasmic peptide serum (lane 8). (C) SWC cells expressing the chimeric subunits $\alpha_x\beta_c$ and $\beta_x\alpha_{cyt}$ were ^{125}I surface labeled, lysed, and immunoprecipitated with the mouse monoclonal anti-chicken β_1 antibody CSAT (lane 9), a rabbit anti- α_5 serum (lane 10), and the monoclonal anti-human α_1 antibody TS2/7 (lane 11). The diagram below the autoradiographs represents the exogenous subunits expressed in the cells.

assembly (Figure 2A, lane 2). Furthermore, the adhesion of these cells to collagen IV (which is entirely α_1 dependent in 3T3 cells; Briesewitz *et al.*, 1993b) involves both mouse and chicken β_1 containing heterodimers as shown by inhibition studies (see below; Figure 4). Hence, exogenous wild-type human α_1 and chicken β_1 subunits assemble with each other as well as with endogenous mouse subunits.

To test the role of opposing α and β cytoplasmic domains in the assembly of integrins, we have coexpressed the two chimeric subunits described above, $\alpha_x\beta_c$ and $\beta_x\alpha_c$. In contrast to the expression of the chimera as single subunits in 3T3 cells, when $\alpha_x\beta_c$ and $\beta_x\alpha_c$ were coexpressed in NIH 3T3 cells, they selectively dimerized with each other. This cell line was named SW1. In addition, the chimeric subunits $\alpha_x\beta_c$ and $\beta_x\alpha_{cyt}$ were also coexpressed in 3T3 cells, and the cell line studied was named SWC. Figure 2 shows the results of an immunoprecipitation of extracts derived from ^{125}I surface-labeled SW1 (Figure 2B) and SWC (Figure 2C) cells. Immunoprecipitation of both extracts with an anti-chicken β_1 antibody or anti- α_1 antibody shows $\beta_x\alpha_c$ (or $\beta_x\alpha_{cyt}$) in complex with $\alpha_x\beta_c$ subunits, with no evidence of assembly with endogenous subunits. The relative amounts of the major FN receptor $\alpha_5\beta_1$ in these cells are similar (Figure 2B, lane 4, and Figure 2C, lane 10). In contrast to the individual chimera-containing cell lines, the surface expression of $\alpha_x\beta_c/\beta_x\alpha_c$ and $\alpha_x\beta_c/\beta_x\alpha_{cyt}$ dimers was comparable to that of native integrins (Table 1).

Thus, we have created an integrin collagen IV receptor in NIH 3T3 cells whose subunits dimerize specifically.

Because the chimeric subunits contain swaps of both transmembrane and cytoplasmic domains, we also wished to test the role of the β transmembrane domain in the assembly of chimeric integrin subunits. We attached the transmembrane domain of the interleukin-2 receptor α subunit to the α_1 extracellular domain and the β_1 intracellular domain, creating the chimeric subunit $\alpha\text{T}\beta$, and expressed this mutant subunit with $\beta_x\alpha_{cyt}$. Like SW1 and SWC cells, the $\alpha\text{T}\beta/\beta_x\alpha_{cyt}$ cells show selective and stable dimerization of chimeric subunits (our unpublished results). The surface content of the heterodimer is high (Table 1). These results indicate that the β transmembrane region is not required for assembly in this system. All chimeric heterodimers that are efficiently expressed on the cell surface contain the KXGFFKR sequence and the β_1 cytoplasmic domain. Thus, both these domains are required during biosynthesis.

Adhesive Properties of Cells Expressing Chimeric Integrins

If the chimeric collagen IV receptors were functional, they would provide a method for performing structure-function studies of the exogenous β_1 subunit without concern for the role of endogenous β_1 subunits, because these subunits would not be involved in the formation of the collagen IV receptor. To study the

function of the chimeric integrins, we examined the adhesion of SW1 and SWC cells to FN and collagen IV. The adhesive properties of these two cell lines were compared with those of NIH 3T3 cells transfected with the wild-type human $\alpha 1$ subunit, HA1 cells (Briesewitz *et al.*, 1993a,b). Normal, untransfected NIH 3T3 cells adhere to fibronectin through $\alpha 3\beta 1$ and $\alpha 5\beta 1$ receptors, and show little adhesion to collagen IV (Briesewitz *et al.*, 1993b). In all cases, heterologous expression of chimeric subunits in NIH 3T3 cells does not result in a major change in the adhesion to FN (Briesewitz *et al.*, 1993b and our unpublished results). The adhesion of these three NIH 3T3 lines to collagen IV, however, is different as shown in Figure 3A. The expression of the wild-type human $\alpha 1$ subunit in HA1 cells results in effective adhesion to collagen IV (Briesewitz *et al.*, 1993b). SW1 and SWC cells also adhere to collagen IV, but their adhesion is reduced as compared with HA1 cells, despite a high level of $\alpha x\beta c/\beta x\alpha c$ and $\alpha x\beta c/\beta x\alpha c$ on their surfaces (Table 1).

To understand the relative contribution of $\alpha 1\beta 1$ heterodimers in the adhesion to collagen IV, inhibition studies were performed (Figure 4). The adhesion of HA1 βav cells (containing wild-type versions of the human $\alpha 1$ subunit and chicken $\beta 1$ subunits) to collagen IV is blocked by either the inhibitory anti-human $\alpha 1$ monoclonal 1B3.1 or a combination of the anti-mouse $\beta 1$ polyclonal serum "Lenny" and the inhibitory anti-chicken $\beta 1$ monoclonal CSAT antibodies.

There is a partial block with the anti-chicken $\beta 1$ antibodies alone, showing that collagen IV adhesion is due to a mixture of heterodimers containing the $\alpha 1$ subunit with either the mouse or chicken $\beta 1$ as a partner. In contrast, both SW1 and SWC cells are effectively blocked by the chicken $\beta 1$ or anti- $\alpha 1$ antibodies, and do not require anti-mouse $\beta 1$ for inhibition. Because all of the collagen IV adhesion is blocked by either the inhibitory anti-human $\alpha 1$ monoclonal 1B3.1 or by the inhibitory anti-chicken $\beta 1$ monoclonal CSAT, we conclude that the adhesion of SW1 and SWC cells to collagen IV is mediated by human $\alpha 1$ and chicken $\beta 1$ containing chimeric subunits, again demonstrating the selectivity of dimerization of the exogenous subunits.

The reduced adhesion of the SW1 and SWC cells to collagen IV could be due to reduced receptor-ligand binding or a defect in post-ligand binding events leading to reduced adhesion. To distinguish between these possibilities, we compared the binding properties of the SW1 and HA1 integrins by affinity chromatography. ^{125}I surface-labeled extracts of HA1 and SW1 cells were incubated with the CB3 fragment of collagen-type IV-Sepharose, washed, and eluted with EDTA (Figure 3B). The affinity binding experiments were performed under two different conditions, either with Mn^{2+} (Figure 3B, lanes 3 and 7) or with Mg^{2+} (Figure 3B, lanes 4 and 8). Both the wild-type and chimeric $\alpha 1\beta 1$ bound to CB3-Sepharose in a Mg^{2+} -dependent manner. The binding was enhanced by the presence of

Figure 3. Analysis of adhesion and binding of chimeric integrins to collagen IV. (A) Increasing amounts of collagen IV were dried down onto microtiter plates. HA1 cells (open circles), SWC cells (filled circles), SW1 cells (open squares), and 3T3 cells (open triangles) were plated, incubated for 30 min at 37°C, and washed. The bound cells were fixed and stained with crystal violet. After washing, the stain was dissolved with Triton X-100 and the optical density was read at 595 nm. (B) Extracts of ^{125}I surface labeled HA1 cells (lanes 1–4) or SW1 cells (lanes 5–8) were prepared using β -octyl-glucoside (see MATERIALS AND METHODS). These extracts were immunoprecipitated with a rabbit anti- $\beta 1$ cytoplasmic domain peptide serum (lanes 1 and 5) or the mouse monoclonal anti-human $\alpha 1$ antibody TS2/7 (lanes 2 and 6). Alternatively, portions of the extracts were incubated with CB3-Sepharose either in a buffer containing Mn^{2+} (lanes 3 and 7) or in a buffer containing Mg^{2+} (lanes 4 and 8), washed, and eluted with 1 ml of buffer containing 10 mM EDTA. The eluates were then immunoprecipitated with the rabbit anti- $\beta 1$ cytoplasmic domain peptide serum using protein A-Sepharose.

Figure 4. Inhibition studies of collagen IV adhesion of 3T3 lines transfected with $\alpha 1$ and chicken $\beta 1$ constructs. Microtiter well surfaces were coated with collagen IV (0.05 $\mu\text{g}/\text{well}$ [HA1, HA1 βav] or 0.1 $\mu\text{g}/\text{well}$ [SW1 or SWC]). Cells were incubated for 30 min at 37°C in the presence of anti-human $\alpha 1$ mAb 1B3.1 (black bars, final dilution 1:200), anti-chicken $\beta 1$ mAb CSAT (hatched bars, final dilution 1:5), a mixture of anti-chicken $\beta 1$ mAb CSAT and rabbit anti-rat $\beta 1$ serum Lenny (black hatched bars, 1:100), or in the absence of antibodies (open bars). Cells were fixed and stained with crystal violet. Optical density was read at 595 nm and compared with the optical density of cells bound to an excess of FN (1 $\mu\text{g}/\text{well}$).

Mn^{2+} in both cases; however, both $\alpha 1$ -containing integrins were capable of binding in solution in the lower affinity state (Kern *et al.*, 1993) produced when the experiment was performed in Mg^{2+} . These results suggest that the observed adhesion defect in SW1 and SWC cells is based on post-ligand binding events, because binding to ligand is not affected. However, we cannot rule out in these experiments that some of the adhesive differences could be due to a modest decrease in affinity for ligand.

Cellular Localization of Integrin Receptors

Further evidence for post-ligand defects within the cell lines containing chimeric integrins was seen in the cellular localization of the receptors when the cells were adherent to collagen IV (Figure 5). When HA1 cells containing wild-type human $\alpha 1$ are plated on collagen IV, the human $\alpha 1$ subunit (Figure 5b) localizes to focal contact sites, as shown by co-localization with the cytoskeletal protein talin (Figure 5c). In addition, the actin cytoskeleton is organized into stress fibers, as shown by phalloidin staining (Figure 5a). In contrast, both the SW1 and SWC cell lines have few detectable stress fibers (Figure 5, d and g) and they do not form $\alpha 1$ -containing focal adhesions (Figure 5, e, f, h, and i). Thus, there is poor cytoskeletal organization in the SW1 and SWC cells when plated on collagen.

Integrin-Mediated Signaling

In addition to cytoskeletal organization, another critical post-ligand event in the function of integrins is the stimulation of the tyrosine kinase focal adhesion ki-

nase, or pp125^{FAK} (Guan *et al.*, 1991). This intracellular kinase is phosphorylated rapidly upon integrin binding to ligand. We have determined the amount of adhesion-dependent increase of tyrosine phosphorylation by performing Western blots using anti-phosphotyrosine antibodies of extracts of cells either maintained in suspension, or plated on FN or collagen IV. HA1 cells (Figure 6) showed a dramatic increase in tyrosine phosphorylation as compared with suspension controls (Figure 6, lane 1) when plated on FN (and adhering via endogenous $\alpha 3\beta 1$ and $\alpha 5\beta 1$; Figure 6, lane 2) or collagen IV (and adhering via transfected $\alpha 1\beta 1$; Figure 6, lane 3). Both SW1 (Figure 6, lane 5) and SWC cells (Figure 6, lane 8) show increased tyrosine phosphorylation when plated on FN similar to that seen in HA1 cells (Figure 6, lane 2). In contrast, SW1 (Figure 6, lane 6) and SWC (Figure 6, lane 9) have less tyrosine phosphorylation when plated on collagen IV as compared with HA1 cells (Figure 6, lane 3). Thus the chimeric integrins are clearly defective in their ability to induce tyrosine phosphorylation when plated on collagen IV.

Inhibition of Ligand-independent Focal Contact Localization

Recently, it has been shown that the α cytoplasmic domain may play a role in the control of integrin focal contact localization. Deletion of most of the $\alpha 1$ cytoplasmic domain (preserving the GFFKR) leads to ligand-independent focal contact localization on substrates that $\alpha 1\beta 1$ does not bind (Briesewitz *et al.*, 1993b). We decided to test whether this same property

Figure 5. The chimeric collagen receptors of SW1 and SWC cells do not support stress fiber or focal contact formation on collagen IV. HA1 cells (a, b, and c), SW1 cells (d, e, and f), and SWC cells (g, h, and i) were plated on collagen-coated coverslips and incubated at 37°C for 3 h. The cells were either stained with rhodamine-conjugated phalloidin (a, d, and g) or double-label immunofluorescence was performed using a mixture of a rabbit polyclonal anti-talin serum as a focal contact marker and the mouse monoclonal anti-human α_1 antibody TS2/7. The primary antibodies were visualized by the use of fluorescein-conjugated goat anti-rabbit IgG for the anti-talin serum (c, f, and i) and rhodamine-conjugated rabbit anti-mouse IgG for TS2/7 (b, e, and h). The diagrams to the right of the photomicrographs represent the exogenous subunits expressed in the cells.

occurred in the chimeric integrin subunits. We plated HA1, SW1, and SWC cells on FN and performed colocalization studies for α_1 integrin subunits, with talin as a marker of focal contact sites. As expected, when HA1 cells were plated on FN, the wild-type human α_1 subunit (Figure 7a) did not localize to focal contacts (Figure 7b). Similarly, when SW1 cells were plated on FN, the chimeric α_1 subunit (Figure 7c) did not localize to focal contact sites (Figure 7d). In contrast, when the SWC cells were plated on FN, the chimeric α_1 subunit lacking the distal α cytoplasmic domain (Figure 7e) exhibited ligand-independent focal contact localization (Figure 7f). Antibody inhibition studies (our unpublished results) reveal that the chimeric $\alpha_1\beta_1$

subunits do not contribute to FN adhesion (which is expected, because wild-type $\alpha_1\beta_1$ does not bind FN), and therefore confirm that this effect is ligand independent.

DISCUSSION

We have shown that the cytoplasmic domains of α and β subunits of integrin receptors are necessary for proper assembly and function. Previous work with soluble integrin subunits demonstrated the association of integrin extracellular domains (Dana *et al.*, 1991; Frachet *et al.*, 1992; Briesewitz *et al.*, 1993a), which is readily apparent in electron micrographs of

Figure 6. SW1 and SWC cells are defective in tyrosine phosphorylation when plated on collagen IV but not on fibronectin. Dishes (60 mm) were coated with FN or collagen IV at 37°C for 1 h. HA1 cells (lanes 1–3), SW1 cells (lanes 4–6), and SWC cells (lanes 7–9) were plated in the presence of BSA (1 mg/ml) on the FN (lanes 2, 5, and 8) or collagen IV (lanes 3, 6, and 9) dishes or left in suspension (lanes 1, 4, and 7). After incubation at 37°C for 1 h cells were washed twice with PBS and lysed in 300 μ l SDS-PAGE sample buffer. After lysis the samples were heated at 95°C for 10 min, the proteins were transferred to nitrocellulose, and Western blotting was performed using a polyclonal rabbit anti-phosphotyrosine serum and the ECL detection system.

integrin preparations (Carrell *et al.*, 1985; Nermut *et al.*, 1988). However, soluble subunits do not associate with endogenous integrins, suggesting that the transmembrane and/or cytoplasmic domains are required for normal integrin assembly. By creating chimeric and truncated forms of α and β subunits, we have shown that the KXGFFKR sequence and the β intracellular domain on opposing subunits are required for assembly of heterodimer. Deletion of the GFFKR sequence in either $\alpha 2$ or $\alpha 5$ leads to a reduced surface expression of $\alpha 2\beta 1$ or $\alpha 5\beta 1$ dimers (Bauer *et al.*, 1993; Kawaguchi and Hemler, 1993). Although this sequence does not appear to be required for the assembly of $\alpha IIb\beta 3$ in Chinese hamster ovary (CHO) cells (O'Toole *et al.*, 1991, 1994), it is our experience that the assembly of the wild-type versions of these two subunits is very inefficient in CHO cells, so that factors other than the KXGFFKR sequence may be rate limiting (our unpublished data).

We further showed that neither the α cytoplasmic domain C-terminal to the KXGFFKR nor the β transmembrane domain is necessary for stable heterodimer formation. In the $\beta 1$ cytoplasmic domain, the assembly motif may include the first 10 amino acids of the cytoplasmic domain (KLLMIIHDDR), because they are juxtaposed to the KXGFFKR sequence in the α cytoplasmic domain and deletion in this region led to a partial defect in dimerization (Hayashi *et al.*, 1990), while preservation of the dibasic (RR) allowed efficient assembly (Hayashi *et al.*, 1990; Marcantonio *et al.*, 1990). It is possible that the KXGFFKR of the α subunit and the KLLMIIHDDR of the $\beta 1$ subunit associate directly via their hydrophobic stretches as has been suggested (Williams *et al.*, 1994). Alternatively, the interaction of the two hydrophobic domains may be indirectly mediated by an assembly factor(s). If such

an association based on hydrophobic sequences occurs, it would likely be very weak and unstable in agents used to solubilize receptors (see also below).

Because integrin subunits may associate intracellularly and extracellularly, we correctly postulated that mutated chains with their transmembrane and cytoplasmic domains exchanged ($\alpha x\beta c$ and $\beta x\alpha c$) would assemble only with each other and mediate collagen adhesion without involving endogenous integrin subunits. These constructs, which were normally expressed and bound ligand normally, allowed us to study the role of cytoplasmic domain interactions in integrin function. When cells expressing both swapped chimeras (SW1) or both swapped chimeras with the α cytoplasmic domain deleted distal to the KXGFFKR (SWC) are plated on collagen IV, they bind ligand through their chimeric receptors, but exhibit poor focal contact formation and cytoskeletal reorganization. They also poorly stimulate tyrosine phosphorylation of FAK through their chimeric integrins. This result could be due to a defective direct or indirect interaction with FAK or a lack of cytoskeletal organization, which is required for integrin clustering and FAK phosphorylation (Shattil *et al.*, 1994).

The disruption of post-ligand binding events in SW1 and SWC cells is similar to that seen when fibroblasts adhere to the 105-kDa or 120-kDa central cell binding fragment of FN. In both instances, adhesion and spreading occur, but no stress fibers or focal contacts are formed (Woods *et al.*, 1986), and FAK phosphorylation is somewhat reduced (Guan *et al.*, 1991). In the case of SW1 and SWC cells, most cells spread less well on collagen IV than they would on FN (our unpublished results). However, we have shown examples of well spread cells to demonstrate that even in these cells there is little cytoskeletal organization. It is unclear whether the cytoskeleton is involved in the spreading of the SW1 and SWC cells on collagen IV or this occurs as a consequence of attachment to ligand. We favor the latter, both in these cell lines and in the adhesion to FN fragments. The adhesion phenotype on FN fragments appears to be due to a lack of protein kinase C activation, which is necessary for proper cytoskeletal organization (Woods and Couchman, 1992). It is tempting to speculate that the $\alpha x\beta c/\beta x\alpha c$ heterodimers cannot mediate or respond to the normal PKC signal produced by collagen IV, but this speculation requires a better understanding of the mechanism of the PKC effect.

We propose a working model to explain our results (Figure 8). The model shows that integrin heterodimers that have opposing α and β cytoplasmic domains have high surface expression, and those that lack either of these domains are not efficiently expressed on the surface. Our model proposes that during receptor assembly there may be an association of the KXGFFKR sequence with the corresponding β mo-

Figure 7. SWC cells display ligand-independent focal contact localization of their chimeric collagen receptor. HA1 cells, SW1 cells, and SWC cells were plated on fibronectin-coated coverslips and incubated at 37°C for 3 h. Double-label immunofluorescence was performed using a mixture of a rabbit polyclonal anti-talin serum as a focal contact marker and the mouse monoclonal anti-human $\alpha 1$ antibody TS2/7. The primary antibodies were visualized by the use of rhodamine-conjugated rabbit anti-mouse IgG for TS2/7 (a, c, and e) and fluorescein-conjugated goat anti-rabbit IgG for the anti-talin serum (b, d, and f). The diagrams to the right of the photomicrographs represent the exogenous subunits expressed in the cells.

tif directly or indirectly via an "assembly factor." This association would allow the α subunit to keep the β cytoplasmic domain function latent until ligand is bound, and hence would regulate ligand-dependent integrin cytoplasmic domain functions. In cells expressing the native $\alpha 1 \beta 1$ integrin (HA1), binding collagen IV would lead to the disruption of the α -dependent inhibition of the β cytoplasmic domain. The β cytoplasmic domain would then be free to mediate integrin post-ligand binding events, including cytoskeletal organization and signaling. It is unclear

why our swapped chimeric constructs are defective in post-ligand binding events, however, one possibility is that the normal change in the α and β cytoplasmic relationship does not occur, because the cytoplasmic domains are on the wrong extracellular domains.

Our results with SW1 cells indicate that merely the opposition of the α and β cytoplasmic domains, irrespective of their attached extracellular domains, is sufficient to prevent ligand-independent focal contact localization, which is an important feature of native integrins. Thus, these integrins do not localize to focal

Figure 8. Working hypothetical model of integrin subunit assembly. Integrin α and β subunits have compatible cytoplasmic domains, which, along with extracellular dimerization motifs, lead to efficient assembly into heterodimers. This assembly requires the cytoplasmic domains and does not require the α cytoplasmic domain beyond the KXGFFKR sequence. Substitution of the β cytoplasmic domain or a cytoplasmic domain with other cytoplasmic sequences ($\beta_x\alpha_c$; $\alpha_x\beta_c$) creates noncompatible cytoplasmic domains, which leads to inefficient assembly and low surface expression. Exchange of the α and β cytoplasmic domains ($\alpha_x\beta_c$ / $\beta_x\alpha_c$) onto the corresponding extracellular domains allows for efficient assembly and surface expression, due to the restoration of opposing compatible cytoplasmic domains. Further-

more, the switching of chains promotes selective assembly of the chimeric subunits. This drawing reflects the idea of directly associating cytoplasmic sequences. Alternatively, an assembly factor(s) may bind to both sequences mediating an indirect interaction.

adhesions formed by the endogenous $\alpha_5\beta_1$ on FN, because α_1 integrins do not bind FN (Figure 7). When the distal portion of the α subunit is deleted, however, the cells (SWC) lose this inhibition and the chimeric subunits localize to preformed focal contacts. This finding, which correlates with work on the native α_1 (Briesewitz *et al.*, 1993b) and α_{IIb} (Ylänne *et al.*, 1993) subunits, strongly suggests that focal contact localization is a direct effect of the uninhibited β_1 cytoplasmic domain, rather than an indirect one, such as a conformational change transmitted through the extracellular domains to the β_1 cytoplasmic domain. The latter mechanism could have occurred in the native α_1 subunit with such a deletion, but is extremely unlikely in the SWC cells due to the swapped intracellular and extracellular domains. We believe that these results support the notion of a functional association between α and β cytoplasmic domains.

Despite the importance of cytoplasmic domains in receptor function, we have been unable to demonstrate an association between α and β cytoplasmic domains independent of extracellular assembly. Using reporter constructs containing α and β cytoplasmic domains, we have found no evidence of association in co-precipitation or co-localization experiments (our unpublished data). We have also failed to demonstrate an association using the yeast two-cell hybrid system. We believe that extracellular association is necessary

to establish the stable association of these two small cytoplasmic domains. Nevertheless, the relationship between the α and β cytoplasmic domains is clearly crucial for several post-ligand binding functions of the receptor, and future work will be directed at understanding the nature of this interaction and the mechanisms of signal transduction involved.

ACKNOWLEDGMENTS

The authors gratefully acknowledge T. Giordano for the IL-2 receptor α subunit cDNA, T. Springer and I. Bank for the anti- α_1 antibodies, C. Buck for CSAT and Lenny antibodies, J. Sap for anti-phosphotyrosine antibodies, and K. Burridge for anti-talin antisera. The expert technical assistance of M. Epstein is much appreciated. Special thanks to Sorina Tugulea, Nicole Sociu-Foca, and Thomas Häupl for their help with the FACS analysis. We also thank M. Epstein, L. Smilenov, F. David, and P. Zage for their critical reading of the manuscript. This work was supported by National Institutes of Health grant GM-44585, and by grant #2906A from the Council for Tobacco Research. A.K. is supported by a long term fellowship (LT 653/93) from the Human Frontier Science organization. E.M. is an Established Scientist of the New York City affiliate of the American Heart Association.

REFERENCES

- Albelda, S.M., Daise, M., Levine, E.N., and Buck, C.A. (1989). Identification and characterization of cell substratum adhesion receptors on cultured human adult large vessel endothelial cells. *J. Clin. Invest.* 83, 992-1002.

- Bank, I., Hemler, M., Brenner, M.B., Cohen, D., Levy, V., Belko, J., Crouse, C., and Chess, L. (1989). A novel monoclonal antibody, 1B3.1, binds to a new epitope of the VLA-1 molecule. *Cell. Immun.* 122, 416–423.
- Bauer, J.S., Varner, J., Schreiner, C., Kornberg, L., Nicholas, R., and Juliano, R.L. (1993). Functional role of the cytoplasmic domain of the integrin $\alpha 5$ subunit. *J. Cell Biol.* 122, 209–221.
- Briesewitz, R., Epstein, M.R., and Marcantonio, E.E. (1993a). The expression of native and truncated forms of the human integrin $\alpha 1$ subunit. *J. Biol. Chem.* 268, 2989–2996.
- Briesewitz, R., Kern, A., and Marcantonio, E.E. (1993b). Ligand-dependent and -independent integrin focal contact localization: the role of the α chain cytoplasmic domain. *Mol. Biol. Cell* 4, 593–604.
- Burridge, K., Fath, K., Kelly, T., Nuckolls, G., and Turner, C. (1988). Focal adhesion: transmembrane junctions between the extracellular matrix and the cytoskeleton. *Annu. Rev. Cell Biol.* 4, 487–525.
- Burridge, K., Turner, C.E., and Romer, L.H. (1992). Tyrosine phosphorylation of paxillin and pp125^{FAK} accompanies cell adhesion to extracellular matrix: a role in cytoskeletal assembly. *J. Cell. Biol.* 119, 893–903.
- Carrell, N.A., Fitzgerald, L.A., Steiner, B., Erickson, H.P., and Phillips, D.R. (1985). Structure of human platelet glycoproteins IIb and IIIa as determined by electron microscopy. *J. Biol. Chem.* 260, 1743–1749.
- Chen, W.T., Hasegawa, C., Weinstock, C., and Yamada, K.M. (1985). Development of cell surface linkage complexes in cultivated fibroblasts. *J. Cell Biol.* 100, 1103–1114.
- Cobb, B.S., Schaller, M.D., Leu, T.-H., and Parsons, J.T. (1994). Stable association of pp60^{src} and pp59^{lyn} with the focal adhesion-associated protein tyrosine kinase, pp125^{FAK}. *Mol. Cell. Biol.* 14, 147–155.
- Damsky, C.M., Knudsen, K.A., Bradley, D., Buck, C.A., and Horwitz, A.F. (1985). Distribution of the CSAT cell-matrix antigen on myogenic and fibroblastic cells in culture. *J. Cell Biol.* 100, 1528–1539.
- Dana, N., Fathallah, D.M., and Arnaout, M.A. (1991). Expression of a soluble and functional form of the human beta 2 integrin CD11b/CD18. *Proc. Natl. Acad. Sci. USA* 88, 3106–3110.
- Dejana, E., Colella, S., Conforti, G., Abbadini, M., Gaboli, M., and Marchisio, P.C. (1988). Fibronectin and vitronectin regulate the organization of their respective Arg-Gly-Asp adhesion receptors in cultured human endothelial cells. *J. Cell Biol.* 107, 1215–1223.
- DeSimone, D.W., and Hynes, R.O. (1988). *Xenopus laevis* integrins: structural conservation and evolutionary divergence of integrin β subunits. *J. Biol. Chem.* 263, 5333–5340.
- Frachet, P., Duperray, A., Delachanal, E., and Marguerie, G. (1992). Role of the transmembrane and cytoplasmic domains in the assembly and surface exposure of the platelet integrin GPIIb/IIIa. *Biochemistry* 31, 2408–2415.
- Geiger, B., Salomon, D., Takeichi, M., and Hynes, R.O. (1992). A chimeric *N*-cadherin/ β_1 -integrin receptor which localizes to both cell-cell and cell-matrix adhesions. *J. Cell Sci.* 103, 943–951.
- Giordano, T., Howard, T.H., Coleman, J., Sakamoto, K., and Howard, B.H. (1991). Isolation of a population of transiently transfected quiescent and senescent cells by magnetic affinity cell sorting. *Exp. Cell Res.* 192, 193–197.
- Guan, J., Trevithick, J.E., and Hynes, R.O. (1991). Fibronectin/integrin interaction induces tyrosine phosphorylation of a 120 kDa protein. *Cell Regul.* 2, 951–964.
- Hartman, S.C., and Mulligan, R.C. (1988). Two dominant-acting selectable markers for gene transfer studies in mammalian cells. *Proc. Natl. Acad. Sci. USA* 85, 8047–8051.
- Hayashi, Y., Haimovich, B., Rezka, A., Boettiger, D., and Horwitz, A.F. (1990). Expression and function of chicken integrin $\beta 1$ subunit and its cytoplasmic domain mutants in mouse NIH 3T3 cells. *J. Cell Biol.* 110, 175–184.
- Horton, R.M., Cai, Z.L., Ho, S.N., and Pease, L.R. (1990). Generation of mutant cDNA by overlapping PCR. *Biotechniques* 5, 528–535.
- Horwitz, A.F., Duggan, E., Buck, C., Beckerle, M.C., and Burridge, K. (1986). Interaction of plasma membrane fibronectin receptor with talin: a transmembrane linkage. *Nature* 320, 531–533.
- Hynes, R.O. (1973). Alteration of cell-surface proteins by viral transformation and by proteolysis. *Proc. Natl. Acad. Sci. USA* 70, 3170–3174.
- Hynes, R.O., Marcantonio, E.E., Stepp, M.A., Urry, L.A., and Yee, G.E. (1989). Integrin heterodimer and receptor complexity in avian and mammalian cells. *J. Cell Biol.* 109, 409–420.
- Jaconi, M.E.E., Theler, J.M., Schlegel, W., Appel, R.D., Wright, S.D., and Lew, P.D. (1991). Multiple elevations of cytosolic-free Ca^{2+} in human neutrophils: initiation by adherence receptors of the integrin family. *J. Cell Biol.* 112, 1249–1257.
- Kawaguchi, S., and Hemler, M.E. (1993). Role of the α subunit cytoplasmic domain in regulation of adhesive activity mediated by the integrin VLA-2. *J. Biol. Chem.* 268, 16279–16285.
- Kern, A., Eble, J., Golbik, R., and Kühn, K. (1993). Interaction of type IV collagen with the isolated integrins $\alpha 1\beta 1$ and $\alpha 2\beta 1$. *Eur. J. Biochem.* 215, 151–159.
- Kornberg, L.J., Earp, H.S., Turner, C.T., Prockop, C., and Juliano, R.L. (1991). Signal transduction by integrins: increased protein tyrosine phosphorylation caused by clustering of $\beta 1$ integrins. *Proc. Natl. Acad. Sci. USA* 88, 8392–8396.
- Laemmli, U.K. (1970). Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature* 227, 680–685.
- LaFlamme, S.E., Akiyama, S.K., and Yamada, K.M. (1992). Regulation of fibronectin receptor distribution. *J. Cell Biol.* 117, 437–447.
- Marcantonio, E.E., and Hynes, R.O. (1988). Antibodies to the conserved cytoplasmic domain of the integrin β_1 subunit react with proteins in vertebrates, invertebrates and fungi. *J. Cell Biol.* 106, 1765–1772.
- Marcantonio, E.E., Guan, J.-L., Trevethick, J.E., and Hynes, R.O. (1990). Mapping of the functional determinants of the integrin β_1 cytoplasmic domain by site-directed mutagenesis. *Cell Regul.* 1, 597–604.
- Nermut, M.V., Green, N.M., Eason, P., Yamada, S.S., and Yamada, K.M. (1988). Electron microscopy and structural model of human fibronectin receptor. *EMBO J.* 7, 4093–4099.
- O'Toole, T.E., Katagiri, Y., Faull, R.J., Peter, K., Tamura, R., Quaranta, V., Loftus, J.C., Shattil, S.J., and Ginsberg, M.H. (1994). Integrin cytoplasmic domains mediate inside-out signal transduction. *J. Cell Biol.* 124, 1047–1059.
- O'Toole, T.E., Mandelman, D., Forsyth, J., Shattil, S.J., Plow, E.F., and Ginsberg, M.H. (1991). Modulation of the affinity of integrin $\alpha \text{IIb}\beta 3$ (GP IIb-IIIa) by the cytoplasmic domain of αIIb . *Science* 254, 845–847.
- Otey, C.A., Pavalko, F.M., and Burridge, K. (1990). An interaction between α -actinin and the $\beta 1$ integrin subunit in vitro. *J. Cell Biol.* 111, 721–729.
- Reszka, A.A., Hayashi, Y., and Horwitz, A.F. (1992). Identification of amino acid sequences in the integrin $\beta 1$ cytoplasmic domain implicated in cytoskeletal association. *J. Cell Biol.* 117, 1321–1329.

- Schaller, M.D., Borgman, C.A., Cobb, B.S., Vines, R.R., Reynolds, A.B., and Parsons, J.T. (1992). pp125^{FAK}, a structurally unique protein tyrosine kinase associated with focal adhesions. *Proc. Natl. Acad. Sci. USA* 89, 5192–5196.
- Schaller, M.D., Hildebrand, J.D., Shannon, J.D., Fox, J.W., Vines, R.R., and Parsons, J.T. (1994). Autophosphorylation of the focal adhesion kinase, pp125^{FAK}, directs SH2-dependent binding of pp60^{src}. *Mol. Cell. Biol.* 14, 1680–1688.
- Schlaepfer, D.D., Hanks, S.K., Hunter, T., and van der Geer, P. (1994). Integrin-mediated signal transduction linked to Ras pathway by GRB2 binding to focal adhesion kinase. *Nature* 372, 786–791.
- Schwartz, M.A. (1993). Spreading of human endothelial cells on fibronectin or vitronectin triggers elevation of intracellular free calcium. *J. Cell Biol.* 120, 1003–1010.
- Schwartz, M.A., Ingber, D.E., Lawrence, M., Springer, T.A., and Lechene, C. (1991a). Multiple integrins share the ability to induce elevation of intracellular pH. *Exp. Cell Res.* 195, 533–535.
- Schwartz, M.A., Lechene, C., and Ingber, D.E. (1991b). Insoluble fibronectin activates the Na/H antiporter by clustering and immobilizing integrin $\alpha 5 \beta 1$, independent of cell shape. *Proc. Natl. Acad. Sci. USA* 88, 7849–7853.
- Shattil, S.J., Haimovitch, B., Cunningham, M., Lipfert, L., Parsons, J.T., Ginsberg, M.H., and Brugge, J.S. (1994). Tyrosine phosphorylation of pp125^{FAK} in platelets requires coordinated signaling through integrin and agonist receptors. *J. Biol. Chem.* 269, 14738–14745.
- Singer, I.I., Scott, S., Kawka, D.W., Kazazis, D.M., Gailit, J., and Rouslahti, E. (1988). Cell surface distribution of fibronectin and vitronectin receptor depends on substrate composition and extracellular matrix accumulation. *J. Cell Biol.* 106, 2172–2182.
- Solowska, J., Guan, J.-L., Marcantonio, E.E., Trevithick, J.E., Buck, C.A., and Hynes, R.O. (1989). Expression of normal and mutant avian integrin subunits in rodent cells. *J. Cell Biol.* 109, 853–861.
- Southern, P.J., and Berg, P. (1982). Transformation of mammalian cells to antibiotic resistance with a bacterial gene under control of the SV40 early region promoter. *J. Mol. Appl. Genet.* 1, 327–341.
- Tamkun, J.W., DeSimone, D.W., Fonda, D., Patel, R.S., Buck, C., Horwitz, A.F., and Hynes, R.O. (1986). Structure of integrin, a glycoprotein involved in the transmembrane linkage between fibronectin and actin. *Cell* 42, 271–282.
- Towbin, H., Staehelin, T., and Gordon, J. (1979). Electrophoretic transfer from polyacrylamide gels to nitrocellulose sheets. *Proc. Natl. Acad. Sci. USA* 76, 4350–4354.
- Vandenberg, P., Kern, A., Ries, A., Luckenbill-Edds, J., Mann, K., and Kühn, K. (1991). Characterization of a type IV collagen major cell binding site with affinity to the alpha 1 beta 1 and the alpha 2 beta 1 integrins. *J. Cell Biol.* 113, 1475–1483.
- Vuori, K., and Ruoslahti, E. (1994). Association of insulin receptor substrate-1 with integrins. *Science* 266, 1576–1578.
- Williams, M.J., Hughes, P.E., O'Toole, T.E., and Ginsberg, M.H. (1994). The inner world of cell adhesion: integrin cytoplasmic domains. *Trends Cell Biol.* 4, 109–112.
- Woods, A., and Couchman, J.R. (1992). Protein kinase C involvement in focal adhesion formation. *J. Cell Sci.* 101, 277–290.
- Woods, A., Couchman, J.R., Johansson, S., and Höök, M. (1986). Adhesion and cytoskeletal organization of fibroblasts in response to fibronectin fragments. *EMBO J.* 5, 665–670.
- Xing, Z., Chen, H.-C., Nowlen, J.K., Taylor, S.J., Shalloway, D., and Guan, J.-L. (1994). Direct interaction of vSrc with the focal adhesion kinase mediated by the Src SH2 domain. *Mol. Biol. Cell* 5, 413–421.
- Ylänne, J., Chen, Y.L., O'Toole, T.E., Loftus, J.C., Takada, Y., and Ginsberg, M.H. (1993). Distinct functions of integrin α and β subunit cytoplasmic domains in cell spreading and formation of focal adhesions. *J. Cell. Biol.* 122, 223–234.