Organelle-Cytoskeletal Interactions: Actin Mutations Inhibit Meiosis-dependent Mitochondrial Rearrangement in the Budding Yeast *Saccharomyces cerevisiae*

Michael G. Smith, Viviana R. Simon, Honor O'Sullivan, and Liza A. Pon*

Department of Anatomy and Cell Biology, Columbia University College of Physicians and Surgeons, New York, New York 10032

Submitted May 1, 1995; Accepted August 8, 1995
Monitoring Editor: David Botstein

During early stages of meiosis I, yeast mitochondria fuse to form a single continuous thread. Thereafter, portions of the mitochondrial thread are equally distributed to daughter cells. Using time-lapse fluorescence microscopy and a membrane potential sensing dye, mitochondria are resolved as small particles at the cell periphery in pre-meiotic, living yeast. These organelles display low levels of movement. During meiosis I, we observed a threefold increase in mitochondrial motility. Mitochondrial movements were linear, occurred at a maximum velocity of 25 ± 6.7 nm/s, and resulted in organelle collision and fusion to form elongated tubular structures. Mitochondria do not co-localize with microtubules. Destabilization of microtubules by nocodazole treatment has no significant effect on the rate and extent of thread formation. In contrast, yeast bearing temperature-sensitive mutations in the actin-encoding *ACT1* gene (*act1-3* and *act1-133*) exhibit abnormal mitochondrial aggregation, fragmentation, and enlargement as well as loss of mitochondrial motility. In *act1-3* cells, mitochondrial defects and actin delocalization occur only at restrictive temperatures. The *act1-133* mutation, which perturbs the myosin-binding site of actin without significantly affecting actin cytoskeletal structure in meiotic yeast, results in mitochondrial morphology and motility defects at restrictive and permissive temperatures. These studies support a role for the actin cytoskeleton in the control of mitochondrial position and movements in meiotic yeast.

INTRODUCTION

Mitochondrial inheritance is the process whereby mitochondria are transferred from mother to developing daughter during cell division. Because mitochondria are essential organelles and are only produced from pre-existing mitochondria, this process is critical for viable progeny. In the budding yeast *Saccharomyces cerevisiae*, spore formation is a meiosis-dependent differentiation process (Esposito and Klapholz, 1981). Mitochondrial inheritance during meiosis proceeds by a defined pattern of rearrangement events (Stevens,

1981; Miyakawa *et al.*, 1984). In pre-meiotic, stationary phase yeast, mitochondria are punctate and localized to the cell periphery. During meiotic prophase, these organelles align and fuse, ultimately forming a single highly branched mitochondrial reticulum. Serial thin section electron microscopy studies indicate that the mitochondrial reticulum is produced by fusion of up to 50 individual organelles (Stevens, 1977). Before meiosis II, the mitochondrial reticulum moves toward the nucleus and associates with the nuclear membrane (Stevens, 1981). This complex mitochondrion remains associated with the nucleus through meiosis II. As a result, a branch of the mitochondrial reticulum is included in each developing spore during prospore wall formation. Thus, meiosis-linked mitochondrial alignment and fusion contribute to mitochondrial inheri-

* Corresponding author: Department of Anatomy and Cell Biology, Columbia University College of Physicians and Surgeons, 630 West 168th Street, New York, NY 10032.

tance by consolidating the organelle before its equal distribution to newly developed daughter spores.

The mechanism of this mitochondrial rearrangement is not well understood. In other systems, organelles not only co-localize with microtubules, microfilaments, and intermediate filaments (Hirokawa, 1982; Bridgmen *et al.*, 1986; Menzel and Schliwa, 1986) but move along microtubules (Allen *et al.*, 1982; Brady *et al.*, 1982; Hayden and Allen, 1984; Koonce and Schliwa, 1985) and microfilaments (Kachar, 1985; Adams and Pollard, 1986; Kachar and Reese, 1988; Kohno and Shimmen, 1988; Hegmann *et al.*, 1989, 1991; Wesels *et al.*, 1989; Kuznetsov *et al.*, 1992). Motor molecules, which are cytoskeleton-associated ATPases, drive this organellar movement (Pollard *et al.*, 1991; reviewed by Schroer and Sheetz, 1991; Titus, 1993). Thus, the cytoskeleton appears to act as a scaffold to immobilize organelles and as a track for energy-dependent, directed movement.

In yeast, cytoplasmic and nuclear microtubules emerge from the spindle pole body, a single microtubule organizing center embedded in the nuclear envelope (Moens and Rapport, 1971; Byers, 1981; Esposito and Klapholz, 1981). Microtubules play a role in chromosome segregation and nuclear migration during cell division and mating (Byers, 1981; Rose and Fink, 1987; Jacobs *et al.*, 1988). Microtubule-vacuole interactions may also occur in yeast, because destabilization of microtubules results in vacuole fragmentation (Guthrie and Wickner, 1988).

The yeast actin cytoskeleton consists of actin cables, bundles of actin filaments, and actin patches, plasma membrane invaginations associated with actin, and the actin-binding proteins fimbrin, Abp1p, cofilin, and capping protein (Drubin *et al.*, 1988; Adams *et al.*, 1991; Amatruda and Cooper, 1992; Moon *et al.*, 1993; Mulholland *et al.*, 1994). Phenotypic analysis of yeast mutants supports a role for the actin cytoskeleton in polarized cell growth (Novick and Botstein, 1985; Drubin *et al.*, 1993), mating projection formation (Read *et al.*, 1992), bud separation during cytokinesis (Novick and Botstein, 1985), control of spindle position during anaphase (Palmer *et al.*, 1992), response to osmotic stress (Chowdhury *et al.*, 1992), as well as vesicle movement during bud expansion, early stages of endocytosis, and late stages of secretion (Johnston *et al.*, 1991; Liu and Bretscher, 1992; Kübler and Riezman, 1993; Govindan *et al.*, 1995). Other studies support a role for the actin cytoskeleton in the control of mitochondrial movement in vegetative, mitotically dividing yeast. Optical microscopic studies indicate that mitochondria co-localize with F-actin bundles, and that actin mutations result in abnormal mitochondrial distribution, movement, and inheritance (Drubin *et al.*, 1993; Lazzarino *et al.*, 1994; Simon *et al.*, 1995). In addition, we find that isolated yeast mitochondria display ATP-sensitive binding to yeast actin filaments.

This binding is reversible, saturable, and occurs through protein-mediated binding of mitochondria to the lateral surface of actin filaments (Lazzarino *et al.*, 1994). Finally, using a microfilament sliding assay, we identified an actin-based motor activity on mitochondrial outer membranes (Simon *et al.*, 1995).

Here, we studied actin and microtubule structure in meiotic yeast, and the role of these cytoskeletal systems in mitochondrial movement and inheritance. The spatial relations between mitochondria and the major yeast cytoskeletal networks were examined using nocodazole to destabilize microtubules and temperature-sensitive lethal mutations (*act1-3* and *act1-133*) to perturb the actin-encoding *ACT1* gene. Our studies support a role for the actin cytoskeleton in control of mitochondrial movements and inheritance during yeast cell meiosis.

MATERIALS AND METHODS

Sporulation and Fixation

Yeast strains used in this study are listed in Table 1. Sporulation of spheroplasts in liquid culture was carried out as described by Miyakawa *et al.* (1984). Yeast cells were grown to late log phase in rich liquid medium containing glucose (YPD). Cells were then concentrated by centrifugation ($1000 \times g$ for 5 min), washed once with water, and incubated with shaking in wash solution (25 mM potassium phosphate buffer, pH 7.5, 0.8 M KCl) supplemented with 10 mM dithiothreitol for 15 min at room temperature (RT). Cells were collected by centrifugation, washed twice, and incubated in wash solution containing 5 mg zymolyase 20,000T (ICN, Costa Mesa, CA) per gram of cells for 15 min at RT. Spheroplasts produced by zymolyase-catalyzed degradation of the yeast cell wall were then washed in ice cold wash solution. Spheroplast recovery was determined by visual analysis using a hemocytometer.

Spheroplasts were suspended in sporulation media (15 mM potassium acetate, 50 mM KPO₄, pH 6.9, 1.2 M sorbitol) at a concentration of 10^7 cells/ml. Samples were incubated at 22, 30, or 37°C with gentle agitation for 0–22 h. To fix samples, paraformaldehyde and potassium phosphate buffer (pH 6.5) were added directly to the sporulation medium to final concentrations of 3.7% and 100 mM, respectively. This mixture was incubated for 15 min at RT and cells were collected by centrifugation. The resulting pellet was resuspended in a second fixative solution (100 mM KPO₄, pH 6.5, 2 mM MgCl₂, and 5% paraformaldehyde) for 1 h at RT. Fixed spheroplasts were washed three times with NS (20 mM Tris-HCl, pH 7.6, 0.25 M sucrose, 1 mM EDTA, 1 mM MgCl₂, 0.1 mM ZnCl₂, 0.1 mM CaCl₂, 0.8 mM phenylmethylsulfonylfluoride, 0.05% [v/v] 2-mercaptoethanol). For some studies, 12.5 µg/ml nocodazole (Sigma Chemical, St. Louis, MO) was added directly to the sporulation media. Because nocodazole was solubilized in dimethyl sulfoxide, dimethyl sulfoxide (0.13% v/v final) was added to control samples.

Visualization of Mitochondrial and Cytoskeletal Structures in Fixed Samples

The method used for indirect immunofluorescence is a modification of published methods (Pringle *et al.*, 1989). Fixed spheroplasts were applied to poly-lysine-coated coverslips and allowed to adhere to the coverslips for 20 min. Immobilized spheroplasts were then gently washed in phosphate-buffered saline (PBS) and incubated in PBT (1× PBS, 0.1% [v/v] Triton X-100, 0.02% [v/v] sodium azide, 1% [w/v] bovine serum albumin) for 5 min at RT. This was followed

Table 1. Yeast strains used in this study

Strain	Genotype	Source
g833-1B	<i>MATa, leu2, can1, HOM3, his1-1, trp2, ADE2, ho, gal</i>	Yeast Genetic Stock Center
g833-2D	<i>MATα, LEU2, CAN1, hom3-10, his1-7, TRP2, ade2, ho, gal</i>	Yeast Genetic Stock Center
MSY1	<i>g833-1B × g833-2D</i>	This study
KWY474	<i>MATa, /MATα, ACT1:: his3/ACT1::his3, ade2/ADE2</i>	K. Wertman
KWY230	<i>leu2-3,112/leu2-3,112, ade4/ADE4, can1-1/can1-1</i>	K. Wertman
DDY437	<i>MATa, /MATα, act1-3/act1-3, his3Δ200/his3Δ200, ura3-52/ura3-52, LEU2/leu2-3,112, LYS2/lys2-801</i>	D. Drubin
DBY877	<i>MATa, /MATα, ura3-52/ura3-52, his3Δ200/his3Δ200</i>	D. Drubin
DBY1691	<i>ADE2/ade2, tub2-201/tub2-201, leu2-3,112/leu2-3,112, ade4/ADE4, act1-133::HIS3/act1-133::HIS3</i>	D. Drubin
MSY101	<i>MATα, his4-619</i>	D. Drubin
MSY106	<i>MATa, his4-619, act1-3</i>	This study
MSY202	<i>DBY877 × DBY1691</i>	This study
	<i>MATa/MATα, his4-619/his4-619</i>	This study
	<i>MATa/MATα, his4-619/HIS4, trp1-1/TRP1, leu2-3,112/LEU2, ura3-1/URA3, act1-3/act1-3</i>	This study

by incubation with primary antibody for 2 h at RT in a humid chamber. YOL 1/34, a rat monoclonal antibody raised against yeast tubulin (Kilmartin and Adams, 1984) was used to detect microtubules. A polyclonal antiserum raised against total mitochondrial outer membrane proteins was used to label mitochondria. The anti-outer membrane antibody was prepared by immunizing rabbits with mitochondrial outer membranes, as described previously (Riezman *et al.*, 1983). Analysis of the antiserum using Western blots indicates that the antibody recognizes all of the major outer membrane proteins (porin, Mas70p, and the 33- and 45-kDa outer membrane proteins). The anti-outer membrane antiserum was pre-adsorbed with intact yeast cells before use.

Subsequent to incubation with primary antibodies, spheroplasts were washed with PBT and incubated with fluorescently labeled secondary antibody for 60 min at RT in the dark. The secondary antibodies used for these studies, fluorescein isothiocyanate-coupled goat anti-rabbit IgG (Boehringer Mannheim, Indianapolis, IN) and rhodamine-coupled goat anti-rat IgG (Kirkegaard and Perry Laboratories, Gaithersburg, MD), were reconstituted, stored, and used according to the manufacturer's instructions. The spheroplasts were washed with PBS to remove unbound secondary antibody and mounted on microscope slides using mounting solution (1 mg/ml of *p*-phenylenediamine, 90% [w/v] glycerol, and 1× PBS) that contained the DNA-binding dye 4,6-diamidino-2-phenylindole (DAPI; Williamson and Fennell, 1975) at a final concentration of 1 μg/ml. Coverslips were then sealed onto glass microscope slides with clear nail polish. Slides were stored at -20°C in the dark before viewing with a Leitz Dialux fluorescence microscope (Rockleigh, NJ).

The actin cytoskeleton was visualized using rhodamine-coupled phalloidin (Molecular Probes, Eugene, OR), a ligand that binds specifically to actin polymers (Cooper, 1987), using published protocols (Pringle *et al.*, 1989). The fluorescent ligand was added to fixed spheroplast samples to a final concentration of 2.5 mM in a solution consisting of a 4:1 ratio of NS to methanol, and samples were allowed to stand in the dark at 4°C for 16 h. Thereafter, the mixture was diluted with 10 volumes of PBS and applied directly to polylysine-coated coverslips. To visualize actin and mitochondria in the same samples, immobilized rhodamine phalloidin-stained spheroplasts were washed with PBS and applied to microscope slides using DAPI-containing mounting solution.

Confocal Microscopy

Confocal images of cells were obtained with a laser scanning confocal microscope (MRC 600; Bio-Rad Microscience, Cambridge, MA)

on a Zeiss Axiovert inverted microscope (Oberkochen, Germany) using a 63× (N.A. 1.4) Zeiss Plan-Apo infinity corrected objective. The illumination sources were the 488- and 514-nm lines from a 25-mW argon laser. Fluorescein-labeled structures were visualized with a 488-nm bandpass excitation filter, a 510-nm dichroic mirror, and a 515-nm long pass emission filter. Rhodamine-phalloidin was visualized with a 514-nm bandpass excitation filter, a 540-nm dichroic mirror, and a 550-nm long pass emission filter. Light was attenuated using neutral density filters in the excitation path. Each field of cells was sectioned three dimensionally by recording images from a series of 0.36 μm optical sections. Movement from one focal plane to the next was achieved using a stepper motor attached to the fine focus control of the microscope. Each optical section collected was the average of 15 scans at the confocal microscope's normal scan rate (1 frame/second). All images were processed and enhanced using the COMOS program (Bio-Rad Microscience) and Adobe Photoshop 2.5 (Mountain View, CA).

Vital Dye Staining and Time Lapse Microscopy

Diploid yeast cells were grown to late log phase in YPD at 22 or 30°C and converted to spheroplasts as described above. Spheroplasts were washed twice and resuspended in sporulation media at a concentration of 10⁷ cells/ml. After incubation in sporulation media, aliquots of spheroplasts were removed and incubated with 3,3'-dihexyloxycarbocyanide iodide (DiOC₆; 50 ng/ml; Polysciences, Warrington, PA) for 5 min at RT. Stained cells were rinsed once, resuspended in sporulation media to a final concentration of 2 × 10⁸ cells/ml, and applied to a microscope slide.

Prepared slides were visualized with a 100×, 1.32 NA oil immersion objective on a Leitz Dialux fluorescence microscope equipped with a 3.1× projection lens (Diagnostics Instruments, Sterling Heights, MI) and a Star-1 cooled CCD camera (Photometrics, Tucson, AZ). Time-lapse images were acquired at 15- to 30-s intervals for a period of 10 min, processed by the National Institutes of Health Image program (v1.55) on a Macintosh Quadra 800 (Apple Computer, Cupertino, CA), and stored using an optical magnetic disk drive (Peripheral Land, Fremont, CA). To minimize sample illumination, light from the 100W mercury arc lamp source was controlled using an automated shutter driver (Uniblitz D122, Vincent and Associates, Rochester, NY), and attenuated using neutral density filters (Omega Optical Corporation, Battleboro, VT).

Electron Microscopy

Preparation of samples for transmission electron microscopy was carried out according to the methods of Stevens (1977). Yeast spheroplasts were fixed by addition of glutaraldehyde (Sigma Chemical) to sporulation medium to a final concentration of 5%. After incubation for 3 h at RT, spheroplasts were concentrated by centrifugation ($10,000 \times g$, 10 min, RT), and washed two times with 0.9% NaCl. Samples were resuspended in 4% KMnO_4 in 0.1 M sodium cacodylate, pH 7.4 (Electron Microscopy Sciences, Fort Washington, PA), and incubated at 4°C for 1 h with gentle rotation. After 2 washes with 0.9% NaCl, samples were resuspended in 2% uranyl acetate (Electron Microscopy Sciences) and incubated for 1 h at RT. After three washes, the samples were dehydrated in a graded series of ethanol solutions, infiltrated with propylene oxide for 10 min, and embedded in Epon-812 (Tousimis Research, Rockville, MD). Ultrathin sections were stained for 5 min with 1% lead citrate before viewing with a JEOL 1200 transmission electron microscope.

RESULTS

Mitochondrial Rearrangements during Early Stages of Meiosis

Mitochondria in pre-meiotic, late log phase spheroplasts are punctate structures approximately $0.5 \mu\text{m}$ in diameter that are localized at the cell periphery. After 2.5 h of incubation under sporulation conditions, mitochondria begin to align and fuse, forming elongated tubular structures (Figure 1). Tubular mitochondria are approximately $0.3 \mu\text{m}$ in diameter and increase in length as a function of incubation time in sporulation medium.

Mitochondrial rearrangement is a highly synchronous, widespread event. In wild-type MSY1 cells, 100% of the stationary phase cells examined contained punctate, peripherally localized mitochondria. Five hours after transfer of cells to sporulation medium, 46% of cells underwent mitochondrial rearrangement resulting in a decrease in the number of cells containing circumferential punctate mitochondria and a corresponding increase in the number of cells containing tubular mitochondria. Mitochondrial thread formation is an intermediate and transient stage of spore formation. After 22 h of incubation in sporulation medium, the percentage of cells displaying mitochondrial threads decreased and the percentage of tetrads (collections of haploid spores) increased (Figure 2). The kinetics and extent of spore formation in spheroplasts is similar to that observed in yeast cells containing complete cell walls. Spheroplasts were used for these studies because internal structures are better resolved upon removal of the cell wall.

Time-lapse Fluorescence Microscopy and Vital Staining Using DiOC₆

A combination of time-lapse fluorescence microscopy and vital staining using the lipophilic membrane potential sensing dye DiOC₆ was used to investigate mitochondrial movements associated with mitochondrial thread formation in living cells. In yeast, the dye

Figure 1. Mitochondrial rearrangement and reticulum formation during early stages of meiosis. To obtain synchronous cultures, wild-type cells (KWY474) were grown to late log phase, converted to spheroplasts, and incubated in sporulation medium at 30°C. Cells were then fixed and mitochondria were visualized by indirect immunofluorescence as described in MATERIALS AND METHODS. (A and B) Mitochondria in yeast spheroplasts 0 and 2.5 h after transfer to sporulation medium. Bar, $2.0 \mu\text{m}$.

stains mitochondria at low concentrations (10–100 ng/ml) (Koning *et al.*, 1993). Specific mitochondrial staining in meiotic yeast was confirmed by double label experiments using DAPI and DiOC₆.

Mitochondrial movements were readily detectable in DiOC₆-stained spheroplasts after 2 h of incubation in sporulation medium (Figure 3A). At this stage of meiosis, mitochondria were resolved as spherical particles or short tubules localized at the periphery. To evaluate the dynamics of mitochondrial motility, we traced the movements of individual organelles over

Figure 2. Time course and synchrony of mitochondrial rearrangement. Meiosis-linked mitochondrial rearrangement was examined in wild-type yeast (MSY1) as for Figure 1. The percentage of cells that contain punctate mitochondria located at the cell perimeter (squares), elongated mitochondrial tubules (circles), or completed spores (triangles) was determined as a function of incubation time in sporulation medium. Cells that contained punctate and tubular mitochondria were scored as punctate or tubular on the basis of the most abundant mitochondrial structure.

3–5 min of real time (Figure 3B). Spherical and short tubular mitochondria exhibited short and long distance linear motions that varied in length from 0.1–1.5 μm . In addition, although mitochondrial movement was continuous, individual organelles were observed to stop and resume motion, and to undergo directional changes.

Mitochondrial fusion is difficult to evaluate by time-lapse imaging. However, previous studies documenting mitochondrial DNA mixing and formation of a continuous mitochondrial reticulum indicate that tubular mitochondria are produced during meiosis by organelle fusion (Stevens, 1977; Uchida and Suda, 1978). Consistent with this, we found that some mitochondrial movements resulted in collision and subsequent production of tubular mitochondria. Because these tubular organelles were able to move as a single unit, they were interpreted as products of mitochondria fusion. Not all collisions resulted in mitochondrial fusion; in many cases organelles were observed to collide, and move away from each other without any obvious change in mass or shape.

The extent of mitochondrial movement in living spheroplasts increases as a function of sporulation time (Table 2). 15.4% of the mitochondria were motile

in pre-meiotic cells. When movements were detected, they were generally short distance and did not result in any net change in the shape or position of the organelle. In contrast, we observed a threefold increase in the percentage of motile mitochondria in meiotic yeast. Although the number of motile elements increased as meiosis proceeded, the maximum velocity of mitochondrial movement ($25 \pm 6.7 \text{ nm/s}$) remained constant. Collectively, the findings that mitochondrial movements are linear, occur at a constant velocity, and are up-regulated during meiosis, suggest that mitochondrial movements are nonrandom, directed events.

Microtubules Are Not Required for Mitochondrial Reorganization

We examined the possible role of microtubules in meiosis-linked mitochondrial motility by examining the following: 1) the spatial relationship between microtubules and mitochondria within intact spheroplasts, and 2) the effect of a microtubule-destabilizing agent on mitochondrial movement and thread formation. After incubation in sporulation medium for 5 h, bar-shaped spindles that colocalize with the nucleus were detected. In the same cells, mitochondria were resolved as short tubular structures. Thus, mitochondrial rearrangement occurs concomitant with spindle formation during meiosis I (Figure 4).

Mitochondria do not co-localize with microtubules. Further, destabilization of microtubules using nocodazole had no significant effect on mitochondrial rearrangements during sporulation (Figures 4 and 5). After 5 h of incubation in nocodazole-containing sporulation medium, the only microtubule-containing structure detected in 95.1% of the spheroplasts examined was the spindle pole body. Thus, treatment with nocodazole resulted in microtubule depolymerization. Mitochondria, however, were not affected by this treatment. Formation of mitochondrial threads occurred to the same extent in the absence as in the presence of microtubules. Consistent with this, time lapse video imaging of nocodazole-treated samples indicated that this treatment had no significant effect on either the velocity or extent of mitochondrial movement (Table 4).

Actin Structures in Meiotic Yeast Cells

We examined the effect of two temperature-sensitive lethal mutations in the actin-encoding *ACT1* gene on actin and mitochondrial morphologies in meiotic yeast. The *act1-3* mutant contains an amino acid substitution of leucine for proline at position 32, and is therefore identical to the *act1-1* allele (Shortle *et al.*, 1984). The *act1-133* mutant (Wertman *et al.*, 1992) contains two amino acid substitutions of

alanine for aspartate at positions 24 and 25, residues within the myosin binding site of yeast actin (Kabsch *et al.*, 1990; Rayment *et al.*, 1993).

Yeast cells entering meiosis contain the same actin structures reported in mitotic cells (Adams and Pringle, 1984): confocal microscopy revealed rhodamine-

Figure 3.

phalloidin-labeled actin cables and cortical actin patches in pre-meiotic yeast and in cells after 5 h of incubation in sporulation medium. These actin structures were present in wild-type yeast upon sporulation at 22 or 37°C (Figure 6). Although actin cables were absent, cortical actin patches were detected in the *act1-3* mutant cells upon sporulation at the permissive temperature. Incubation of *act1-3* cells at elevated temperatures resulted in delocalization of the actin cytoskeleton, i.e., loss of detectable actin patches and accumulation of diffuse, cytoplasmic staining (Figure 6). Quantitation of these results indicates that actin delocalization occurs to some extent in *act1-3* cells at permissive temperatures, and increases by a factor of 3 upon short term shift to restrictive conditions (Table 3). In contrast, actin cables and cortical patches were detected in the *act1-133* mutant cells upon sporulation at permissive or nonpermissive temperatures (Figure 6). Although the patches cluster in some cells, the distribution of patches and cables were not altered significantly by the *act1-133* allele.

Defects in Mitochondrial Arrangement and Morphology in Actin Mutants

To examine the role of the actin cytoskeleton in mitochondrial motility, we studied the effect of temperature shift on meiosis-linked mitochondrial rearrangements in *act1-3* and *act1-133* strains (Figure 7 and Table 3). The *act1-3* mutant displayed temperature-dependent defects in mitochondrial morphology and localization during the early stages of meiosis. Upon sporulation at permissive temperatures, tubular mitochondria were detected in the *act1-3* mutant at levels similar to those of wild-type cells. Low levels of mitochondrial thread formation were also observed

Table 2. Up-regulated mitochondrial motility during early stages of meiosis

Time (h)	Motile mitochondria (%)	Velocity (nm/s)
0-1	15.4	25.5 ± 9.8
3-4	27.0	20.4 ± 4.6
5-6	45.0	28.2 ± 4.1

Late log phase yeast were converted to spheroplasts and incubated in sporulation medium at 22°C. At the times indicated, aliquots of the cultures were stained with DiOC₆ and movements of stained mitochondria were visualized at 30°C using time lapse video microscopy. The velocity of mitochondrial movement was determined by measuring the change in position of a particle as a function of time. The limit of resolution of our system is 1 pixel/20 s or approximately 5.5 nm/s. Mitochondrial movements that resulted in displacement less than 1 pixel, or which persisted for less than three consecutive still frames, were assigned a velocity of 0 nm/s. The linear distances between successive positions of mitochondria were calculated using Microsoft Excel, and instantaneous velocities were averaged to give the mean velocity. Velocities are mean values obtained from three experiments and analysis of over 60 moving particles.

upon sporulation of *act1-3* cells at restrictive temperatures for 1-2 h. However, after 5 h of sporulation at 37°C we observed the following: 1) low levels of mitochondrial aggregation, and 2) a decrease in the number of cells containing tubular mitochondrial structures coupled with a corresponding increase in mutants with abnormally large spherical mitochondria. These findings suggest that prolonged incubation of *act1-3* mutants at restrictive conditions results in defects in mitochondrial morphology and

Figure 3 (cont). (A) Time-lapse images of meiosis-linked mitochondrial movements. Late log phase cells were converted to spheroplasts and incubated in sporulation medium as for Figure 1. Aliquots were removed at 2-5 h after transfer to sporulation medium. Mitochondria were stained with DiOC₆ and images of stained mitochondria were collected at 20-s intervals, as described in MATERIALS AND METHODS. Motile mitochondria were defined as particles that display linear motion for at least 60 s of real time. Panels A-C and D-F are consecutive images of single DiOC₆-stained spheroplasts at 40-s intervals. White arrowheads point to mitochondrial particles displaying linear motion; in panels D-F, these motions lead to production of tubular organelles. Black arrowheads indicate the positions of mitochondria undergoing collision; in panels D-F, this organelle collision does not result in fusion. Black arrows point to the leading tip of motile tubular mitochondria. Bar, 1.0 μm. (B) Dynamics of mitochondrial movement. Tracings of the movements of individual organelles in DiOC₆-stained,

sporulating spheroplasts were made by following single particles during the time in which they were in the plane of focus. Points denote the position of organelles at arbitrary time points and do not connote velocity. Bar, 1.0 μm.

Figure 4. Effect of nocodazole on microtubule integrity and mitochondrial rearrangements. Late log phase cells (MSY106) were converted to spheroplasts, and incubated in sporulation medium in the absence (A, C, and E) or presence (B, D, and F) of nocodazole (12.5 $\mu\text{g/ml}$) for 5 h at 30°C. Fixed samples were stained with the DNA binding dye DAPI (A and B) and by indirect immunofluorescence using antibodies specific for tubulin (C and D) or mitochondrial outer membrane proteins (E and F). The arrow in panel C marks the meiotic spindle. n, nucleus. Bar, 2.0 μm .

collapse of mitochondrial tubules into large spherical structures.

Electron microscopy was used to characterize further mitochondrial rearrangement and structure in

meiotic wild-type and mutant cells. In ultrathin sections of wild-type cells, mitochondria were resolved as spherical or tubular structures located at the cell periphery (Figure 8). Immunofluorescence studies in

Figure 5. Mitochondrial thread formation occurs at the same rate in nocodazole-treated and untreated samples. Spheroplasts were incubated under sporulation conditions in the presence or absence of nocodazole as for Figure 4. Samples were fixed at various intervals after transfer to sporulation medium, and mitochondria and microtubules were visualized by indirect immunofluorescence as for Figure 4. Mitochondrial rearrangement was measured by determining the percent of total cells containing tubular mitochondria (circles). Microtubule integrity was measured by determining the percent cells containing intact spindles (squares). Open symbols, control samples; closed symbols, nocodazole-treated samples.

whole cells indicated that mitochondria had undergone rearrangement to form elongated tubular structures at this stage of incubation (Figure 7). Therefore, the spherical structures may represent cross-sections through tubular organelles. In addition, we observed end-to-end association of tubular mitochondria, which were interpreted as organelles undergoing collision and possible fusion. Ultrastructural analysis confirmed the increased size of *act1-3* mitochondria, and indicated that the large spherical mitochondria were two- to fivefold greater in diameter than the spherical mitochondria detected in pre-meiotic, wild-type cells (Figure 8D). Although mitochondrial size and spatial arrangement were affected in the actin mutant, mitochondrial infrastructure did not appear to be significantly affected: cristae were detectable in both enlarged and aggregated mitochondria.

Mutation within the myosin binding site of actin produces severe defects in meiosis-linked mitochondrial rearrangement. Defects in mitochondrial spatial organization occurred at both permissive and nonpermissive temperatures in the *act1-133* strain (Table 3). Although these cells also contained the spherical structures found in the *act1-3* mutant, the predominant phenotype of the *act1-133* spheroplasts was mi-

tochondrial aggregation. The nature of these mitochondrial aggregates was difficult to discern by conventional and confocal light microscopy (Figure 7); however, ultrastructural analysis confirmed that mitochondrial aggregates consist of multiple individual organelles (Figure 8). Unlike wild-type cells in which mitochondria formed a single layer at the cell cortex, mitochondrial aggregates in *act1-133* cells consisted of clusters of organelles at the cell cortex and deeper within the cell. Twenty-two percent of the *act1-133* mutant spheroplasts had mitochondrial organization defects at the onset of sporulation. After 5 h of incubation under sporulation conditions, we observed a twofold increase in the level of mitochondrial morphology defects (Table 3).

The actin mutants studied are temperature-sensitive lethal mutations that inhibit spore formation at nonpermissive temperatures. Therefore, the defects in mitochondrial arrangement observed in these cells may be due to cell death, or failure to enter meiosis I. To examine these possibilities, cell viability and spindle formation in actin mutant and wild-type cells was assessed. Both of the actin mutants were viable upon sporulation for 7.5 h at restrictive temperatures (Table 3). This finding is consistent with the observation that viability of the *act1-3* mutant at restrictive temperatures varies as a function of growth state: stationary phase and G_1 -arrested *act1-3* cells are significantly less sensitive to temperature shift than mitotic *act1-3* cells (Novick and Botstein, 1985). In addition, both mutants underwent spindle formation at the restrictive temperature (Table 3). Because spindle formation occurred in cells that do not contain buds, it is clear that entry into meiosis I occurred in wild-type cells and actin mutants at 37°C.

Mitochondrial Movement Is Impaired in Actin Mutants

Time lapse video imaging of DiOC₆-stained mitochondria in actin mutants revealed defects in meiosis-linked mitochondrial motility. We observed a 17-fold decrease in the level of mitochondrial motility in *act1-3* and *act1-133* mutants compared with wild-type cells (Table 4). Thus, mutations that delocalize the actin cytoskeleton or perturb the myosin binding site of actin inhibit mitochondrial motility. In addition, we found that mitochondrial movement was different in the actin mutants and wild-type cells. First, many mitochondrial movements in actin mutants appeared to be Brownian motions. In contrast to the linear movements that result in organelle collision and fusion observed in wild-type cells, mitochondria underwent abrupt directional changes and rarely collided in the actin mutants. In addition, although the average velocity of mitochondrial movement was not significantly affected by *act1-3* and *act1-133* mutations, we

Figure 6. Actin structures in sporulating wild-type cells and actin mutants. Wild-type (MSY106) (A and B), *act1-3* (MSY202) (C and D), and *act1-133* (DDY437) (E and F) spheroplasts were incubated in sporulation medium at 22°C (A, C, and E) or 37°C (B, D, and F). Samples were fixed after 5 h of incubation, stained with rhodamine-phalloidin, and visualized by confocal microscopy as described in MATERIALS AND METHODS. Each image was produced by superimposition of three 0.36- μ m confocal optical sections. The arrow in panel D marks an *act1-3* cell containing delocalized actin. Arrowheads, actin cables. Bar, 3.0 μ m.

observed a significantly greater range of velocities for mitochondrial movement in the actin mutants as compared with wild-type cells. These findings support a model in which mitochondrial motility and meiosis-linked rearrangement are dependent on the actin cytoskeleton.

DISCUSSION

Mitochondrial alignment, collision, and fusion are integral components of meiotic mitochondrial inheritance: this process consolidates mitochondria for equal distribution to the four developing spores. Using a

Table 3. Cell viability, spindle formation, and mitochondrial thread formation in wild-type yeast and actin mutants

Strain	Temperature (°C)	Cell viability	Spindle formation	Percent total cells		
				Mitochondrial		Actin delocalization
				Rearrangement	Defects	
MSY106	22	100	95.3	95.3	1.9	4.0
(<i>ACT1</i>)	37	83.5	86.9	80.8	13.8	2.5
MSY202	22	100	94.0	85.9	9.3	10.4
(<i>act1-3</i>)	37	81.1	71.3	26.6	70.9	30.9
DDY437	22	100	91.8	20.8	37.8	2.9
(<i>act1-133</i>)	37	76.5	74.0	31.2	41.2	5.9

Late log phase actin mutants and wild-type yeast were converted to spheroplasts and incubated in sporulation medium at 22°C and 37°C as for Figure 5. After 5 h of incubation, aliquots of each culture were fixed. Mitochondria and microtubule-containing structures were visualized as for Figure 3. Spindle formation was evaluated by determining the percentage of cells that contained bar-shaped microtubule-containing structures. Mitochondrial rearrangement was evaluated by determining the percentage of cells containing elongated, tubular mitochondria. Mitochondrial defects were evaluated by determining the percentage of cells containing enlarged, aggregated, or fragmented mitochondria. The actin cytoskeleton was defined as delocalized in cells if actin patches and cables were not detectable using rhodamine-phalloidin stain. To determine cell viability, aliquots of each culture were removed after 7.5 h of incubation, diluted with osmotically stabilized media, and grown on rich, glucose-containing solid medium at 22°C. Analysis of colony recovery on solid media indicates that viability assay conditions had no significant effect of spheroplast viability. The numbers shown are the number of colonies from 37°C sporulation cultures compared with those obtained from 22°C sporulation cultures.

membrane potential sensing dye (DiOC₆) and time lapse fluorescence microscopy to examine mitochondrial movements during meiosis-linked mitochondrial rearrangement, we conclude that mitochondrial movement is regulated. Fifteen percent of the mitochondria within pre-meiotic stationary phase yeast display movement. However, entry into meiosis I results in a threefold increase in the number of motile mitochondria. Movements observed in meiotic yeast appear to be directed; mitochondria display linear, continuous movements with an average velocity of 25 nm/s. Finally, these movements result in the collision and fusion of mitochondria to form elongated, tubular organelles.

Analysis of the distribution of microtubules and mitochondria, and of the effect of microtubule destabilization on mitochondrial motility and rearrangement indicates that microtubules do not play a major role in this process. In contrast, under restrictive conditions, *act1-3* cells display actin delocalization, a 17-fold decrease in mitochondrial motility, and 60% inhibition in mitochondrial thread formation. Several observations indicate that these defects are direct effects of the *act1-3* mutation rather than indirect effects on cell viability. First, in contrast to vegetative *act1-3* cells, which lose viability after 1–2 h of incubation at restrictive temperatures (Novick and Botstein, 1985), *act1-3* mutants are viable up to 7.5 h at restrictive temperatures under the sporulation conditions used. In addition, it is clear that the *act1-3* strain can enter meiosis I: this mutant undergoes spindle formation to the same extent at permissive and restrictive temperatures. Finally, we previously reported that mitochon-

dria isolated from vegetative yeast display direct reversible ATP-sensitive binding to F-actin, in reactions that are mediated by an actin-dependent motor activity on the mitochondrial outer membrane (Lazzarino *et al.*, 1994; Simon *et al.*, 1995). Therefore, our interpretation of these observations is as follows: 1) mitochondria use actin filaments as tracks for directed movement during meiosis, and 2) perturbation of the actin cytoskeleton in *act1-3* cells disrupts actin-mitochondrial interactions and results in loss of mitochondrial motility.

In addition, sporulation of *act1-3* cells at restrictive temperatures results in formation of large spherical mitochondria. These mitochondria are similar to those observed upon mutation of *YME1* (Campbell *et al.*, 1994), as well as *MDM10* and *MMM1*, genes encoding mitochondrial outer membrane proteins implicated in mitochondrial inheritance in mitotic yeast (Burgess *et al.*, 1994; Sogo and Yaffe, 1994). In the previous studies, the authors suggested that *MDM10* and *MMM1* are required for organelle-cytoskeletal interactions and that mutations of these genes disrupt these interactions, effectively altering mitochondrial shape. Potentially, delocalization of the actin cytoskeleton in the *act1-3* mutant may also release mitochondria from actin structures and cause collapse of tubular organelles into spheres of various diameters.

Studies in many laboratories indicate that myosin, the barbed-end directed actin-dependent motor molecule, utilizes the energy of ATP hydrolysis to drive organelle movement along the lateral surface of microfilaments. This model is based on the following findings: 1) muscle and nonmuscle myosins are capa-

Figure 7. Actin mutants display defects in meiosis-linked mitochondrial rearrangements. Wild-type (A and B), *act1-3* (C and D), and *act1-133* (E and F) spheroplasts were incubated at either 22°C (A, C, and E) or 37°C (B, D, and F) as for Figure 6. Samples were fixed after 5 h of incubation and mitochondria were labeled by indirect immunofluorescence and visualized by confocal microscopy. Each image was produced by superimposition of three 0.36- μm confocal optical sections. The arrows in panel D mark enlarged mitochondria in *act1-3* cells incubated at 37°C. The arrows in panels E and F mark aggregated mitochondria in the *act1-133* cells at permissive and nonpermissive temperatures. Bar, 3.0 μm .

ble of driving ATP-dependent particle movement along actin filaments (Sheetz and Spudich, 1983; Albanesi *et al.*, 1985); 2) proteins that are immunologically cross-reactive with myosin are associated with vesicles and organelles that bind to or move along

microfilaments (Burrige and Phillips, 1975; Grolig *et al.*, 1988; Fath and Burgess, 1993); 3) antibodies raised against myosin block organelle movement along actin bundles in vitro (Adams and Pollard, 1986); and 4) mutations in myosin genes inhibit particle movement

Figure 8. Ultrastructure of mitochondria in sporulating wild-type cells and actin mutants. Wild-type (A and B), *act1-3* (C and D), and *act1-133* (E and F) yeast strains were converted to spheroplasts and incubated in sporulation medium at permissive and nonpermissive conditions as for Figure 6. After 5 h of incubation, samples were fixed and prepared for ultrastructural analysis as described in MATERIALS AND METHODS. (A, C, and E) Samples from 22°C incubations. (B, D, and F) Samples from 37°C incubations. (A) Tubular mitochondria that are alone (arrow) or in end-to-end arrangements with other mitochondria (arrowhead) at the cell periphery in wild-type cells. (D) Arrowhead marks enlarged mitochondrion produced during sporulation of *act1-3* cells at nonpermissive temperatures. (E and F) Arrowheads mark mitochondrial aggregates in *act1-133* cells. Bar, 1.0 μm .

Table 4. Effect of microtubule-destabilization and actin mutations on mitochondrial movement

Strain	Motile mitochondria (%)	Velocity (nm/s)	n
MSY106 (<i>ACT1</i>)	48.8	26 ± 4.3	67
MSY106 + nocodazole	51.5	29 ± 7.9	78
MSY202 (<i>act1-3</i>)	2.6	31 ± 20.0	48
DDY437 (<i>act1-133</i>)	2.6	41 ± 24.0	66

Sporulation of yeast spheroplasts, DiOC₆-staining of mitochondria, and time lapse video microscopy to visualize mitochondrial movements were carried out as for Table 2. Microtubule destabilization using nocodazole in the wild strain (MSY106) was carried out as for Figure 4. The velocity of mitochondrial movement and % motile mitochondria were determined as for Table 2. The values shown are mean values from three experiments. n = number of moving particles analyzed for velocity determinations.

in *Dictyostelium* and budding yeast (Wessels and Soll, 1990; Johnston *et al.*, 1991; Govindan *et al.*, 1995).

We find that mitochondria isolated from vegetative yeast contain an ATP-driven, actin-dependent motor activity (Simon *et al.*, 1995). However, mutation of the known yeast myosin genes has no significant effect on mitochondrial motility in vegetative yeast (Simon *et al.*, 1995). To examine the possible role of actin-myosin interactions in meiosis-linked mitochondrial motility, we studied the effect of the *act1-133* allele. This mutant contains amino acid substitutions of alanine for D24 and D25, residues within the myosin binding site of actin. Johara *et al.* (1993) found that *Dictyostelium* actin containing histidine in place of D24 and D25 binds to myosin but will not support ATP-driven heavy meromyosin sliding in vitro. Phenotypic analysis of the *act1-133* mutation in mitotic yeast reveals low growth rates, enlarged cell size, multiple buds, randomized bud site selection, mislocalized actin patches, and mitochondrial aggregation at permissive temperature (Drubin *et al.*, 1993). We find that this allele has no significant effect on entry into meiosis I: *act1-133* cells are viable and display spindle formation under permissive and restrictive conditions. In addition, the *act1-133* allele has no significant effect on actin patches and cables upon sporulation at permissive and non-permissive temperatures. However, we observe mitochondrial aggregation, loss of all directed mitochondrial movement, and defects in mitochondrial rearrangement upon sporulation of *act1-133* cells at either permissive or restrictive temperatures. Upon long term incubation of the *act1-133* mutant at elevated temperatures, we observed loss of tubular mitochondria and accumulation of large spherical mitochondria. These findings support a role for actin-myosin interactions in control of meiosis-linked

mitochondrial movement, rearrangement, and morphology.

The velocity of organelle movement described here, 25 nm/s (Table 2), is 6–30 times lower than actin-driven organelle motility in systems including *Dictyostelium*, extruded squid axoplasm, and cell-free *Nitella* systems (Adams and Pollard, 1986; Wessels *et al.*, 1989; Kuznetsov *et al.*, 1992). Because sporulation is a starvation response, the low rates of mitochondrial movement may reflect the nutrient-limited state of the cell samples. However, several findings indicate that the rates of mitochondrial movements may reflect inherently lower motility activity of elements of the yeast cytoskeleton. First, using the microfilament sliding assay to measure myosin motor activity, Kron *et al.* (1992) found that yeast actin displays low affinity for, and low sliding rates along, rabbit skeletal muscle myosin. Second, we found that the velocity of actin-dependent mitochondrial movement in vegetative yeast, 49 nm/s, is not significantly different than that of meiosis-linked mitochondrial movement (Simon *et al.*, 1995). Finally, the velocity of mitochondrial motor-driven actin-filament sliding, 25 nm/s, is similar to those of mitotic and meiotic mitochondrial movement in living cells (Simon *et al.*, 1995).

These studies document regulated mitochondrial movements in meiotic yeast, and suggest a role for the actin cytoskeleton in control of meiosis-linked mitochondrial motility and morphology. In addition, we propose a mechanism for mitochondrial inheritance in meiotic yeast: mitochondria bind to actin for control of organelle shape, and utilize actin cytoskeletal elements as tracks for regulated and directed movement.

ACKNOWLEDGMENTS

We thank G. Schatz, D. Drubin, K. Wertman, and L. Symington for yeast strains, antibodies, and invaluable suggestions; and A. Mikhailov and G.G. Gundersen for assistance with video microscopy. We also thank R. Ghosh, K. Dunn, and F. Maxfield for their assistance with confocal microscopy; and P.W. Brandt and A.J. Silverman for critical review of the manuscript. This work has been supported by a National Research Service Award (1 F31 GM15644) to V.S., a Junior Faculty Research Award (ACS JFRA-377) from the American Cancer Society, the Basil O'Connor Starter Scholar Research Award (MDBDF #5-FY92-1143) from the March of Dimes, a Grant-in-aid Award (AHA CU50706601) from the American Heart Association, and research grants from the American Cancer Society (ACS CB-41) and the National Institutes of Health (1 RO1 GM45735).

REFERENCES

- Adams, A.E.M., Botstein, D., and Drubin, D.G. (1991). Requirement of yeast fimbrin for actin organization and morphogenesis in vivo. *Nature* 354, 404–408.
- Adams, A.E.M., and Pringle, J. (1984). Relationship of actin and tubulin distribution in wild-type and morphogenetic mutant *Saccharomyces cerevisiae*. *J. Cell Biol.* 98, 934–945.

- Adams, R.J., and Pollard, T.D. (1986). Propulsion of organelles isolated from *Acanthamoeba* along actin filaments by myosin I. *Nature* 322, 754–756.
- Albanesi, J.P., Fujisaki, J., Hammer, J.A., III, Korn, E.D., Jones, R., and Sheetz, M.P. (1985). Monomeric *Acanthamoeba* myosins-I support movement in vitro. *J. Biol. Chem.* 260, 8649–8652.
- Allen, R.D., Metuzals, J., Tasake, I., Brady, S.T., and Gilbert, S.P. (1982). Fast axonal transport in squid giant axon. *Science* 218, 1127–1128.
- Amatruda, J.F., and Cooper, J.A. (1992). Purification, characterization, and immunofluorescence localization of *Saccharomyces cerevisiae* capping protein. *J. Cell Biol.* 117, 1067–1076.
- Brady, S.T., Lasek, R.J., and Allen, R.D. (1982). Fast axonal transport in extruded axoplasm from squid giant axon. *Science* 218, 1129–1131.
- Bridgman, P.C., Kachar, B., and Reese, T.S. (1986). The structure of cytoplasm in directly frozen cultured cells. II. Cytoplasmic domains associated with organelle movements. *J. Cell Biol.* 102, 1510–1521.
- Burgess, S.M., Delannoy, M., and Jensen, R.E. (1994). *MMM1* encodes a mitochondrial outer membrane protein essential for establishing and maintaining the structure of yeast mitochondria. *J. Cell Biol.* 126, 1375–1391.
- Burridge, K., and Phillips, J.H. (1975). Association of actin and myosin with secretory granule membranes. *Nature* 254, 526–529.
- Byers, B. (1981). Cytology of the yeast life cycle. In: *The Molecular Biology of the Yeast Saccharomyces: Life Cycle and Inheritance*, ed. J.N. Strathern, E.W. Jones, and J.R. Broach, Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 59–96.
- Campbell, C.L., Tanaka, N., White, K.H., and Thorsness, P.E. (1994). Mitochondrial morphological and functional defects in yeast caused by *yme1* are suppressed by mutation of a 26S protease subunit homologue. *Mol. Biol. Cell* 5, 899–905.
- Chowdhury, S., Smith, K.W., and Gustin, M.C. (1992). Osmotic stress and the yeast cytoskeleton: phenotype-specific suppression of an actin mutation. *J. Cell Biol.* 118, 561–571.
- Cooper, J.A. (1987). Effects of cytochalasin and phalloidin on actin. *J. Cell Biol.* 105, 1473–1478.
- Drubin, D.G., Jones, H.D., and Wertman, K.F. (1993). Actin structure and function: roles in mitochondrial organization and morphogenesis in budding yeast and identification of the phalloidin-binding site. *Mol. Biol. Cell* 4, 1277–1294.
- Drubin, D.G., Miller, K.G., and Botstein, D. (1988). Yeast actin-binding proteins: evidence for a role in morphogenesis. *J. Cell Biol.* 107, 2551–2561.
- Esposito, R.E., and Klapholz, S. (1981). Meiosis and ascospore development. In: *The Molecular Biology of the Yeast Saccharomyces: Life Cycle and Inheritance*, ed. J.N. Strathern, E.W. Jones, and J.R. Broach, Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 211–287.
- Fath, K.R., and Burgess, D.R. (1993). Golgi-derived vesicles from developing epithelial cells bind actin filaments and possess myosin-I as a cytoplasmically oriented peripheral membrane protein. *J. Cell Biol.* 120, 117–127.
- Govindan, B., Bowser, R., and Novick, P. (1995). The role of *MYO2*, a yeast class V myosin, in vesicular transport. *J. Cell Biol.* 128, 1055–1068.
- Grolig, F., Williamson, R.E., Parke, J., Miller, C., and Anderton, B.H. (1988). Myosin and calcium-sensitive streaming in the alga *Chara*: detection of two polypeptides reacting to a monoclonal anti-myosin and their localization in the streaming endoplasm. *Eur. J. Cell Biol.* 47, 22–31.
- Guthrie, B.A., and Wickner, W. (1988). Yeast vacuoles fragment when microtubules are disrupted. *J. Cell Biol.* 107, 115–120.
- Hayden, J.H., and Allen, R.D. (1984). Detection of single microtubules in living cells: particle transport can occur in both directions along a single microtubule. *J. Cell Biol.* 99, 1785–1793.
- Hegmann, T.E., Lin, J.L., and Lin, J.J. (1989). Probing the role of non-muscle tropomyosin isoforms in intracellular granule movement by microinjection of monoclonal antibodies. *J. Cell Biol.* 109, 1141–1152.
- Hegmann, T.E., Schulte, D.L., Lin, J.L., and Lin, J.J. (1991). Inhibition of intracellular movement by microinjection of monoclonal antibodies against caldesmon. *Cell Motil. Cytoskeleton* 20, 109–120.
- Hirokawa, N. (1982). Cross-linker system between neurofilament, microtubules and membranous organelles in frog axons revealed by the quick-freeze, deep-etching method. *J. Cell Biol.* 27, 255–272.
- Jacobs, C.J., Adams, A.E.M., Szanislo, P.J., and Pringle, J.R. (1988). Functions of microtubules in the *Saccharomyces* cell cycle. *J. Cell Biol.* 107, 1409–1426.
- Johara, M., Toyoshima, Y.Y., Ishijima, A., Kojima, H., Yanagida, T., and Sutoh, K. (1993). Charge-reversion mutagenesis of *Dictyostelium* actin to map the surface recognized by myosin during ATP-driven sliding motion. *Proc. Natl. Acad. Sci. USA* 90, 2127–2131.
- Johnston, G.C., Prendergast, J.A., and Singer, R.A. (1991). The *Saccharomyces cerevisiae* *MYO2* gene encodes an essential myosin for vectorial transport of vesicles. *J. Cell Biol.* 113, 539–551.
- Kabsch, W., Mannherz, H.G., Suck, D., Pai, E.F., and Holmes, K.C. (1990). Atomic structure of the actin:DNAseI complex. *Nature* 347, 37–44.
- Kachar, B. (1985). Direct visualization of organelle movement along filaments dissociated from *Characean* algae. *Science* 227, 1355–1357.
- Kachar, B., and Reese, T.S. (1988). The mechanism of cytoplasmic streaming in *Characean* cells: sliding of endoplasmic reticulum along actin filaments. *J. Cell Biol.* 106, 1545–1552.
- Kilmartin, J.V., and Adams, A.E.M. (1984). Structural rearrangements of tubulin and actin during cell cycle of *Saccharomyces cerevisiae*. *J. Cell Biol.* 98, 922–933.
- Kohn, T., and Shimmer, T. (1988). Accelerated sliding of pollen tube organelles along *Characeae* actin bundles regulated by Ca^{++} . *J. Cell Biol.* 106, 1539–1543.
- Koning, A.J., Lum, P.Y., Williams, J.M., and Wright, R. (1993). DiOC₆ staining reveals organelle structure and dynamics in living yeast cells. *Cell Motil. Cytoskeleton* 25, 111–128.
- Koonce, M.P., and Schliwa, M. (1985). Bidirectional organelle transport occurs in cell processes that contain single microtubules. *J. Cell Biol.* 100, 322–326.
- Kron, S.J., Drubin, D.G., Botstein, D., and Spudich, J.A. (1992). Yeast actin filaments display ATP-dependent sliding movement over surfaces coated with rabbit muscle myosin. *Proc. Natl. Acad. Sci. USA* 89, 4466–4470.
- Kübler, E., and Riezman, H. (1993). Actin and fimbrin are required for the internalization step of endocytosis in yeast. *EMBO J.* 12, 2855–2862.
- Kuznetsov, S.A., Langford, G.M., and Weiss, D.G. (1992). Actin-dependent organelle movement in squid axoplasm. *Nature* 356, 722–725.
- Lazzarino, D., Boldogh, I., Smith, M.G., Rosand, J., and Pon, L.A. (1994). Yeast mitochondria contain ATP-sensitive, reversible actin-binding activity. *Mol. Biol. Cell* 5, 807–818.
- Liu, H., and Bretscher, A. (1992). Characterization of *TPM1* disrupted yeast cells indicates an involvement of tropomyosin in directed vesicular transport. *J. Cell Biol.* 118, 285–299.

- Menzel, D., and Schliwa, M. (1986). Motility in the siphonous green alga *Bryopsis*. I. Spatial organization of the cytoskeleton and organelle movements. *J. Cell Biol.* 40, 275–285.
- Miyakawa, I., Aoi, H., Sando, N., and Kuroiwa, T. (1984). Fluorescence microscopic studies of mitochondrial nucleoids during meiosis and sporulation in the yeast, *Saccharomyces cerevisiae*. *J. Cell Sci.* 66, 21–38.
- Moens, P.B., and Rapport, E. (1971). Spindles, spindle plaques and meiosis in the yeast *Saccharomyces cerevisiae* (Hansen). *J. Cell Biol.* 50, 344–361.
- Moon, A.L., Janmey, P.A., Louie, K.A., and Drubin, D.G. (1993). Cofilin is an essential component of the yeast cortical cytoskeleton. *J. Cell Biol.* 120, 421–435.
- Mulholland, J., Preuss, D., Moon, A., Wong, A., Drubin, D., and Botstein, D. (1994). Ultrastructure of the yeast actin cytoskeleton and its association with the plasma membrane. *J. Cell Biol.* 125, 381–391.
- Novick, P., and Botstein, D. (1985). Phenotypic analysis of temperature-sensitive yeast actin mutants. *Cell* 40, 405–416.
- Palmer, R.E., Sullivan, D.S., Huffaker, T., and Koshland, D. (1992). Role of astral microtubules and actin in spindle orientation and migration in the budding yeast, *Saccharomyces cerevisiae*. *J. Cell Biol.* 119, 583–593.
- Pollard, T.D., Doberstein, S.K., and Zot, H.G. (1991). Myosin-I. *Annu. Rev. Physiol.* 53, 653–681.
- Pringle, J.R., Preston, R.A., Adams, A.E.M., Stearns, T., Drubin, D.G., and Haarer, B.K. (1989). Fluorescence microscopy methods for yeast. *Methods Cell Biol.* 31, 357–435.
- Rayment, I., Holden, H.M., Whittaker, M., Yohn, C.B., Lorenz, M., Holmes, K.C., and Milligan, R.A. (1993). Structure of the actin-myosin complex and its implications for muscle contractions. *Science* 261, 58–65.
- Read, E.B., Okamura, R.H., and Drubin, D.G. (1992). Actin-and tubulin-dependent functions during *Saccharomyces cerevisiae* mating projection and formation. *Mol. Biol. Cell* 3, 429–444.
- Riezman, H., Hay, R., Gasser, S., Daum, G., Schneider, G., Witte, C., and Schatz, G. (1983). The outer membrane of yeast mitochondria: isolation of outside-out sealed vesicles. *EMBO J.* 2, 1105–1111.
- Rose, M.D., and Fink, G.R. (1987). KAR1, a gene required for function of both intranuclear and extranuclear microtubules in yeast. *Cell* 48, 1047–1060.
- Schroer, T.A., and Sheetz, M.P. (1991). Functions of microtubule-based motors. *Annu. Rev. Physiol.* 53, 629–652.
- Sheetz, M.P., and Spudich, J.A. (1983). Movement of myosin-coated fluorescent beads on actin cables in vitro. *Nature* 303, 31–35.
- Shortle, D., Novick, P., and Botstein, D. (1984). Construction and genetic characterization of temperature-sensitive mutant alleles of yeast actin gene. *Proc. Natl. Acad. Sci. USA* 81, 4889–4893.
- Simon, V.R., Swayne, T.C., and Pon, L.A. (1995). Actin-dependent mitochondrial motility in mitotic yeast and cell-free systems: identification of a motor activity on the mitochondrial surface. *J. Cell Biol.* 130, 345–354.
- Sogo, L.F., and Yaffe, M.P. (1994). Regulation of mitochondrial morphology and inheritance by Mdm10p, a protein of the mitochondrial outer membrane. *J. Cell Biol.* 126, 1361–1373.
- Stevens, B. (1977). Variation in number and volume of the mitochondria in yeast according to growth conditions: a study based on serial sectioning and computer graphics reconstruction. *Biol. Cell* 28, 37–56.
- Stevens, B. (1981). Mitochondrial structure. In: *The Molecular Biology of the Yeast Saccharomyces: Life Cycle and Inheritance*, ed. J.N. Strathern, E.W. Jones, and J.R. Broach, Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 471–488.
- Titus, M.A. (1993). Myosins. *Curr. Opin. Cell Biol.* 5, 77–81.
- Uchida, A., and Suda, K. (1978). Distribution of mitochondrially inherited drug-resistance genes to tetrads from young zygotes in yeast. *Mol. Gen. Genet.* 165, 191–197.
- Wertman, K.F., Drubin, D.G., and Botstein, D. (1992). Systematic mutational analysis of the yeast *ACT1* gene. *Genetics* 132, 337–350.
- Wessels, D., Schroeder, N.A., Voss, E., Hall, A.L., Condeelis, J., and Soll, D.R. (1989). cAMP-mediated inhibition of intracellular particle movement and actin reorganization in *Dictyostelium*. *J. Cell Biol.* 109, 2841–2851.
- Wessels, D., and Soll, D.R. (1990). Myosin II heavy chain null mutant of *Dictyostelium* exhibits defective intracellular particle movement. *J. Cell Biol.* 111, 1137–1148.
- Williamson, D.H., and Fennell, D.J. (1975). The use of fluorescent DNA-binding agent for detecting and separating yeast mitochondrial DNA. *Methods Cell Biol.* 12, 335–351.