Published in final edited form as:

Arch Otolaryngol Head Neck Surg. 2010 June ; 136(6): 634–636.

Pathology Quiz Case 1

Dr Sara B. Immerman, MD,

New York University School of Medicine, New York, New York

Dr Luc G. T. Morris, MD,

New York University School of Medicine, New York, New York

Dr Mark D. DeLacure, MD,

New York University School of Medicine, New York, New York

Dr Beverly Y. Wang, MD, and

New York University School of Medicine, New York, New York

Dr Zoya Kurago, DDS, PhD

New York University College of Dentistry, New York, New York

An 89-year-old white woman with a history of schizophrenia presented with a 1-year history of progressive left facial swelling accompanied by local bleeding, pain, and dysphagia. A review of systems revealed a 9-kg unintentional weight loss and lower lip paresthesias. She denied trismus, fever, trauma, and recent dental work. She did not use tobacco or alcohol. Physical examination revealed a 6-cm deforming mass in the left mandibular alveolar ridge that extended into the floor of the mouth and overlying skin, causing ulceration and bleeding. There were several “floating” teeth in the left anterior and posterior mandible and anesthesia over the left mental nerve distribution. The results of the rest of the head and neck examination were normal. Computed tomography revealed an expansile, ill-defined lytic lesion in the body of the left mandible, with extensive erosion into the buccal cortex (Figure 1).

The findings of fine-needle aspiration of the mass were inconclusive, and the results of an incisional biopsy were consistent with a malignant epithelial neoplasm of odontogenic origin. Final histopathologic examination of the composite resection revealed a mass that was composed of epithelial cells with intraosseous and extraosseous components. The intraosseous portion demonstrated variably sized nests of epithelial cells with a variety of patterns, most notably plexiform (Figure 2) and spindling. Other nests showed prominent peripheral columnar cells with nuclear palisading away from the basement membrane and stellate reticulumlike architecture of the central areas (Figure 3 and Figure 4). Nuclear pleomorphism (Figure 3 and Figure 4), atypical mitotic figures, and necrosis, all features that support an impression of malignancy, were present throughout the tumor nests.

What is your diagnosis?

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Figure 1.

Figure 2.

Figure 3.

Figure 4.