Adapting North American wheat production to climatic challenges, 1839–2009

Alan L. Olmstead^a and Paul W. Rhode^{b,1}

^aDepartment of Economics, University of California, Davis CA 95616; and ^bDepartment of Economics, University of Michigan, Ann Arbor, MI 48109

Edited* by Richard A. Easterlin, University of Southern California, Los Angeles, CA, and approved November 12, 2010 (received for review June 10, 2010)

The Intergovernmental Panel on Climate Change projects that temperatures in the major grain-growing areas of North America will rise by 3–4 °C by 2100. Such abrupt changes will create major challenges, significantly altering the area suitable for wheat. The historical record offers insight into the capability of agriculture to adapt to climatic challenges. Using a new county-level dataset on wheat production and climate norms, we show that during the 19th and 20th centuries North American grain farmers pushed wheat production into environments once considered too arid, too variable, and too harsh to cultivate. As summary measures, the median annual precipitation norm of the 2007 distribution of North American wheat production was one-half that of the 1839 distribution, and the median annual temperature norm was 3.7 °C lower. This shift, which occurred mostly before 1929, required new biological technologies. The Green Revolution associated with the pioneering work of Norman Borlaug represented an important advance in this longer process of biological innovation. However, well before the Green Revolution, generations of North American farmers overcame significant climatic challenges.

adaptation | agricultural technology | food security

The Intergovernmental Panel on Climate Change (IPCC) projects that by 2100 annual mean temperatures in North America will increase by 2 to 3 °C at midlatitude coastal regions and by “up to more than” 5 °C at more northern latitudes. In the main grain-growing areas, the IPCC forecasts temperatures will rise 3 to 4 °C (1). A more recent MIT study suggests far greater changes will occur (2). There are a wide range of estimates of how climate changes will impact agricultural production (3–7). Numerous researchers have speculated about how farmers might change cultivars, cropping patterns, and farming methods to mitigate some of the costs of abrupt climatic changes (8). Researchers at the International Maize and Wheat Improvement Center (CIMMYT) anticipate that North American wheat farmers may extend the margin of wheat production roughly 1,000 km north into northern Canada and Alaska, whereas heat and drought will make cultivation untenable in many areas of the southern Great Plains (9). To provide perspective on these and other predictions, this paper asks how farmers responded to past climatic challenges.

The spread of wheat cultivation across North America required that farmers repeatedly adapt to unfamiliar and hostile climatic conditions. The variations in climatic conditions that settlers encountered rivaled the magnitude of the predicted changes at given locations over the next century. We quantified the extent of the geographic variations and decipher how wheat growers learned to produce in new environments. Because of the paucity of Mexican data before 1929, most of our analysis of “North America” refers to Canada and the United States. Inclusion of Mexico in the later part of the 20th century highlights the role of the Green Revolution in pushing production into hotter and drier zones.

Results

The quantification of the geoclimatic conditions of wheat production between 1839 and 2007 shows that farmers made striking changes to adjust to climate; most of the changes in location and the technological adjustments that made settlement possible occurred before the Green Revolution.

Quantifying Geographic Changes in Production and Climate. Between 1839 and 2009, wheat output increased 26-fold in the United

States and more than 270-fold in Canada. In 1839, the geographic center (mean) of North American wheat production was located in eastern Ohio. Cultivation was concentrated in Ohio and New York; relatively little wheat was grown as far west as Illinois. In 2007 the center of production had moved 1,800 km west, into west-central South Dakota (*Methods*). Almost all this movement occurred when plant sciences were in their infancy.

The change in the location entailed large shifts in growing conditions. The six panels of Fig. 1 display the main features of the changing geographic distribution of the North American wheat crop across latitude; longitude; annual, January, and July temperature norms; and annual precipitation norms. The series cover the period 1839–2007, using county-level information from United States and Canada. The distributions summarized in Fig. 1 weight the fixed county-level geoclimatic characteristics by output in each locality at each date (*Methods*).

Fig. 1A summarizes the changing longitude of wheat production in North America over roughly 170 y. The median production shifted west by 21° of longitude (nearly 1,800 km) between 1839 and 1929, with little movement thereafter. By 1879, the median was beyond the extreme western boundary of production in 1839. The median latitude of production (Fig. 1B) was relatively constant until the 1890s, when the northern Plains and the Canadian Prairies began to enter cultivation. The median latitude of production in 1929 was near the northern fringe of production (the 95% line) in 1839. The northernmost one-quarter of production (reflected by the 75% line) moved north by 8° of latitude (>880 km) between 1839 and 1929.

Fig. 1C–F quantifies the changes in climatic conditions associated with these geographic shifts in production. In 1839 median production took place in an environment with a (1941–1970) norm of nearly 100 cm of precipitation (Fig. 1C). In 2007, median production took place on land with a norm of less than 50 cm of precipitation; this environment was drier than virtually any place growing wheat in the United States or Canada in 1839. Almost all the changes in the distribution of production, as measured by annual precipitation, had occurred by 1929. In that year the marginal fringe (the 10% line) with 35 cm or less of precipitation produced about one-fifth more wheat than North America’s total output in 1839. The range of annual moisture conditions widened substantially, as indicated by the growing spread between the 10% and 90% lines. Data on precipitation in January and July (Figs. S1 and S2) show that the decline was apparent across the year, creating an array of challenges in new production zones. As an indicator of the extent of the changes, the driest 10% of North American production moved from areas with 7.8 cm of rain in July in 1839 to areas that averaged 0.9 cm in 1889.

The median annual and January temperature norms fell by 3.7 °C and 5.9 °C, respectively, between 1839 and 2007 (Fig. 1D and E). The range of temperature conditions widened greatly, with a pronounced movement into colder domains. The 90–10 percentile differential in annual temperature doubled, from 6.3 to 13.1 °C,

Author contributions: A.L.O. and P.W.R. designed research; A.L.O. and P.W.R. performed research; A.L.O. and P.W.R. contributed new reagents/analytic tools; A.L.O. and P.W.R. analyzed data; and A.L.O. and P.W.R. wrote the paper.

The authors declare no conflict of interest.

*This Direct Submission article had a prearranged editor.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. E-mail: pwrhode@umich.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1008279108/-DCSupplemental.

SOCIAL SCIENCES

remind us of the revolutionary accomplishments that he and other agronomists made during the past 70 y. However, in 1944, when Borlaug started his work for the Rockefeller Foundation in Mexico, there already was a long history of agriculturalists who had successfully improved wheat to meet climatic conditions.

Biological Innovation to Adapt to Environmental Challenges. Wheat cultivation was introduced into Mexico in 1521, but it did not appear in the territory that would become Canada and the United States until 1602 (10). Discovering wheats suitable for new areas was a reoccurring struggle. Farmers in eastern Canada and New England continuously experimented to find cold-tolerant and pest-resistant wheats (11). In the more temperate Middle colonies, the cultivars (cultivated varieties) transplanted from Western Europe fared better. However, the challenges were particularly acute when pioneers moved wheat cultivation westward onto the northern Prairies, Great Plains, and Pacific Coast. All these regions eventually became major wheat suppliers, but only after farmers learned to overcome climatic conditions far different from those prevailing to the East and in Western Europe. The initial attempts to grow traditional wheat cultivars frequently failed.

The experiences of the Selkirk colonists who settled near Lake Winnipeg offer an example. The winter wheat, first tried in 1811–1812, failed. Successive crops of spring wheat also succumbed to drought, freezing, and insects. To obtain sufficient seed for the 1820 crop, a band of Selkirk settlers had to trek more than 2,000 km (round trip) to Prairie du Chien on the upper Mississippi River during the dead of winter. After about a decade of hungry times, the colony began to sustain itself (12). The prolonged troubles of the Selkirk colonists represented a clear case of settlers leapfrogging beyond the limits of their climatic knowledge. However, even when settlers inched west in a more orderly fashion, the challenge of adapting was daunting. In the 1840s, attempts to grow soft winter wheat on the Wisconsin Prairie failed repeatedly, and wheat culture succeeded only after farmers switched to a new hard spring wheat cultivar (13).

The Great Plains were depicted as the “Great American Desert” and were considered incapable of supporting agriculture. The first waves of settlers from the humid East and Midwest moved into the High Plains during the relatively wet years of the 1880s. These farmers, along with railroad and government officials, significantly miscalculated the climatic obstacles that had to be overcome (14). Success required decades of experimentation and frequently depended on knowledge and cultivars introduced by immigrants from frigid and arid locales of Eurasia.

The spread of wheat cultivation across the Great Plains and Canadian Prairies was made possible by mechanization and the extension of railroads. A host of mechanical innovations cut in half the labor required to produce a unit of wheat between 1840 and 1913. In the 1860s, railroad builders began expanding into this region; by 1913, about 133,000 km of track in the Plains and Prairies served to link farmers to expanding distant markets (15, 16). However, the expansion of wheat cultivation also depended on the introduction and breeding of hard red winter and spring wheats that were entirely new to North America. During the late 19th century, the premier hard spring wheat cultivated in North America was *Red Fife* (which probably made its way from Eastern Europe to Ontario via Scotland). According to the standard account, David and Jane Fife of Otonabee, ON, selected and increased the grain stock from a single wheat plant grown on their farm in 1842. It was not introduced into the United States until the mid-1850s. *Red Fife* was the first hard spring wheat grown in North America and became the basis for the spread of wheat into Wisconsin, Minnesota, the Dakotas, and Canadian Prairies (13).

Another notable breakthrough was the introduction of *Turkey* wheat, which, like *Red Fife*, allowed production to shift into more marginal environments (*SI Text*). This hard red winter cultivar was especially well suited to the harsh growing conditions of the southern wheat belt. The standard histories credit German Mennonites, who migrated from southern Russia to Kansas, with importing *Turkey* in 1873. This transfer was far from haphazard, because local railroads recruited these migrants for their knowledge of farming in such environments (ref. 16, pp. 88–89.). Furthermore, before departing for Kansas, the Mennonites tediously

selected high-quality seed considered suitable for the new lands. Earlier Kansas settlers experimented with soft winter cultivars common to the eastern states, but these wheats proved unreliable in the cold winters and hot, dry summers. Tests at the Kansas Agricultural Experiment Station (AES) demonstrated *Turkey*'s superiority and helped popularize the wheat. In 1919, *Turkey*-type wheat made up more than 80% of the wheat acreage in Nebraska and Kansas and nearly 70% of the wheat acreage in Colorado and Oklahoma (17, 18 and ref. 19, pp. 208–218). At this time, S. C. Salmon, a leading US Department of Agriculture (USDA) breeder, concluded that, without *Turkey* cultivars, “The wheat crop of Kansas today would be no more than half what it is, and the farmers of Nebraska, Montana and Iowa would have no choice but to grow spring wheat” (20).

In both Canada and the United States many varietal innovations were the result of government investments. In 1886 Parliament created the Canadian Federal Experiment Station system. Its most acclaimed breeder, William Saunders, commenced a systematic program of hybridizing high-quality cultivars with early-maturing wheats introduced from around the world. In 1903 his son, Charles Saunders, took over the work at the Dominion Experimental Farm, near Ottawa. The most valuable result of their combined research efforts was *Marquis*, a cross between *Red Fife* and *Red Calcutta*, a very early wheat from India. Released in 1909, *Marquis* was an immediate success and accounted for about 90% of Canada's wheat acreage in 1920 (21). After testing the wheat in its network of experiment stations, the USDA released the spring wheat in 1912–1913. By 1919, this cultivar accounted for 17% of United States acreage, and its range stretched from Washington State to northern Illinois (19). The rapid spread of *Marquis* was not an isolated case. Around 1900, Mark Alfred Carleton, the USDA's most prominent wheat breeder and plant explorer, introduced scores of cultivars from the Russian Empire. He explicitly was seeking cultivars that thrived in harsh environments. After proving drought and rust resistant in controlled tests, several durum introductions diffused rapidly in Minnesota and the Dakotas (13, 22).

Table 1, which lists a number of important new wheat cultivars, hints at the growing importance of public breeding (18, 23, 24). (The distinction between the public and private sectors is not precise, because breeders in both sectors use germplasm from the other.) In 1919, 27% of the wheat acreage in the United States was planted in cultivars introduced or bred by the public sector, and almost all Canadian wheat was a product of government breeders (ref. 19, pp. 208–218, and ref. 21, pp. 73–74). Public breeding activities in the United States increased during much of the 20th century. In 2008 varietal surveys of United States and Canadian wheat (*SI Text*) indicate that roughly 84% of wheat acreage is planted to cultivars bred by the public sector. Improvements in breeding technologies along with environmental and economic factors have accelerated the rate of turnover of wheat cultivars. The surveys show that the median vintage of modern wheats is 12 y. By comparison, the 1919 median vintage was about 3 decades. Changes in technologies have allowed farmers to adopt seeds tailored to narrower geoclimatic niches; none of the more than 300 cultivars reported in the 2008 surveys accounted for more than 6% of total United States and Canadian wheat acreage. Breeding advances in the United States and Mexico accelerated the evolution of wheat germplasm and the plant's adaptability to more varied conditions.

Home of the Green Revolution. Although wheat appeared early in Mexico, the crop always was secondary to maize. By the early 1940s, rust diseases were taking a heavy toll, and national production accounted for less than one-half of consumption. In 1943, the Mexican government together with the Rockefeller Foundation established a wheat-breeding program, hiring Norman Borlaug as the chief breeder the following year. His invention of shuttle breeding (whereby researchers took seed grown at high elevations over the summer to grow at low elevations over the winter and then planted the resulting seed at high elevations the following spring, and then repeated the process) allowed Borlaug to cut the breeding time in half and develop plants adapted to a wider range of environmental conditions. Borlaug bred a series of early-ripening, fertility-responsive, rust-resistant cultivars, but

sundry provincial sources to fill gaps in the Census data between 1950 and 1976. To link the output data for the period before 1978, we combine the Canadian 1850 production with the United States production for 1849, and so forth. The number of local entities varies over time; in 1929, for example, we record data on 3,070 counties in the United States and 216 units in Canada. We have wheat production data for every 10 y from 1839–1978 and for every 5 y thereafter. We base our 1839 Canadian output estimate on data for Upper Canada (Ontario) in 1842 and Lower Canada (Quebec) in 1844.

For the United States, we linked production to each county's location based on its 1970 population centroid from ref. 41. For Canada the data are less standardized, and the local units reported by official sources include counties, census divisions, and agricultural districts depending on the province and year. We linked production to a fixed location with each local unit.

Climate Norm Data. The geoclimatic variables reflect average 1941–1970 conditions in each county or agricultural district as recorded by the US National Oceanic and Atmospheric Administration in ref. 41 or the Canadian Atmospheric Environment Service (42). These climate norms largely predate the more recent climate changes associated with the global warming.

Mexican Data. We do not include Mexico in our primary long-run analysis because of a paucity of data until 1929 and Mexico's relatively low wheat output for most of the period. Production data for 1929 are from ref. 43, those for 1969 are from ref. 44, and those for 2007 are from Sistema de

Información Agroalimentaria de Consulta, www.siap.gob.mx. Weather norms are from ref. 45.

Baseline Climate Data. The baseline (1981–1990) annual temperatures and precipitation for Columbus, OH, Dickerson, ND, Edmonton, AB, Fort Dodge, KS, and Ciudad Obregón are from World Weather Records, <http://dss.ucar.edu/datasets/ds570.1/>.

Regression Analysis. Fig. 2 connects the latitude at which spring wheat output equaled winter wheat output for each longitude between 87° W and 105° W. The points are derived from regressions of the proportion of spring wheat in total county wheat production in the 1869 and 1929 for United States and Canada; observations are binned by degree of longitude.

ACKNOWLEDGMENTS. We thank participants at the National Bureau of Economic Research Climate Change Conferences, the 2010 World Economic History Congress, 2010 Allied Social Science Associations meetings, the 2010 Western Economic Association International Conference, and colloquia at Oxford, Copenhagen, University of Guelph, Duke University, University of California, Berkeley, University of California, Davis, and Can Tho University. Julian Alston, P. Stephen Baenziger, Brady Deaton, Ron DePauw, Helen Goldstein, Jeffrey Graham, Michael Haines, Brian Hubner, Kris Inwood, Josh MacFayden, Philip Pardey, Carol Perry, Calvin Qualset, Kathleen Stroud, and Chuanliang Su provided data and/or comments. The research was supported by National Science Foundation Grants SES-0550913, SES-0551130, and SES-0921732, and by the University of California Agricultural Issues Center and the Institute of Governmental Affairs, University of California, Davis.

- Christensen JH, et al. (2007) *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, eds Solomon S, et al. (Cambridge Univ Press, Cambridge, UK), pp 889–890.
- Sokolov AP, et al. (2009) Probabilistic forecast for 21st century climate based on uncertainties in emissions (without policy) and climate parameters. *J Clim* 22: 5175–5204.
- Rosenzweig C, Parry ML (1994) Potential impact of climate change on world food supply. *Nature* 367:133–138.
- Mendelsohn R, Nordhaus WD, Shaw D (1994) The impact of global warming on agriculture: A Ricardian analysis. *Am Econ Rev* 84:371–411.
- Cline W (2007) *Global Warming and Agriculture: Impact Estimates by Country* (Center for Global Development, Washington, DC), pp 7–21.
- Nelson GC, et al. (2009) *Climate Change: Impact on Agriculture and Costs of Adaptation* (International Food Policy Research Institute, Washington, DC), pp 7–21.
- US Climate Change Science Program (2008) *The Effects of Climate Change on Agriculture. Land Resources, Water Resources, and Biodiversity in the United States*, (Climate Change Science Program, Washington, DC), pp 71–78.
- Howden SM, et al. (2007) Adapting agriculture to climate change. *Proc Natl Acad Sci USA* 104:19691–19696.
- Ortiz R, et al. (2008) Climate change: Can wheat beat the heat? *Agric Ecosyst Environ* 126:46–58.
- Del Río Moreno J, López y Sebastián L (1996) El trigo en las ciudad de México. *Revista Complutense de Historia de América* 22:33–51.
- White WJ (1995) *Harvest of Gold: The History of Field Crop Breeding in Canada*, eds Slinkard AE, Knott DR (University Extension Press, Univ of Saskatchewan, Saskatoon, SK, Canada), p 6.
- Pritchett J (1942) *The Red River Valley, 1811–1849: A Regional Study* (Yale Univ Press, New Haven, CT), pp 113–228.
- Olmstead AL, Rhode PW (2008) *Creating Abundance: Biological Innovation and American Agricultural Development* (Cambridge Univ Press, New York).
- Libecap GD, Hansen ZK (2002) Rain follows the plow and dryfarming doctrine: The climate information problem and homestead failure in the upper Great Plains, 1890–1925. *J Econ Hist* 62:86–120.
- Carter SB, et al., eds (2006) *Historical Statistics of the United States* (Cambridge Univ Press, New York), Vol 4, p 209.
- Clark I (1958) *Then Came the Railroads* (Univ of Oklahoma Press, Norman, OK) 65–72, 84–90, 105–106, 205–206.
- Malin JC (1944) *Winter Wheat in the Golden Belt of Kansas* (Univ of Kansas Press, Lawrence, KS), pp 40–44, 96–101, 156–519, 162–187.
- Dalrymple DG (1988) Changes in wheat varieties and yields in the United States, 1919–1984. *Agric Hist* 62:20–36.
- Clark JA, Martin JH, Ball CR (1922) *Classification of American Wheat Varieties*. US Department of Agriculture Bulletin 1074. (US Department of Agriculture, Washington, DC) pp 90–91, 208–218.
- Salmon SC (1920) Developing better varieties of wheat for Kansas. *Wheat in Kansas* (Kansas State Board of Agriculture, Topeka, KS), pp 210–217.
- Pomeroy EM (1956) *William Saunders and His Five Sons: The Story of the Marquis Wheat Family* (Ryerson, Toronto, Canada), pp 48–52, 73–74.
- Paulsen GM, Shroyer JP (2008) Early history of wheat improvement in the Great Plains. *Celebrate the Centennial. Agron J* 100 (Suppl):S70–S78.
- Smith CW (1995) *Crop Production: Evolution, History, and Technology* (John Wiley, Hoboken, NJ), pp 66–74.
- McCallum BD, DePauw RM (2008) A review of wheat cultivars grown in the Canadian prairies. *Can J Plant Sci* 88:649–677.
- Borlaug NE (1954) Mexican wheat production and its role in the epidemiology of stem rust in North America. *Phytopathology* 44:398–404.
- Dalrymple DG (1986) *Development and Spread of Wheat Varieties in Developing Countries* (Agency for International Development, Washington, DC), pp 13–19.
- Rajaram S, Hettel GP, eds (1994) *Wheat Breeding at CIMMYT: Commemorating 50 Years of Research in Mexico for Global Wheat Improvement* (International Maize and Wheat Improvement Center, Ciudad Obregón, Mexico).
- Reynolds MP, Hayes D, Chapman (2010) Breeding for adaptation to heat and drought stress. *Climate Change and Crop Production*, ed Reynolds MP (CABI, Wallingford, UK), pp 71–91.
- Rosenzweig C, Hillel D (2008) *Climate Variability and the Global Harvest: Impacts of El Niño and Other Oscillations on Agroecosystems* (Oxford Univ Press, New York), pp 70–82.
- Rosenberg NJ (1982) The increasing CO₂ concentration in the atmosphere and its implications on agricultural productivity II. Effects through CO₂-induced climatic change. *Clim Change* 4:139–154.
- Covert JR (1912) *Seedtime and Harvest: Cereals, Flax, Cotton, and Tobacco*. Bureau of Statistics Bulletin 85 (US Department of Agriculture, Washington, DC).
- US Department of Agriculture (1997) *Usual Planting and Harvesting Dates for U.S. Field Crops*. Agricultural Handbook 628 US Department of Agriculture.
- Kamiguchi K, Kitoh A, Uchiyama T, Mizuta R, Noda A (2006) Changes in precipitation-based extremes indices due to global warming projected by a global 20-km-mesh atmospheric model. *SOLA* 2:64–67.
- Uchiyama T, Mizuta R, Kamiguchi K, Kitoh A, Noda A (2006) Changes in temperature-based extremes indices due to global warming projected by a global 20-km-mesh atmospheric model. *SOLA* 2:68–71.
- Alexander WH (1923) *A Climatological History of Ohio*. Ohio State University Engineering Experiment Station Bulletin 26 (Columbus, OH), pp 112, 176–179, 308–315, 594–597, 623–626, 665–670.
- Alston JM, Beddow JM, Pardey PG (2009) Agriculture. Agricultural research, productivity, and food prices in the long run. *Science* 325:1209–1210.
- Inter-University Consortium for Political and Social Research (2005) *Historical Demographic, Economic, and Social Data, 1790–2002* ICPSR 2896.
- US Census Bureau (1942) *Sixteenth Census: Agriculture* (GPO, Washington, DC).
- Canada Census Office (1842–1951) *Agricultural Census* (Ottawa, Ontario).
- Statistics Canada (2009) *Crops Small Area Data* (Statistics Canada, Ottawa, Ontario, Canada).
- Inter-University Consortium for Political and Social Research (1991) *Bureau of Health Professions Resource File* ICPSR 9075.
- Canada Atmospheric Environment Service (1972) *Temperature and precipitation, 1941–1970*. 6 vols (Department of the Environment, Downsview, Ontario, Canada).
- Banco Nacional de Crédito Agrícola (1938) *El Trigo en México, Parte I, Análisis Estadístico de la Producción* (BNCA, México, D.F.), pp 30–34.
- México, Dirección General de Estadística (1975) *Quinto Censos Agrícola-Ganadero Y Ejidal, 1970 Resumen General* (Dirección General de Estadística, México, D.F.).
- México, Servicio Meteorológico Nacional (1976) *Normales Climatológicas, Período 1941–1970* (Secretaría de Agricultura y Ganadería, México, D.F.).