Published in final edited form as:

J Aging Health. 2011 February ; 23(1): 86–111. doi:10.1177/0898264310383156.

Older Adults' Inpatient and Emergency Department Utilization For Ambulatory-Care-Sensitive Conditions: Relationship with Alcohol Consumption

Elizabeth S. Levy Merrick, Ph.D., MSW¹, Dominic Hodgkin, Ph.D.¹, Deborah W. Garnick, Sc.D.¹, Constance M. Horgan, Sc.D.¹, Lee Panas, MS¹, Marian Ryan, RRT, MPH, CHES¹, Frederic C. Blow, Ph.D.², and Richard Saitz, M.D., M.P.H.³

Dominic Hodgkin: Hodgkin@brandeis.edu; Deborah W. Garnick: garnick@brandeis.edu; Constance M. Horgan: horgan@brandeis.edu; Lee Panas: lpnas@brandeis.edu; Marian Ryan: mryan@brandeis.edu; Frederic C. Blow: fredblow@umich.edu; Richard Saitz: Richard.Saitz@bmc.org

¹ Institute for Behavioral Health, Heller School for Social Policy and Management, Brandeis University. Fax 781-736-3985. Dominic Hodgkin, Ph.D, Phone 781-736-8551; Deborah W. Garnick, Sc.D, Phone 781-736-3840; Constance M. Horgan, Sc.D., Phone 781-736-3916; Lee Panas, MS, Phone 781-736-3943; Marian Ryan, RRT, MPH, CHES, Phone 781-736- 8493

² Department of Psychiatry, University of Michigan, Department of Veterans Affairs, Health Services Research and Development, Serious Mental Illness Treatment Research and Evaluation Center. Frederic C. Blow, Ph.D., Phone 734-845-5046

³ Boston University School of Medicine, Clinical Addiction Research and Education (CARE) Unit, Section of General Internal Medicine, Boston Medical Center, Youth Alcohol Prevention Center & Department of Epidemiology, Boston University School of Public Health. Richard Saitz, M.D., M.P.H, Phone 617-414-774, Fax 617-414-4676

Abstract

Objectives—This study examined the relationship between drinking that exceeds guideline-recommended limits and acute-care utilization for ambulatory-care-sensitive conditions (ACSCs) by older Medicare beneficiaries.

Methods—This secondary data analysis utilized the 2001–2006 Medicare Current Beneficiary Survey (unweighted n=5,570 community dwelling, past-year drinkers, 65 years and older). Self-reported alcohol consumption (categorized as within-guidelines, exceeding monthly but not daily limits, or heavy episodic) and covariates were used to predict ACSC hospitalization, emergency department visit not resulting in admission, and emergency department visit that did result in admission.

Results—Heavy episodic drinking was significantly associated with higher likelihood of an ACSC emergency department visit not resulting in admission (adjusted odds ratio 1.91, 95% CI 1.11 – 3.30; p<.05). Drinking pattern was not significant for other ACSC measures.

Discussion—Results partially support the hypothesis that excessive drinking may be related to ACSC acute-care utilization among older adults, suggesting increased risk of lower-quality outpatient care.

Keywords

older adults; alcohol; ambulatory-care-sensitive conditions; health care utilization; quality of care

Introduction

Older adults frequently have unhealthy drinking patterns, ranging from risky drinking (which incurs increased risk of adverse consequences) to alcohol disorders (Saitz, 2003). Recent national prevalence estimates for adults aged 65 and older range from 9% drinking in excess of national guidelines (Merrick et al., 2008a) to gender-specific rates of 13% and 8% with at-risk use and 14% and 3% with binge drinking (five or more drinks on same occasion in past 30 days) for men and women respectively (Blazer & Wu, 2009). Among the problems associated with excessive drinking is the possibility of increased risk for inadequate medical care. This study investigated the relationship between excessive drinking and acute-care utilization for ambulatory-care-sensitive conditions (ACSCs).

Excessive drinking could be connected to inadequate medical care in several ways. Patients with excessive drinking may under-utilize routine care including primary care (Ford, Trestman, Tennen, & Allen, 2005; Girard, Partridge, Becker, & Bock, 2004; Kunz, 1997; Rice & Duncan, 1995; Rice et al., 2000), for reasons including concerns about stigma or financial barriers to care. Excessive alcohol consumption could reflect generalized self-neglect of health (Blow, Brockmann, & Barry, 2004; Hazelton, Sterns, & Chisholm, 2003). In any case, the result may be lower use of recommended services or delays in seeking care until problems are more severe. At the same time, providers sometimes have attitudinal barriers towards patients with alcohol problems (Anderson et al., 2004; Deehan, Templeton, Taylor, Drummond, & Strang, 1998; Freidman, McCullough, Chin, & Saitz, 2000; Kaner et al., 2009) which could potentially result in differential treatment. These patients may be more difficult to treat, or scarce clinical time may be taken up with competing demands.

While the mechanisms are not well understood, there is some evidence of lower-quality medical care for persons with mental disorders, and in some cases substance use disorders specifically, in disparate clinical contexts (Clark, Weir, Ouellette, Zhang, & Baxter, 2009; Desai, Rosenheck, Druss, & Perlin, 2002; Druss, Rask, & Katon, 2008; Rathore, Wang, Druss, Masoudi, & Krumholz, 2008). Older adults who drink excessively according to various criteria are less likely to receive preventive medical care (Merrick et al., 2008b; Moore et al., 2001; Ozminkowski et al., 2006).

However, acute-care utilization for ACSC has seldom been explored in relation to excessive drinking. ACSCs are conditions for which “the provision of timely and effective outpatient care can help to reduce the risks of hospitalization by either preventing the onset of an illness or condition, controlling an acute episodic illness or condition, or managing a chronic disease or condition” (Billings et al., 1993). Congestive heart failure, diabetes, asthma and chronic obstructive pulmonary disease are examples of ACSCs. Measuring hospitalization for ACSCs is one approach to examining access to adequate outpatient care. Numerous ACSC indicators have been developed, including the, widely used Agency for Healthcare Research and Quality (AHRQ) Prevention Quality Indicators. These are designed to identify admissions that evidence suggests could have been avoided, at least in part, through better access to high-quality outpatient care (Agency for Healthcare Research and Quality, 2008). Researchers have considered which ACSC indicators are most appropriate for an older adult population (McCall, 2004). The concept and diagnosis specifications have more recently been extended to emergency department visits (Logan, Riley, & Barker, 2008; McCall, 2004).

The relationship between alcohol use and ACSC inpatient and emergency department utilization is important to investigate in older adults, who experience adverse health events at high rates and face special issues with alcohol. The National Institute on Alcohol Abuse and Alcoholism (NIAAA) and American Geriatrics Society define risky drinking amounts for those 65 years and older as more than seven drinks per week, or more than 3 drinks on any single day (American Geriatrics Society Clinical Practice Committee, ; NIAAA). Exceeding these limits is associated with interpersonal and functioning problems for elders (Moos, Brennan, Schutte, & Moos, 2004) who have higher sensitivity and impaired ability to metabolize alcohol (Saitz, 2003). Limits may be lower or abstinence may be best for older adults who use medications that interact with alcohol or have medical problems that can be adversely affected by alcohol consumption.

Many studies have examined the occurrence and predictors of ACSC hospitalization (and, more recently, ACSC emergency department visits) but few have examined the role of alcohol consumption. Li and colleagues examined New York state hospital discharge data for adults aged 20–64 years (Li, Glance, Cai, & Mukamel, 2008). Inpatients with mental disorders (including substance use disorders) were significantly more likely than other admitted inpatients to have been admitted for an ACSC and also had longer average length of stay. Another study with a mixed-age inpatient sample found that binge drinking (five or more alcoholic beverages per occasion during the past 30 days) was predictive of preventable hospitalization (Arozullah et al., 2006).

Other factors found to be associated with ACSC hospitalization include health status, comorbidity and functional limitations (Culler, Parchman, & Przybylski, 1998; Niefeld et al., 2003), income (Billings, Anderson, & Newman, 1996; Billings et al., 1993; Blustein, Hanson, & Shea, 1998), health insurance status (Weissman, Gatsonis, & Epstein, 1992; Zeng et al., 2006), mental disorders (Bynum et al., 2004; Himelhoch, Weller, Wu, Anderson, & Cooper, 2004), access to regular primary care or community health centers (Culler et al., 1998; Epstein, 2001; Falik, Needleman, Wells, & Korb, 2001), continuity of care (Gill & Mainous, 1998), and race and ethnicity (Culler et al., 1998; Friedman & Basu, 2004; Laditka, Laditka, & Mastanduno, 2003). McCall et al., using the 1999 Medicare Current Beneficiary Survey, found that prior ACSC hospitalization, comorbidity, and health status were among the significant predictors for some ACSCs among older beneficiaries (McCall, 2004). No variables for alcohol use or mental disorders were included. For ACSC emergency department use, barriers to primary care may contribute to higher utilization for ACSCs among black adults and Medicaid patients (Oster & Bindman, 2003). Among older adults, emergency department and observation stay utilization increased between 1992 and 2000 for ten of eleven ACSCs studied (including congestive heart failure, pneumonia and cellulitis) (McCall, 2004).

The current study aimed to help address the lack of research in this area by focusing on the relationship between self-reported alcohol consumption and ACSC acute-care utilization in a nationally representative sample of older adults. We hypothesized that drinking that exceeded guideline-recommended limits, particularly heavy episodic drinking, would be positively related to ACSC hospitalization and emergency department use.

Method

Data and Sample

The primary data source was the Medicare Current Beneficiary Survey (MCBS) for 2001–2006. The MCBS is an ongoing survey of a representative national sample of the Medicare population by the Centers for Medicare and Medicaid Services. The sample is selected using a stratified, multistage probability design to represent the national Medicare population

(Centers for Medicare and Medicaid Services). Sample weights are provided to achieve nationally representative estimates, in this analysis for the continuously enrolled Medicare population. Beneficiaries sampled from Medicare enrollment files (or proxies) are interviewed three times a year including in-person, computer-assisted interviewing. There is a 4-year rotating panel design. Beneficiaries were randomly selected according to age strata from a nationally representative set of 107 geographic primary sampling units, with oversampling of the disabled (age <65) and those aged 65 and older. Normalized sampling weights were assigned to represent the population. The survey content includes sociodemographics, health and functional status, and utilization. The 2001, 2003 and 2005 MCBS included items regarding alcohol consumption as well as data for covariates (detailed in “Measures”) (Centers for Medicare and Medicaid Services). Subjects’ Medicare claims were linked to survey data for this analysis. We used data from 2001, 2003 and 2005 for baseline characteristics including drinking and covariates, and from 2002, 2004, and 2006 to identify hospitalization and emergency department services for ACSCs. This approach seeks to avoid reverse causality bias between ACSC use and other variables. To maximize sample size, we included beneficiaries present in the MCBS and continuously enrolled for any two consecutive years: 2001 and 2002; 2003 and 2004; or 2005 and 2006.

The analytic sample identification started with 20,482 community-dwelling beneficiaries who were 65 years or older. Of these, 13,612 were non-drinkers and another 129 were missing alcohol data, and were excluded. This analysis focused on persons who reported drinking alcohol in a typical month in the past year, since non-drinkers would constitute an especially heterogeneous group including lifetime abstainers and those who quit due to health problems, and these differences would be unobservable. Health maintenance organization enrollees were excluded because their claims were not available (N=1,171). Our final study sample included 5,570 persons representing a weighted N of 15,128,450. Bivariate comparison totals varied due to item-missing data (all <4%). The logistic regression sample consisted of 5,046 individuals.

Measures

Dependent Variables: Measures of Acute-Care Utilization for ACSCs—The three dependent variables were binary measures of any ACSC hospitalization, ACSC emergency department visit not resulting in inpatient admission (thus not overlapping with ACSC hospitalization), and ACSC emergency department visit that did result in admission. We identified ACSCs by applying the specifications from two sources. First, following AHRQ specifications, we used all 13 relevant PQIs (excluding low birth weight): diabetes short-term complications, perforated appendix, diabetes long-term complications, chronic obstructive pulmonary disease (COPD), hypertension, congestive heart failure (CHF), dehydration, bacterial pneumonia, urinary tract infection, angina without procedure, uncontrolled diabetes, adult asthma, and rate of lower extremity amputation among patients with diabetes (Agency for Healthcare Research and Quality, 2008). Second, we adopted 15 indicators identified by McCall et al. through literature review and deemed appropriate by clinical experts for application to older adults: asthma/COPD, cellulitis, CHF, dehydration, diabetes, hypertension, hypoglycemia, hypokalemia, influenza, urinary tract infection, malnutrition, perforated or bleeding ulcer, pneumonia, seizure disorder, severe ear/nose/throat infection (McCall, 2004). The specifications were adopted from the original source and an expanded version published in a later ACSC study (Zeng et al., 2006).

There is substantial overlap between the two indicator lists. If a utilization event qualified in terms of either source, we counted it as an ACSC event. For sensitivity analysis purposes, we examined frequencies for each list of indicators separately, and found that the differences were quite small in magnitude. We identified hospitalizations with an ACSC principal

diagnosis. While the PQI set was developed for application to inpatient hospitalization, we follow previous research that has extended the ACSC concept and diagnostic specifications to emergency department visits (Logan et al., 2008; McCall, 2004).

We identified emergency department visits based on procedure and location codes. For emergency department utilization that did not result in hospital admission, we used the visit primary diagnosis. Emergency department visits that resulted in admission were necessarily identified through the inpatient file, and the primary admitting diagnosis was attributed to the preceding emergency department visit.

Alcohol Consumption Variables—The 2001, 2003 and 2005 MCBS included three alcohol consumption items. Quantity and frequency were ascertained by asking, “Please think about a typical month in the past year. On how many days did you drink any type of alcoholic beverage? On those days that you drank alcohol, how many drinks did you have?” Heavy episodic drinking was assessed by asking, “Please think about a typical month in the past year. On how many days did you have 4 or more drinks in a single day?” Alcoholic beverages were described as including “liquor such as whiskey or gin, mixed drinks, wine, beer, and any other type of alcoholic beverage.”

To assess unhealthy drinking in terms of consuming risky amounts of alcohol (regardless of whether alcohol consequences or disorders were present), we defined alcohol measures reflecting two parameters in the NIAAA and American Geriatrics Society guidelines (American Geriatrics Society Clinical Practice Committee). First, to be consistent with the weekly guideline we defined “exceeding monthly limits” as more than 30 drinks per typical month. (Forty-two respondents reporting 31 drinks per month whose responses were clearly based on a 31-day month were coded as negative, since the items did not specify standardized number of days per month.) Second, we constructed a “heavy episodic drinking” variable indicating whether an individual reported four or more drinks in any single day during a typical month in the past year, according to either drinking quantity item.

We categorized respondents into three mutually exclusive categories: within-guidelines drinkers (not exceeding the monthly limit or the three-drink single-day limit); drinkers who exceeded the monthly limit but not the single-day limit; and heavy episodic drinkers who exceeded the single-day drinking limit, with or without exceeding the monthly limit. For secondary analyses, we calculated a continuous measure of drinks per month, based on the quantity-frequency responses. To address nonlinearity, we included a squared term in the regressions.

Covariates—Covariates were selected that previous research found to affect health care utilization.

Sociodemographic variables: We included gender, race, Hispanic ethnicity, annual household income, age, education, marital status, region, and residence in a metropolitan area. Living arrangement was not included due to high correlation with marital status.

Health status variables: We controlled for health status by utilizing DxCG (diagnostic cost group) risk adjustment software that uses sex, age and diagnosis codes from claims to construct a continuous measure of relative risk of healthcare resource use (DxCG, 2009). Compared to other illness burden indices or scales, the DxCG score contains higher specificity related to the individual’s clinical profile in projecting future health care costs and estimating an individual’s care management needs (Zhao, Ash, Ellis, & Slaughter, 2002; Zhao et al., 2001). Thus, it may be used as a proxy for health status in that higher DxCG risk scores denote higher health care resource use risk and presumably poorer health (Wang et

al., 2000). A value of 1 indicates the individual's predicted cost equals the population average for all persons with Medicare claims; higher values indicate higher-than-average predicted costs. For bivariate analyses, we created categories: no claims or claims not indicative of significant health risk (DxCG score < 0.1); claims indicative of lower-than-average health risk ($0.1 \leq \text{DxCG score} \leq 1$), claims indicative of higher-than-average health risk (DxCG score > 1). For logistic regression models, we used the continuous measure; increasing scores indicate higher risk of health care resource use (poorer health status). We also included dichotomous variables for current smoking, and for presence of a chronic disease explicitly related to ACSC indicators. We used diagnosis codes from the Centers for Medicare and Medicaid Services' Chronic Condition Data Warehouse (COPD, diabetes, CHF) (Buccaneer Computer Systems and Services, 2009) and the AHRQ PQI specifications (hypertension, asthma); two outpatient claims or one inpatient claim during the baseline year were required for all but CHF in which one claim of any type was required.

We controlled for functional status utilizing a modified Katz Index of Independence in Activities of Daily Living (ADL) variable constructed from survey data (Katz, Ford, Moskowitz, Jackson, & Jaffe, 1963; Shelkey & Wallace, 1999). Respondents were asked whether they had trouble or needed assistance with six ADLs: bathing, dressing, transferring, toileting, continence or feeding. If no difficulty was indicated, that activity received a score of 1. The resulting variable reflects a 7-position scale (0–6) of the number of independent ADLs.

Two mental health variables were used. First, a self-reported depression variable was created. Respondents were asked, "In the past 12 months, how much of the time did you feel sad, blue, or depressed?" (all, most, some, little, or none of the time), and "In the past 12 months, have you had 2 weeks or more when you lost interest or pleasure in things that you usually cared about or enjoyed?" (yes, no). Respondents who answered "all" or "most of the time" to the first question, and/or "yes" to the second question were categorized as having self-reported depression. This approximates the modified PHQ-2 validated for older adults (Li, Friedman, Conwell, & Fiscella, 2007). Second, a dichotomous variable indicating behavioral health diagnosis was constructed based on the presence of one inpatient claim or two outpatient claims with an ICD-9 mental health or substance use disorder diagnosis, excluding alcohol disorders. We excluded alcohol diagnosis from our main models as we wanted to observe the full effect of self-reported drinking, but we included this in alternate versions. We found that although in bivariate analyses the presence of a claim with an alcohol disorder diagnosis was associated with significantly higher rates of ACSC acute-care utilization, it was not statistically significant in a multivariate context.

Access variables: We included measures on self-reported trouble getting needed care during the past 12 months, private supplemental insurance, and Medicaid coverage. We also included a measure for living in a county designated as a full-county primary care health professional shortage area according to Health Resources and Services Administration data (Health Resources and Services Administration). In addition, we constructed variables indicating type of usual care: no particular "medical person or clinic" usually seen when respondent is sick or for advice, except for hospital emergency room or walk-in urgent care center; primary care physician (family practice, general practice, geriatrics, internal medicine); or non-primary care physician. We constructed a variable indicating a prior-year ACSC event. Due to multicollinearity it was not included in final models. When we tested its inclusion, this variable was positive and significant, but did not affect the direction, magnitude or significance of alcohol variables.

Statistical Analysis

Results are weighted estimates that represent the continuously enrolled, community-dwelling, non-HMO, elderly Medicare population of current drinkers. Chi-square tests were used to assess bivariate differences; chi-square statistics were corrected for the survey design and converted to F-statistics. The continuous drinks per month variable was significantly skewed and bivariate testing accounted for skewness. We conducted logistic regression analyses to model occurrence of an ACSC hospitalization or emergency room visit as a function of alcohol consumption (drinking category) and covariates. We also constructed a continuous measure of alcohol consumption—number of drinks per month—and included in a separate set of logistic regression models. We explored the use of an instrumental variables approach to address any potential bias due to unobserved differences between drinking groups. Among several exogenous variables (e.g., Sunday ban on alcohol sales, beer tax) that have been previously used as instruments for alcohol use, none were found for this sample of older drinkers that met the assumptions required for instrumental variables analysis (Wooldridge, 2006) and were robust to specific geographic inclusion (notably Puerto Rico).

Due to the complex sampling design, using procedures which assumed equal probability of selection would likely lead to underestimating standard errors (Cohen, 1997; Lemeshow et al., 1998). The SVY:LOGIT procedure of the statistical package STATA version 9.0 was used to more accurately determine the statistical significance of observed differences (STATA Corporation).

Results

As shown in Table 1, the weighted sample was predominantly white (93.0%), male (55.2%), and married (65.4%). Only 16.2% were over 80 years old. More than half had some education beyond high school. Three-quarters (75.2%) reported within-guidelines alcohol consumption in a typical month in the past year. Ten percent exceeded the monthly limit only, and 14.8% reported heavy episodic drinking. Drinking patterns varied significantly by sociodemographic, health and access variables.

Overall, 3.5% had an ACSC hospitalization during the year, 2.1% had an ACSC emergency department visit that did not result in inpatient admission, and 2.1% had an emergency department visit that resulted in admission (Table 2). Drinking category was not significantly related to ACSC hospitalization in bivariate analyses. Drinking category was significantly associated with ACSC emergency department visits not resulting in admission (1.9% for within-guidelines drinkers, 2.8% for those exceeding monthly limits only, and 3.1% for heavy episodic drinkers, $p < .05$). Drinking category was not significantly related to ACSC emergency department visits that did result in inpatient admission. The continuous measure of drinks per month was not significantly associated with ACSC measures in bivariate tests.

The most common ACSCs for hospitalization were congestive heart failure (0.9%), bacterial pneumonia (0.8%), and asthma/chronic obstructive pulmonary disease (0.7%; data not shown). For emergency department visits that did not result in admission, urinary tract infections (0.4%), asthma/chronic obstructive pulmonary disease (0.3%), and cellulitis (0.3%) were most common. For emergency department visits that resulted in admission, asthma/chronic obstructive pulmonary disease (0.4%), congestive heart failure (0.4%), and bacterial pneumonia (0.4%) were most common. While the numbers within each specific ACSC were small, thus limiting statistical power to identify significant differences by drinking pattern, there was a significant bivariate relationship between specific ACSC and

drinking pattern in several cases. For example, there was a significant relationship between hypertension and drinking pattern across all types of utilization ($p < .01$) with heavy episodic drinkers having the highest utilization. Urinary tract infection and cellulitis were also significantly associated with drinking pattern ($p < .01$), with persons who exceeded monthly guidelines only showing the highest rates of emergency department visits that did not result in admission.

In the logistic regression model predicting ACSC hospitalization, drinking variables were not significant. Being over 80 years of age, African-American, having higher relative health risk score, independence in 0 to 4 ADLs, and chronic disease were associated with higher likelihood of an ACSC hospitalization; living in the West region of the country was associated with lower likelihood relative to the South. In the model predicting any ACSC emergency department visit that did not result in inpatient admission, heavy episodic drinking was significantly associated with higher likelihood (adjusted odds ratio 1.91, 95% confidence interval 1.11 – 3.30, $p < .05$). Other factors associated with greater likelihood of utilization included older age, greater relative health risk, independence in 0 to 4 ADLs, and chronic disease. Finally, in the model predicting any ACSC emergency department visit that resulted in inpatient admission, drinking variables were not significant. Greater relative health risk and independence in 0–4 ADLs were associated with higher likelihood of this type of utilization. Being a high school graduate (relative to having less than a high school education), annual household income of over \$40,000 per year, and living in the Midwest or West (relative to South) were associated with lower likelihoods.

In another set of alternate logistic regression analyses using a continuous measure of drinks per month as well as a squared term to address non-linearity, there was no significant effect on ACSC hospitalization or emergency department utilization resulting in admission (data not shown). However, for emergency department utilization that did not result in admission, there was a significant, positive relationship (OR 1.01, $p < .05$)

Discussion

Heavy episodic drinking was a predictor of ACSC emergency department use that did not result in admission, but drinking variables did not predict ACSC hospitalization or emergency department visits resulting in hospital admission. Thus, we found partial support for the hypothesis that older adults whose drinking exceeded guideline-recommended limits would be at greater risk of ACSC acute-care utilization. This is consistent with prior studies that found evidence for lower quality of medical or preventive care for persons with substance abuse problems defined in various ways (Arozullah et al., 2006; Clark et al., 2009; Desai et al., 2002; Li et al., 2008; Merrick et al., 2008; Ozminkowski et al., 2006), and extends this line of inquiry by finding this connection between older adults' self-reported drinking and ACSC emergency department use. Since ACSC severity level for emergency department visits not resulting in admission is likely lower, our findings suggest that people with heavy episodic drinking may experience some deficiencies in access to or quality of outpatient care but in ways that do not affect the most severe outcomes. Heavy episodic drinkers may be using the emergency department for issues that should normally have been addressed earlier in primary care but are not so far advanced as to require hospitalization. Small numbers greatly inhibited our ability to examine specific ACSCs. However, we did find that for some ACSCs (notably hypertension) that can be worsened by excessive drinking, utilization varied by drinking pattern with heavy episodic drinkers more likely to have ACSC utilization.

The fact that this relationship occurs only for those with heavy episodic drinking rather than those who exceed monthly limits only echoes findings from a previous study on receipt of

preventive services (Merrick et al., 2008b). The level of drinking represented in the group exceeding monthly limits only is not extremely high, with most respondents indicating consumption of two drinks per day most days of the month. Older adults whose drinking exceeds monthly limits only may reflect a population without current impairment affecting use of outpatient services that prevents ACSC acute-care utilization. Heavy episodic drinkers may be more likely to have alcohol disorders, compared to persons exceeding monthly guidelines only, some of whom who may be continuing to drink at levels that were acceptable for younger ages.

The alternate, exploratory analysis we conducted using drinks per month as the key explanatory variable, when we accounted for non-linearity, yielded results similar to the results for heavy episodic drinking. Future research to investigate the potentially complex influences of drinking pattern as well as drinking quantity in more detail would be fruitful. Future work should also investigate the causal mechanisms underlying the association we identified between episodic heavy drinking (or greater number of drinks per month) and ACSC emergency department visits not resulting in admission. Identifying the provider, patient and system roles in barriers to care for this group will be useful.

Study limitations include possible underreporting of alcohol consumption, although self-reported alcohol consumption is in general considered to be as accurate as other drinking measures (Babor, Steinberg, Anton, & Del Boca, 2000). Relatively small sample size limited statistical power to detect small differences. As noted earlier, it is possible that unobserved differences between drinking groups could create bias, but we were unable to identify satisfactory instruments for our sample and measures. However, one main possible source of bias in this type of analysis is health status differences, and the multivariate models we used included multiple health and functional status variables, somewhat reducing risk this bias. Furthermore, we focused on past-year drinkers, avoiding the bias which could have resulted from including a large, heterogeneous group of non-drinkers, many of whom might have quit drinking due to ill health. Previous research on effects of drinking on acute-care utilization (but not ACSC specifically) that used instrumental variables found no evidence of bias in a different nationally representative sample of older adults (Balsa, Homer, Fleming, & French, 2008). Another limitation is the imprecision of the continuous drinking measures for heavy episodic drinkers in our secondary analyses. This is due to lack of harmonization across alcohol variables and lack of information on specific number of drinks consumed on heavy drinking days. We also note that the application of ACSC indicators to emergency department utilization is a more recent extension of the concept and specifications originally applied to hospitalization.

The study findings add to the understanding of the full range of risks associated with drinking that exceeds recommended guidelines for older adults. They may be useful to outpatient and emergency room providers in raising awareness of this issue, which in turn may heightened vigilance and effective interventions. This may contribute to efforts to both address alcohol issues and reduce acute-care utilization for ACSCs in older adults. The findings also provide a useful contribution in analyzing the effects of a relevant factor largely absent from the large body of research on predictors of ACSC hospitalization and emergency department utilization.

References

- Agency for Healthcare Research and Quality. Prevention Quality Indicators: Technical Specifications. 2008. D. o. H. a. H. Services o. Document Number

- American Geriatrics Society Clinical Practice Committee. Clinical guidelines for alcohol use disorders in older adults. Retrieved <http://www.americangeriatrics.org/products/positionpapers/alcohol.shtml>, from <http://www.americangeriatrics.org/products/positionpapers/alcohol.shtml>
- Anderson P, Kaner E, Wutzke S, Funk M, Heather N, Wensing M, et al. Attitudes and managing alcohol problems in general practice: an interaction analysis based on findings from a WHO collaborative study. *Alcohol* 2004;39(4):351–356.
- Arozullah AM, Lee SY, Khan T, Kurup S, Ryan J, Bonner M, et al. The roles of low literacy and social support in predicting the preventability of hospital admission. *J Gen Intern Med* 2006;21(2):140–145. [PubMed: 16336616]
- Babor TF, Steinberg K, Anton R, Del Boca F. Talk is cheap: measuring drinking outcomes in clinical trials. *J Stud Alcohol* 2000;61(1):55–63. [PubMed: 10627097]
- Balsa AI, Homer JF, Fleming MF, French MT. Alcohol consumption and health among elders. *Gerontologist* 2008;48(5):622–636. [PubMed: 18981279]
- Billings J, Anderson GM, Newman LS. Recent findings on preventable hospitalizations. *Health Aff (Millwood)* 1996;15(3):239–249. [PubMed: 8854530]
- Billings J, Zeitel L, Lukomnik J, Carey TS, Blank AE, Newman L. Impact of socioeconomic status on hospital use in New York City. *Health Aff (Millwood)* 1993;12(1):162–173. [PubMed: 8509018]
- Blazer DG, Wu LT. The epidemiology of at-risk and binge drinking among middle-aged and elderly community adults: National Survey on Drug Use and Health. *Am J Psychiatry* 2009;166(10):1162–1169. [PubMed: 19687131]
- Blow FC, Brockmann LM, Barry KL. Role of alcohol in late-life suicide. *Alcohol Clin Exp Res* 2004;28(5 Suppl):48S–56S. [PubMed: 15166636]
- Blustein J, Hanson K, Shea S. Preventable hospitalizations and socioeconomic status. *Health Aff (Millwood)* 1998;17(2):177–189. [PubMed: 9558796]
- Buccaneer Computer Systems and Services. Chronic Condition Data Warehouse User Manual, version 1.5. Center for Medicare and Medicaid Services; 2009. Document Number
- Bynum JP, Rabins PV, Weller W, Niefeld M, Anderson GF, Wu AW. The relationship between a dementia diagnosis, chronic illness, medicare expenditures, and hospital use. *J Am Geriatr Soc* 2004;52(2):187–194. [PubMed: 14728626]
- Centers for Medicare and Medicaid Services. Medicare Current Beneficiary Survey. from <http://www.cms.gov/mcbs>
- Clark RE, Weir S, Ouellette RA, Zhang J, Baxter JD. Beyond health plans: behavioral health disorders and quality of diabetes and asthma care for Medicaid beneficiaries. *Med Care* 2009;47(5):545–552. [PubMed: 19319000]
- Cohen SB. An evaluation of alternative PC-based software packages developed for the analysis of complex survey data. *Am Stat* 1997;51(3):285–292.
- Culler SD, Parchman ML, Przybylski M. Factors related to potentially preventable hospitalizations among the elderly. *Med Care* 1998;36(6):804–817. [PubMed: 9630122]
- Deehan A, Templeton L, Taylor C, Drummond C, Strang J. Low detection rates, negative attitudes and the failure to meet the “Health of the Nation” alcohol targets: findings from a national survey of GPs in England and Wales. *Drug Alcohol Rev* 1998;17(3):249–258. [PubMed: 16203491]
- Desai MM, Rosenheck RA, Druss BG, Perlin JB. Mental disorders and quality of diabetes care in the veterans health administration. *Am J Psychiatry* 2002;159(9):1584–1590. [PubMed: 12202281]
- Druss BG, Rask K, Katon WJ. Major depression, depression treatment and quality of primary medical care. *Gen Hosp Psychiatry* 2008;30(1):20–25. [PubMed: 18164936]
- DxCG. Products and services, RiskSmart Stand Alone. 2009. Retrieved October 27, 2009, from <http://www.dxcg.com/products-services/risksmart-stand-alone.asp>
- Epstein AJ. The role of public clinics in preventable hospitalizations among vulnerable populations. *Health Serv Res* 2001;36(2):405–420. [PubMed: 11409820]
- Falik M, Needleman J, Wells BL, Korb J. Ambulatory care sensitive hospitalizations and emergency visits: experiences of Medicaid patients using federally qualified health centers. *Med Care* 2001;39(6):551–561. [PubMed: 11404640]

- Ford JD, Trestman RL, Tennen H, Allen S. Relationship of anxiety, depression and alcohol use disorders to persistent high utilization and potentially problematic under-utilization of primary medical care. *Soc Sci Med* 2005;61(7):1618–1625. [PubMed: 16005791]
- Friedmann PD, McCullough D, Chin MH, Saitz R. Screening and intervention for alcohol problems. A national survey of primary care physicians and psychiatrists. *J Gen Intern Med* 2000;15(2):84–91. [PubMed: 10672110]
- Friedman B, Basu J. The rate and cost of hospital readmissions for preventable conditions. *Med Care Res Rev* 2004;61(2):225–240. [PubMed: 15155053]
- Gill JM, Mainous AG 3rd. The role of provider continuity in preventing hospitalizations. *Arch Fam Med* 1998;7(4):352–357. [PubMed: 9682689]
- Girard DD, Partridge RA, Becker B, Bock B. Alcohol and tobacco use in the elder emergency department patient: assessment of rates and medical care utilization. *Acad Emerg Med* 2004;11(4):378–382. [PubMed: 15064212]
- Hazelton LD, Sterns GL, Chisholm T. Decision-making capacity and alcohol abuse: clinical and ethical considerations in personal care choices. *Gen Hosp Psychiatry* 2003;25(2):130–135. [PubMed: 12676427]
- Health Resources and Services Administration. Primary care health professional shortage areas. from <http://datawarehouse.hrsa.gov/hpsadetail.aspx>
- Himelhoch S, Weller WE, Wu AW, Anderson GF, Cooper LA. Chronic medical illness, depression, and use of acute medical services among Medicare beneficiaries. *Med Care* 2004;42(6):512–521. [PubMed: 15167319]
- Kaner E, Bland M, Cassidy P, Coulton S, Deluca P, Drummond C, et al. Screening and brief interventions for hazardous and harmful alcohol use in primary care: a cluster randomised controlled trial protocol. *BMC Public Health* 2009;9:287. [PubMed: 19664255]
- Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of Illness in the Aged. The Index of Adl: A Standardized Measure of Biological and Psychosocial Function. *JAMA* 1963;185:914–919. [PubMed: 14044222]
- Kunz JL. Alcohol use and reported visits to health professionals: an exploratory study. *J Stud Alcohol* 1997;58(5):474–479. [PubMed: 9273911]
- Laditka JN, Laditka SB, Mastanduno MP. Hospital utilization for ambulatory care sensitive conditions: health outcome disparities associated with race and ethnicity. *Soc Sci Med* 2003;57(8):1429–1441. [PubMed: 12927473]
- Lemeshow S, Letenneur L, Dartigues JF, Lafont S, Orgogozo JM, Commenges D. Illustration of analysis taking into account complex survey considerations: the association between wine consumption and dementia in the PAQUID study. *Personnes Ages Quid. Am J Epidemiol* 1998;148(3):298–306. [PubMed: 9690368]
- Li C, Friedman B, Conwell Y, Fiscella K. Validity of the Patient Health Questionnaire 2 (PHQ-2) in identifying major depression in older people. *J Am Geriatr Soc* 2007;55(4):596–602. [PubMed: 17397440]
- Li Y, Glance LG, Cai X, Mukamel DB. Mental illness and hospitalization for ambulatory care sensitive medical conditions. *Med Care* 2008;46(12):1249–1256. [PubMed: 19300315]
- Logan JE, Riley AW, Barker LE. Parental mental and pain-related health and pediatric ambulatory care sensitive emergency department visits and hospitalizations. *Health Serv Res* 2008;43(2):656–674. [PubMed: 18370972]
- McCall, N. Investigation of increasing rates of hospitalization for ambulatory care sensitive conditions among Medicare Fee-for-Service beneficiaries. RTI International; 2004. CMS o. Document Number
- Merrick EL, Hodgkin D, Garnick DW, Horgan CM, Panas L, Ryan M, et al. Unhealthy drinking patterns and receipt of preventive medical services by older adults. *J Gen Intern Med* 2008;23(11):1741–1748. [PubMed: 18716847]
- Merrick EL, Hodgkin D, Garnick DW, Horgan CM, Panas L, Ryan M, et al. Unhealthy drinking patterns and receipt of preventive medical services by older adults. *J Gen Intern Med* 2008b;23(11):1741–1748. [PubMed: 18716847]

- Merrick EL, Horgan CM, Hodgkin D, Garnick DW, Houghton SF, Panas L, et al. Unhealthy drinking patterns in older adults: prevalence and associated characteristics. *J Am Geriatr Soc* 2008a;56(2): 214–223. [PubMed: 18086124]
- Moore AA, Morgenstern H, Harawa NT, Fielding JE, Higa J, Beck JC. Are older hazardous and harmful drinkers less likely to participate in health-related behaviors and practices as compared with nonhazardous drinkers? *J Am Geriatr Soc* 2001;49(4):421–430. [PubMed: 11347786]
- Moos RH, Brennan PL, Schutte KK, Moos BS. High-risk alcohol consumption and late-life alcohol use problems. *Am J Public Health* 2004;94(11):1985–1991. [PubMed: 15514241]
- NIAAA. National Institute on Alcohol Abuse and Alcoholism. Helping patients who drink too much. A Clinician's Guide. 2005. Edition.. Retrieved June 5, 2008., from <http://pubs.niaaa.nih.gov/publications/Practitioner/CliniciansGuide2005/guide.pdf>
- Niefeld MR, Braunstein JB, Wu AW, Saudek CD, Weller WE, Anderson GF. Preventable hospitalization among elderly Medicare beneficiaries with type 2 diabetes. *Diabetes Care* 2003;26(5):1344–1349. [PubMed: 12716786]
- Oster A, Bindman AB. Emergency department visits for ambulatory care sensitive conditions: insights into preventable hospitalizations. *Med Care* 2003;41(2):198–207. [PubMed: 12555048]
- Ozminkowski RJ, Goetzel RZ, Shechter D, Stapleton DC, Baser O, Lapin P. Predictors of preventive service use among Medicare beneficiaries. *Health Care Financ Rev* 2006;27(3):5–23. [PubMed: 17290645]
- Rathore SS, Wang Y, Druss BG, Masoudi FA, Krumholz HM. Mental disorders, quality of care, and outcomes among older patients hospitalized with heart failure: an analysis of the national heart failure project. *Arch Gen Psychiatry* 2008;65(12):1402–1408. [PubMed: 19047527]
- Rice C, Duncan DF. Alcohol use and reported physician visits in older adults. *Prev Med* 1995;24(3): 229–234. [PubMed: 7644444]
- Rice DP, Conell C, Weisner C, Hunkeler EM, Fireman B, Hu TW. Alcohol drinking patterns and medical care use in an HMO setting. *J Behav Health Serv Res* 2000;27(1):3–16. [PubMed: 10695237]
- Saitz, R. Medical and surgical complications of addiction. In: Graham, AST.; Mayo-Smith, M.; Ries, R.; Wilford, BB., editors. *Principles of Addiction Medicine*. 3. Chevy Chase, Maryland: American Society of Addiction Medicine; 2003. p. 1027-1052.
- Shelkey M, Wallace M. Katz Index of Independence in Activities of Daily Living. *J Gerontol Nurs* 1999;25(3):8–9. [PubMed: 10362969]
- STATA Corporation. STATA version 9.0 College Station, TX.
- Wang MC, Rosen AK, Kazis L, Loveland S, Anderson J, Berlowitz D. Correlation of risk adjustment measures based on diagnoses and patient self-reported health status. *Health Serv Outcomes Res Method* 2000;1(3–4):351–365.
- Weissman JS, Gatsonis C, Epstein AM. Rates of avoidable hospitalization by insurance status in Massachusetts and Maryland. *JAMA* 1992;268(17):2388–2394. [PubMed: 1404795]
- Wooldridge, JM. *Introductory Econometrics*. 3. Thomson Southwestern; 2006.
- Zeng F, O'Leary JF, Sloss EM, Lopez MS, Dhanani N, Melnick G. The effect of medicare health maintenance organizations on hospitalization rates for ambulatory care-sensitive conditions. *Med Care* 2006;44(10):900–907. [PubMed: 17001260]
- Zhao Y, Ash AS, Ellis RP, Slaughter JP. Disease burden profiles: an emerging tool for managing managed care. *Health Care Manag Sci* 2002;5(3):211–219. [PubMed: 12363048]
- Zhao Y, Ellis RP, Ash AS, Calabrese D, Ayanian JZ, Slaughter JP, et al. Measuring population health risks using inpatient diagnoses and outpatient pharmacy data. *Health Serv Res* 2001;36(6 Pt 2): 180–193. [PubMed: 16148968]

Table 1
Sample Description and Distribution Among Drinking Categories

	Total % of sample (weighted column %)	Weighted Row Percent		
		Drinks within guidelines	Exceeds monthly limit only	Heavy episodic drinking
Unweighted n	5,570	4,215	543	812
Weighted N	15,128,450	11,373,766	1,516,640	2,238,044
Total	100.0	75.2	10.0	14.8
<i>Sociodemographics</i>				
Gender *				
Female	44.8	85.1	7.6	7.3
Male	55.2	67.1	12.0	20.9
Age *				
65 to 70 years	39.1	70.5	10.9	18.6
71 to 80 years	44.7	76.6	9.5	13.9
≥81 years	16.2	82.6	9.2	8.1
Hispanic *				
Non-Hispanic	95.7	75.4	10.2	14.4
Race *				
White	93.0	75.5	10.4	14.2
African-American and Other ^d	7.0	72.5	4.8	22.7
Education *				
< H.S. diploma	15.7	70.3	7.3	22.4
H.S. graduate	27.6	75.6	9.3	15.1
Some college/voc/tech	27.9	77.0	10.2	12.8
College degree	28.8	75.6	12.2	12.2
Annual Income *				
< \$25,000	37.5	75.7	7.4	16.9
\$25,000 to \$40,000	27.1	76.2	8.9	15.0
> \$40,000	35.4	73.5	13.1	13.4
Marital Status *				
Married	65.4	74.7	10.5	14.8
Widowed	22.8	79.9	8.4	11.7
Divorced, separated, single	11.8	69.0	10.2	20.8
Metropolitan area *				
Lives in metro area	81.0	76.2	10.0	13.8
Non-metropolitan area	19.0	71.0	10.0	19.0
Region *				
Northeast	21.1	76.7	11.4	12.0
Midwest	27.4	77.5	7.3	15.2

	Total % of sample (weighted column %)	Weighted Row Percent		
		Drinks within guidelines	Exceeds monthly limit only	Heavy episodic drinking
West	19.3	72.9	13.4	13.7
South and Other ^a	32.2	73.7	9.4	16.9
Health and functional status				
Relative health risk[*]				
No risk indication	36.1	72.8	9.5	17.7
< average risk	45.5	76.6	10.2	13.1
> average risk	18.4	76.3	10.5	13.2
Self-reported depression-most of time				
Depressed	8.4	72.3	9.4	18.2
Not depressed	91.6	75.4	10.1	14.5
Mental health diagnosis				
Has mental health dx	7.3	77.9	9.1	13.0
No mental health dx	92.7	75.0	10.1	14.9
Current smoker[*]				
Is current smoker	13.0	57.6	13.0	29.4
Not current smoker	87.0	77.8	9.6	12.6
Functional status				
Independence in 6 ADLs	70.1	74.4	10.3	15.2
Independence in 5 ADLs	22.7	76.4	10.0	13.6
Independence in 0 to 4 ADLs	7.3	79.3	6.5	14.1
Chronic disease (selected)^{**}				
Has chronic disease	56.3	77.0	9.4	13.6
No chronic disease	43.7	73.8	10.5	15.7
Access variables				
Lives in primary care shortage area (PCSA)[*]				
Lives in PCSA	10.1	68.6	11.1	20.4
Not in PCSA	89.9	76.0	9.9	14.2
Trouble getting care				
Had trouble	2.7	80.0	6.3	13.7
Did not have trouble	97.3	75.0	10.1	14.8
Usual source of care[*]				
No usual source (except ER/urgent care)	5.4	63.3	10.6	26.1
Primary care	85.6	76.2	10.1	13.7
Non-primary care	9.0	76.2	9.6	14.1
Private insurance[*]				
Has private insurance	82.9	76.3	10.4	13.3
No private insurance	17.1	69.7	8.2	22.1
Medicaid coverage[*]				

	Total % of sample (weighted column %)	<i>Weighted Row Percent</i>		
		Drinks within guidelines	Exceeds monthly limit only	Heavy episodic drinking
Has Medicaid coverage	4.5	68.4	5.0	26.6
No Medicaid coverage	95.5	75.5	10.3	14.2

^a=Some values not displayed separately due to containing unweighted cell n < 11

Ns for bivariate comparisons varied due to item-missing data, <3.9% for all variables. Percents are for non-missing data.

*
p <.01

**
p <.05

Table 2

Utilization for Ambulatory-Care-Sensitive Conditions (ACSC), By Drinking Pattern

<i>Weighted Percent with Any:</i>			
Unweighted n=5,570 Weighted N= 15, 128,450	ACSC Hospitalization	ACSC Emergency Dept. Visit—no inpatient admission	ACSC Emergency Dept. Visit— resulted in inpatient admission
Overall Percent	3.5	2.1	2.1
Drinking Category		**	
Within guidelines	3.5	1.9	2.2
Drinks over monthly limit only	3.7	2.8	2.3
Heavy episodic	3.8	3.1	1.8
Sociodemographics			
Gender			
Female	3.2	2.1	1.9
Male	3.8	2.2	2.3
Age	*	*	*
65 to 70 years	2.3	1.2	1.6
71 to 80 years	3.8	2.5	2.0
≥81 years	5.8	3.5	3.8
Ethnicity			
Hispanic	3.5	2.3	3.0
Non-Hispanic	5.0	2.1	2.1
Race	*		
White	3.3	2.1	2.0
African-American or Other ^a	6.8	2.6	3.4
Education	*		*
< H.S. diploma	5.5	2.7	3.7
H.S. graduate	3.4	2.4	1.5
Some college/voc/tech	3.5	1.8	2.0
College degree	2.6	1.9	2.0
Annual income	*		*
< \$25,000	5.0	2.2	3.3
\$25,000 to \$40,000	3.3	2.2	1.6
> \$40,000	2.3	2.1	1.2
Marital Status	*		
Married	2.9	2.0	1.8
Widowed	5.2	2.7	2.8
Divorced, separated, single	3.7	1.8	2.4
Metropolitan Area			
In metropolitan area	3.6	2.0	2.3
Not in metro area	3.3	2.7	1.4
Region	**		*
Northeast	2.9	2.4	2.1

<i>Weighted Percent with Any:</i>				
Unweighted n=5,570 Weighted N= 15, 128,450	ACSC Hospitalization	ACSC Emergency Dept. Visit—no inpatient admission	ACSC Emergency Dept. Visit— resulted in inpatient admission	
Midwest	3.5	2.2	1.5	
West	2.5	2.2	1.5	
South and Other ^a	4.6	1.9	3.0	
Health and functional status				
Relative Health Risk	*	*	*	
No risk indication	1.2	1.0	0.8	
< average risk	3.1	1.9	1.7	
> average risk	9.2	4.8	5.8	
Self-reported depression		**	**	
Depressed most of time	4.7	3.7	3.5	
Not depressed most of time	3.4	2.0	2.0	
Mental health diagnosis	*		**	
MH diagnosis	6.4	3.0	3.7	
No MH diagnosis	3.3	2.1	2.0	
Current smoker	*			
Current smoker	4.9	2.0	2.9	
Not current smoker	3.3	2.2	2.0	
Functional status	*	*	*	
Independence in 6 ADLs	2.9	1.9	1.7	
Independence in 5 ADLs	4.4	2.1	2.4	
Independence in 0 to 4 ADLs	7.9	4.8	5.9	
Chronic disease (selected)	*	*	*	
Has chronic disease	5.6	3.1	3.2	
Does not have chronic disease	1.9	1.4	1.3	
Access variables				
Primary care shortage area				
Lives in PCSA	4.2	2.8	2.3	
Not in PCSA	3.4	2.1	2.1	
Trouble getting care	4.9	1.7	3.4	
No trouble getting care	3.5	2.2	2.1	
Usual source of care				
No usual source except ER/urgent care	4.1	1.3	1.5	
Primary care	3.5	2.2	2.1	
Non-primary care	3.1	2.3	3.0	
Private insurance		**		
Private insurance	3.4	2.3	2.0	
No private insurance	4.2	1.3	2.6	
Medicaid coverage	*		**	
Had Medicaid coverage	7.2	1.9	4.4	

<i>Weighted Percent with Any:</i>			
Unweighted n=5,570 Weighted N= 15, 128,450	ACSC Hospitalization	ACSC Emergency Dept. Visit—no inpatient admission	ACSC Emergency Dept. Visit— resulted in inpatient admission
No Medicaid coverage	3.4	2.1	2.0

Weighted sample used; significance based on chi-square tests, corrected for survey design. N for each bivariate comparison varied slightly due to item-missing data, <4% for all variables. Percents shown are for non-missing data.

^a values combined to avoid unweighted cell sizes <11

*
p < .01,

**
p < .05,

ACSC measure varies significantly by independent variable

Table 3

Logistic Regression Results (Weighted): Predictors of hospitalization and emergency department visits for ambulatory-care-sensitive conditions (ACSCs)

	Odds Ratio (95% confidence interval)		
Unweighted n= 5,052 Weighted N=13,677,112	Any ACSC hospitalization	Any ACSC emergency dept. visit—no inpatient admission	Any ACSC emergency dept. visit—resulted in inpatient admission
<i>Drinking Category</i> (ref.: drinks within guidelines)			
Drinks over monthly limit only	1.14 (.63–2.07)	1.53 (.80–2.94)	.91 (.46–1.83)
Heavy episodic	.89 (.53–1.48)	1.91 (1.11–3.30)**	.70 (.36–1.32)
<i>Sociodemographics</i>			
Female	.74 (.52–1.07)	.95 (.56–1.60)	.70 (.47–1.05)
Age (reference: 65–70 yrs)			
71 to 80 years	1.33 (.86–2.04)	1.84 (1.05–3.25)*	1.04 (.62–1.74)
≥81 years	1.77 (1.11–2.81)**	2.41 (1.30–4.47)*	1.54 (.91–2.61)
Hispanic	1.27 (.46–3.54)	.97 (.30–3.10)	.98 (.25–3.80)
Race (ref: white)			
African-American	2.38 (1.23–4.60)**	1.98 (.86–4.56)	1.38 (.65–2.96)
Other	.90 (.33–2.47)	1.59 (.43–5.90)	.66 (.20–2.22)
Education (ref. < H.S. diploma)			
H.S. graduate	.81 (.51–1.29)	.96 (.58–1.61)	.54 (.30–1.00)**
Some college/voc/tech	.83 (.53–1.32)	.79 (.43–1.44)	.79 (.47–1.32)
College degree	.77 (.44–1.35)	.78 (.41–1.46)	.95 (.51–1.75)
Annual income (ref: < \$25,000)			
\$25,000 to \$40,000	.93 (.59–1.48)	1.24 (.75–2.05)	.58 (.32–1.04)
> \$40,000	.74 (.46–1.18)	1.37 (.80–2.34)	.41 (.23–73)*
Marital Status (ref: married)			
Widowed	1.39 (.98–1.97)	1.11 (.64–1.93)	1.01 (.63–1.61)
Divorced, separated, single	1.05 (.64–1.71)	1.26 (.59–2.68)	.94 (.48–1.2)
Metropolitan area	.94 (.63–1.40)	.75 (.45–1.24)	1.37 (.75–2.49)
Region (ref: South)			
Northeast	.68 (.43–1.05)	1.28 (.76–2.16)	.68 (.41–1.12)
Midwest	.79 (.55–1.14)	1.17 (.69–1.98)	.44 (.28–70)*
West	.55 (.33–92)**	.99 (.60–1.64)	.48 (.25–90)**
Other	.83 (.18–3.86)	1.56 (.29–8.44)	1.06 (.13–8.34)
<i>Health and functional status</i>			
Relative Health Risk	1.46 (1.31–1.62)*	1.22 (1.08–1.38)*	1.46 (1.29–1.64)*
Depressed most of time	.97 (.55–1.70)	1.44 (.83–2.52)	1.23 (.62–2.42)
Mental health diagnosis	1.02 to (.60–1.72)	.97 (.48–1.94)	.87 (.43–1.78)

	Odds Ratio (95% confidence interval)		
Unweighted n= 5,052 Weighted N=13,677,112	Any ACSC hospitalization	Any ACSC emergency dept. visit—no inpatient admission	Any ACSC emergency dept. visit—resulted in inpatient admission
Current smoker	1.40 (.83–2.35)	.81 (.40–1.63)	1.54 (.81–2.93)
Functional status (ref: Indep 6 ADLs)			
Independence in 5 ADLs	1.21 (.85–1.72)	1.08 (.64–1.80)	1.28 (.77–2.11)
Independence in 0 to 4 ADLs	1.1 (1.09–3.00) **	2.20 (1.19–4.04) **	2.20 (1.26–3.81) *
Chronic disease	1.89 (1.33–2.69) *	1.64 (1.03–2.62) **	1.48 (.97–2.27)
Access variables			
Lives in primary care shortage area	.86 (.36–2.04)	1.15 (.43–3.05)	.57 (.13–2.45)
Trouble getting care	1.07 (.40–2.86)	.27 (.07–1.05)	.88 (.30–2.54)
Usual source of care (ref=primary care)			
None (except ER/urgent care)	1.31 (.62–2.77)	1.01 (.38–2.74)	.51 (.17–1.52)
Non-primary care	.68 (.39–1.18)	1.29 (.77–2.17)	.88 (.44–1.76)
Private insurance	1.22 (.68–2.19)	1.91 (.88–4.14)	1.06 (.55–2.01)
Medicaid coverage	1.24 (.61–2.51)	.84 (.29–2.41)	0.93 (.38–2.26)
Baseline year 2001	1.10 (.80–1.52)	1.25 (.80–1.97)	1.00 (.64–1.57)

*
p<.01

**
p<.05