Published in final edited form as:

Synthesis (Stuttg). 2010 April 20; 41(16): i-. doi:10.1002/chin.201016160.

A New Route to Substituted Pyrimido[5,4-*e*]-1,2,4-triazine-5,7(1*H*, 6*H*)-diones and Facile Extension to 5,7(6*H*,8*H*) Isomers

Anjanette J. Turbiak and H. D. Hollis Showalter*

Department of Medicinal Chemistry, University of Michigan, Ann Arbor, MI 48109-1065, USA

Abstract

A new route to substituted pyrimido[5,4-*e*]-1,2,4-triazine-5,7(1*H*,6*H*)-diones is outlined. The synthesis proceeds via pre-formed hydrazone intermediates, which are then condensed with an activated chlorouracil to build up the entire molecular framework, followed by a reductive ring closure to provide the parent series. The route has been extended to the isomeric pyrimido[5,4-*e*]-1,2,4-triazine-5,7(6*H*,8*H*)-dione class via the use of methylhydrazine as hydrazine surrogate, followed by regiospecific alkylation of the *N*⁸-H pyrimidotriazinediones with a range of alkyl and alkaryl substituents. This new methodology permits the generation of a wide range of compounds with variable substitution at the *N*¹, *C*³, and *N*⁸ positions for a heterocyclic scaffold with demonstrated pharmacological activity.

Keywords

pyrimido[5,4-*e*]-1,2,4-triazine-5,7(1*H*,6*H*)-diones; pyrimido[5,4-*e*]-1,2,4-triazine-5,7(6*H*,8*H*)-diones; heterocycles; hydrazones; cyclizations

Pyrimido[5,4-*e*]-1,2,4-triazine-5,7-diones such as the natural products toxoflavin (**1**) and fervenulin (**2**) (Figure 1) exhibit antibiotic and other useful pharmacological activity, motivating interest in their exploration since the early 1960s.¹ While a variety of analogues have been synthesized and evaluated within both templates, one subclass conspicuously absent is that with aryl substituents at the *N*¹-position, **3**, whereas analogues with similar substituents at *N*⁸, **4**, have been reported.² We recently have reported on a novel synthesis of compounds related to **1**, including hitherto inaccessible *N*¹-phenyl analogues **3**.³ In this paper, we report on the further application of this synthesis to a variety of *N*¹-alkyl and *N*¹-aryl compounds, and an efficient process for extending this synthesis to the generation of compounds within the isomeric fervenulin series.

The strategy and range of analogues synthesized within the pyrimido[5,4-*e*]-1,2,4-triazine-5,7(1*H*,6*H*)-dione (toxoflavine) series is shown in Table 1. First, monosubstituted hydrazines **5**, either as the free base or hydrochloride salt, were condensed under reflux with aromatic aldehydes **6** in ethanol or tetrahydrofuran to form hydrazones **7**, which usually precipitated out of solution upon cooling. The resulting hydrazones **7** were then treated with 6-chloro-3-methyl-5-nitrouracil (**8**), which had been formed by nitration of commercially available 6-chloro-3-methyluracil according to the published method.⁴ The addition to **8** required catalysis with aluminum trichloride for those hydrazones formed from arylhydrazines. The resultant 6-(2-arylidene-1-substituted-hydrazinyl)-3-methyl-5-nitrouracils **9** were then cyclized to 3-(4-aryl)-6-methyl-1-substituted-pyrimido[5,4-*e*]-1,2,4-triazine-5,7(1*H*,6*H*)-diones **10** upon treatment with zinc dust (4 equiv) and ammonium

chloride (2 equiv) while exposed to the air and with vigorous stirring in refluxing 50% aqueous ethanol. The N¹-substituted pyrimidotriazinediones were then isolated cleanly, usually in >60% yield, by washing with aqueous 1 M hydrochloric acid. Other solvents and combinations thereof were tried for the zinc-mediated cyclization, but the yields were considerably diminished. Further, additional standard methods were evaluated to reduce the nitro group and promote cyclization. These included standard hydrogenation over palladium on carbon, transfer hydrogenation (Pd/C, ammonium formate), tin in ethyl acetate, tin(II) chloride dihydrate in ethanol, iron in glacial acetic acid, and zinc/hydrochloric acid in ethanol, but none provided clean product.

For Ar and R = aryl, the method was amenable to a variety of substituents on either ring (Table 1, entries 1–8). Both electron-donating and electron-withdrawing substituents were tolerated on either component and there was variation in overall yield depending on the combination of moieties examined. However, the hydrazone formed from 2-pyridylhydrazine was unreactive, perhaps due to catalyst deactivation via complexation of the adjacent pyridyl nitrogen with aluminum trichloride. An attempt to conduct this reaction in the absence of catalyst and at higher temperatures led only to recovered hydrazone.

In optimizing the reaction sequence, it was found that hydrazones **7**, where R = aryl, did not need to be isolated in a separate step (Table 1, Method A) and that yields of **9** were often improved by forming the hydrazone in situ. Hence, this became the preferred method. In the improved protocol (Table 1, Method B), 6-chloro-3-methyl-5-nitrouracil (**8**) and aluminum trichloride were added after 1–2 hours of reaction time between the arylhydrazine **5** and arylaldehyde **6** in an appropriate solvent. After a solvent screen that included standard chlorinated solvents (no reaction) and acetonitrile (modest reaction), tetrahydrofuran was found to provide superior yields. In addition to arylhydrazines, the methodology was investigated for a variety of alkylhydrazines **5** (R = alkyl), and here the hydrazones **7** formed in situ were found to be sufficiently reactive with chlorouracil **8** that activation with aluminum trichloride was unnecessary. Table 1 (Method C; entries 9–13) summarizes reactions for successful alkylhydrazines. In addition to those shown, the hydrazones of methylhydrazine also reacted quickly and efficiently with **8**; however, in the subsequent cyclization of **9** to **10** with zinc and ammonium chloride, demethylation at N¹ was a significant side reaction. Varying the reaction time and temperature could minimize this, but it was never completely suppressed. Hence, for R = methyl, the preferred method of synthesis would be one of the standard methods commonly used to make N¹-methylpyrimidotriazinediones,⁵ but for most other R substituents, our new methodology is advantageous with regard to both synthesis time and efficiency. Moreover, for Methods B and C, the sequence of in situ formation of hydrazone **7** followed by condensation with chloronitrouracil **8** and subsequent cyclization to **10** can be carried out with yields comparable to or better than when **7** is isolated separately.

There are some shortcomings to this methodology, however. The reaction between hydrazones **7** and 6-chloro-3-methyl-5-nitrouracil (**8**) was the more general of the last two steps. An example where the desired Michael-type addition to **8** did not occur included the hydrazone of *tert*-butylhydrazine. For the cyclization of **9** to **10**, reaction conditions were robust for aryl and some alkyl substituents at the N¹-position. However, problems were encountered for both benzyl and methyl (as described above) substituents. While N¹-methylpyrimidotriazinediones could be isolated to some extent, despite the competing demethylation, no desired N¹-benzylpyrimidotriazinediones **10** could be detected as debenzylation appeared to rapidly follow cyclization. A similar outcome was observed for R = (CH₂)₂NEt₂ with N¹-C bond cleavage likely occurring via an aziridinium intermediate.

The demethylation that occurred when R = Me in the formation of **10** could be used to advantage, however, when the goal was an N⁸-H pyrimidotriazinedione. While hydrazine itself could be used in the initial condensation with aldehydes **6** to form hydrazones **7** (R = H), subsequent reaction of these with chlorouracil **8** followed by reductive ring closure with zinc and ammonium chloride never provided clean N¹(N⁸)-H pyrimidotriazinediones. Hence, methylhydrazine could be used as a surrogate to attain the same objective. Table 2 lists several examples where 3-aryl-substituted pyrimidotriazinediones were made in this two-pot three-step process. Here, slightly lower yields were obtained in the ring closure/demethylation step when there was an electron-donating substituent at the 4-position of the aryl substituent (Table 2, entries 6 and 7), whereas the highest yields were obtained with the strongly electron-withdrawing fluoro group at the same position (Table 2, entries 3 and 8). Both of these observations are consistent with the ease with which demethylation of N¹-methylpyrimidotriazinediones occurred. This involves a likely mechanism of S_N2 nucleophilic attack of the N¹-methyl group by solvent followed by C–N bond cleavage with the core pyrimidotriazinedione heterocycle serving as a leaving group.

The demethylated pyrimidotriazinediones could then be alkylated to produce a variety of novel N⁸-substituted pyrimidotriazinedione analogues, as shown in Table 3. This reaction, which has been previously documented in the literature,⁶ can be subjected to conditions that provide either N¹- or N⁸-alkyl products or mixtures thereof. Our conditions, which involve the use of cesium carbonate as base, provide N⁸ products exclusively for a wide range of alkyl and alkaryl substituents. While yields are variable, these were not optimized except for **11i** and **11j**.

In summary, we have discovered a facile route to pyrimido[5,4-*e*]-1,2,4-triazine-5,7-diones with variable substitution at the N¹, C³, and N⁸ positions, including previously undisclosed N¹-aryl variants. We also have exemplified the use of methylhydrazine as a surrogate for hydrazine to provide an entry to a variety of N¹(N⁸)-unsubstituted congeners, which can be mildly and regioselectively functionalized at the N⁸-position with alkyl and alkaryl halides. We expect this chemistry to find wide application toward the discovery of novel pharmacological agents.

All starting materials were obtained from commercial suppliers and were used without further purification. THF was distilled prior to use over Na/benzophenone. Reactions were run under a blanket of N₂ unless specified otherwise. Glassware was oven-dried before use for reactions run under anhydrous conditions. Melting points were determined in open capillary tubes on a Laboratory Devices Mel-Temp apparatus and are uncorrected. The NMR spectra were recorded on a Bruker instrument at 500 MHz for ¹H, 125 MHz for ¹³C, and 470 MHz for ¹⁹F spectra. Mass spectra were recorded on a Micromass TofSpec-2E Matrix-Assisted, Laser-Desorption, Time-of-Flight mass spectrometer.

All known compounds had spectroscopic (NMR and MS) and physical properties identical to those given in the literature. There are preparative procedures in the literature for the synthesis of hydrazones **7a**,⁷ **7b**⁸ and **7c**⁹ of Table 1, (hydrazinyl)nitropyrimidinediones **9n**,¹⁰ **9q**¹⁰ and **9s**¹⁰ of Table 2, pyrimidinotriazines **10n**,¹⁰ **10q**¹⁰ and **10s**¹⁰ of Table 2, and alkylated pyrimidinotriazines **11a**¹¹ and **11c**¹² of Table 3.

2-Substituted 1-(4-Arylidene)hydrazines **7**; General Procedure for Method A

A mixture of aryl- or alkylhydrazine **5**, arylaldehyde **6** (1.1 equiv), and abs EtOH (0.5–0.7 M) was heated at reflux for 90 min. The soln was left standing at r.t. for several hours and the formed precipitate was collected by filtration, rinsed thoroughly with EtOH, and dried. It was used directly in the next step.

1-(4-Chlorobenzylidene)-2-phenylhydrazine (7c)⁹

Pale peach powder; yield: 290 mg (52%); mp 124–126 °C.

¹H NMR (DMSO-*d*₆): δ = 6.76 (t, *J* = 7.3 Hz, 1 H), 7.07 (d, *J* = 8.2 Hz, 1 H), 7.22 (dd, *J* = 7.5 Hz, *J'* = 8.1 Hz, 2 H), 7.44 (d, *J* = 8.2 Hz, 2 H), 7.66 (d, *J* = 8.5 Hz, 2 H), 7.85 (s, 1 H), 10.45 (s, 1 H).

¹³C NMR (DMSO-*d*₆): δ = 112.52, 119.42, 127.60, 129.13, 129.59, 132.52, 135.29, 135.43, 145.54.

MS: *m/z* = 231.1 (M + H).

6-(2-Arylidene-1-substituted-hydrazinyl)-3-methyl-5-nitropyrimidine-2,4(1*H*, 3*H*)-diones 9; General Procedure for Method A

Hydrazone **7** was suspended in anhyd THF (0.3M), and treated with AlCl₃ (1.0 equiv) and chlorouracil **8** (0.9 equiv). The mixture was heated at reflux for 4 h, during which time it became homogeneous and usually a darker color in appearance. The mixture was cooled on ice for several hours and the thus formed precipitate was collected by filtration, rinsed successively with THF and EtOH, and dried.

6-[2-(4-Chlorobenzylidene)-1-phenylhydrazinyl]-3-methyl-5-nitropyrimidine-2,4(1*H*,3*H*)-dione (9c)

Yellow powder; yield: 95 mg (63%); mp 210–212 °C (dec).

¹H NMR (DMSO-*d*₆): δ = 3.14 (s, 3 H), 7.42 (d, *J* = 7.7 Hz, 2 H), 7.50 (m, 3 H), 7.58 (m, 3 H), 7.70 (d, *J* = 7.7 Hz, 2 H), 11.90 (br s, 1 H).

¹³C NMR (DMSO-*d*₆): δ = 27.72, 112.52, 122.27, 127.83, 128.78, 129.11, 130.22, 131.17, 132.73, 134.97, 135.29, 142.61, 145.53, 157.75.

MS: *m/z* = 422.1 (M + Na).

6-(2-Arylidene-1-substituted-hydrazinyl)-3-methyl-5-nitropyrimidine-2,4(1*H*, 3*H*)-diones 9 via In Situ Generation of Hydrazone **7; General Procedure for Methods B and C**

Method B: A mixture of aryl- or alkylhydrazine **5** and arylaldehyde **6** (1.1 equiv) was dissolved in anhyd THF (0.3 M) and the soln was heated at reflux for 1.5 h. The soln was cooled to r.t, and chlorouracil **8** (0.9 equiv) was added, followed by AlCl₃ (1.0 equiv). The mixture was returned to reflux for 4 h and then cooled on ice for several hours. The formed precipitate was collected by filtration, rinsed with THF and EtOH, and dried.

Method C: The same initial steps for Method B were followed, except that the catalyst was omitted. A precipitate began to form within 5–10 min after addition of **8**. The mixture was heated for an additional 20–45 min. Further processing as per Method B provided product **9**.

6-[2-(4-Fluorobenzylidene)-1-(4-methoxyphenyl)hydrazinyl]-3-methyl-5-nitropyrimidine-2,4(1H,3H)-dione (9h)

Fluorescent yellow powder; yield: 556 mg (73%); mp 228–229 °C.

¹H NMR (DMSO-*d*₆): δ = 3.15 (s, 3 H), 3.86 (s, 3 H), 7.14 (d, *J* = 8.85 Hz, 2 H), 7.23 (dd, *J* = 8.85 Hz, *J'* = 8.85 Hz, 2 H), 7.33 (d, *J* = 8.8 Hz, 2 H), 7.46 (s, 1 H), 7.75 (dd, *J* = 8.3 Hz, *J'* = 5.7 Hz, 2 H).

¹³C NMR (DMSO-*d*₆): δ = 27.66, 55.99, 115.83, 124.39, 126.33, 129.68, 130.12, 131.64, 143.71, 147.29, 149.11, 158.83, 160.83, 162.18, 165.48.

¹⁹F NMR (DMSO-*d*₆): δ = −107.2.

MS: *m/z* = 414.1 (M + H).

6-[1-Isopropyl-2-(4-methoxybenzylidene)hydrazinyl]-3-methyl-5-nitropyrimidine-2,4(1H,3H)-dione (9m)

Pastel yellow powder; yield: 74 mg (80%); mp 280–285 °C (dec).

¹H NMR (DMSO-*d*₆): δ = 1.40 (d, *J* = 6.1 Hz, 6 H), 3.20 (s, 3 H), 3.79 (s, 3 H), 4.76 (m, 1 H), 6.97 (d, *J* = 8.3 Hz, 2 H), 8.15 (d, *J* = 8.2 Hz, 2 H), 12.25 (s, 1 H).

¹³C NMR (DMSO-*d*₆): δ = 22.04, 27.57, 49.37, 55.59, 95.54, 113.88, 124.76, 129.86, 143.21, 147.78, 151.18, 158.64, 160.04.

MS: *m/z* = 362.1 (M + H), 384.0 (M + Na).

6-[2-(4-Fluoro-3-methoxybenzylidene)-1-methylhydrazinyl]-3-methyl-5-nitropyrimidine-2,4(1H,3H)-dione (9u)

Pastel yellow powder; yield: 295 mg (84%); mp 218–222 °C.

¹H NMR (DMSO-*d*₆): δ = 3.15 (s, 3 H), 3.41 (s, 3 H), 3.95 (s, 3 H), 7.35 (m, 1 H), 7.55 (m, 1 H), 7.90 (d, *J* = 8.5 Hz, 1 H), 8.71 (s, 1 H), 11.30 (br s, 1 H).

¹³C NMR (DMSO-*d*₆): δ = 27.63, 34.98, 56.34, 111.01, 116.49, 122.42, 131.20, 131.65, 142.36, 149.47, 157.62.

¹⁹F NMR (DMSO-*d*₆): δ = −129.7.

MS: *m/z* = 350.1 (M − H).

Anal. Calcd for C₁₄H₁₄FN₅O₅: C, 47.87; H, 4.02; N, 19.94. Found: C, 47.82; H, 3.89; N, 19.87.

Pyrimidotriazinediones 10; General Procedure

Table 1: To a suspension of **9** in 50% aq EtOH (0.2 M) was added Zn dust (4.0 equiv) and NH₄Cl (2.0 equiv). The vigorously stirred suspension was heated at reflux overnight with exposure to the air. The mixture was cooled to r.t. and diluted with aq 1 M HCl (1–3 mL per mmol of Zn used). After stirring at r.t. for 1 h, the formed precipitate was collected by filtration, washed successively with aq 1 M HCl and EtOH, and dried to provide **10**.

Table 2: The same procedure as described above was followed except that reaction was carried out in 50% aq DMF (0.08–0.1 M).

1-(4-Fluorophenyl)-3-(4-methoxyphenyl)-6-methylpyrimido[5,4-*e*]-1,2,4-triazine-5,7(1*H*,6*H*)-dione (10e)

Bright red-orange solid; yield: 188 mg (79%); mp 310–315 °C (dec).

¹H NMR (DMSO-*d*₆): δ = 3.29 (s, 3 H), 3.86 (s, 3 H), 7.14 (d, *J* = 8.8 Hz, 2 H), 7.51 (t, *J* = 8.7 Hz, 2 H), 7.82 (dd, *J*₁ = 8.8 Hz, *J*₂ = 4.9 Hz, 2 H), 8.14 (d, *J* = 8.8 Hz, 2 H).

¹³C NMR (DMSO-*d*₆): δ = 26.03, 52.04, 112.45, 115.67, 117.07, 119.19, 124.06, 124.19, 128.12, 129.45, 144.84, 154.96, 162.24.

¹⁹F NMR (DMSO-*d*₆): δ = −105.6.

MS: *m/z* = 380.1 (M + H), 402.1 (M + Na).

Anal. Calcd for C₁₉H₁₄FN₅O₃: C, 60.16; H, 3.72; N, 18.46. Found: C, 59.88; H, 3.50; N, 18.19.

1-(2-Hydroxyethyl)-6-methyl-3-(4-methylphenyl)pyrimido[5,4-*e*]-1,2,4-triazine-5,7(1*H*,6*H*)-dione (10i)

Bright reddish-orange powder; yield: 57 mg (70%); mp 199–201 °C (dec).

¹H NMR (DMSO-*d*₆): δ = 2.41 (s, 3 H), 3.34 (s, 3 H), 3.93 (t, *J* = 5.2 Hz, 2 H), 4.53 (t, *J* = 5.3 Hz, 2 H), 4.98 (br s, 1 H), 7.39 (d, *J* = 7.3 Hz, 2 H), 8.21 (d, *J* = 7.3 Hz, 2 H).

¹³C NMR (DMSO-*d*₆): δ = 21.50, 28.75, 57.15, 58.09, 127.07, 130.28, 132.64, 141.67, 146.92, 149.95, 151.75, 154.59, 159.49.

MS: *m/z* = 314.1 (M + H).

3-(4-Methoxyphenyl)-6-methylpyrimido[5,4-*e*]-1,2,4-triazine-5,7(6*H*,8*H*)-dione (10s)¹⁰

Mustard yellow powder; yield: 897 mg (43%); mp >325 °C.

¹H NMR (TFA-*d*): δ = 3.55 (s, 3 H), 3.90 (s, 3 H), 7.09 (d, *J* = 7.2 Hz, 2 H), 8.31 (d, *J* = 7.2 Hz, 2 H).

¹³C NMR (TFA-*d*): δ = 19.49, 45.95, 116.54, 121.91, 132.06, 138.08, 147.51, 150.95, 161.72, 166.36.

MS: *m/z* = 284.1 (M − H).

8-Alkylpyrimido[5,4-*e*]-1,2,4-triazine-5,7(6*H*,8*H*)-diones 11; General Procedure

A mixture of 3-arylpyrimidotriazinedione **10**, Cs₂CO₃ (1.5 equiv), alkyl halide (1.2 equiv), and acetone (0.1 M) was stirred at r.t. overnight (bromides) or heated to 50 °C in a sealed vessel overnight (chlorides). The mixture was diluted with H₂O, and refrigerated for 3–4 h. The resulting precipitate was collected, washed with H₂O, and dried to provide **11**.

8-(3-Fluorobenzyl)-6-methyl-3-phenylpyrimido[5,4-e]-1,2,4-triazine-5,7(6H,8H)-dione (11g)

Yield: 47 mg (66%); mp 229–232 °C.

¹H NMR (DMSO-*d*₆): δ = 3.40 (s, 3 H), 5.60 (s, 2 H), 7.11 (t, *J* = 7.2 Hz, 1 H), 7.31 (m, 2 H), 7.39 (m, 1 H), 7.63 (d, *J* = 2.6 Hz, 3 H), 8.43 (m, 2 H).

¹³C NMR (DMSO-*d*₆): δ = 29.31, 45.42, 114.42, 123.70, 127.67, 129.77, 130.80, 131.99, 134.42, 139.46, 149.93, 150.19, 159.99, 160.20, 160.51.

MS: *m/z* = 364.1 (M + H).

Acknowledgments

A.J.T. gratefully acknowledges financial support of the American Chemical Society Division of Medicinal Chemistry, Sheila B. Cresswell, and Fred and Dee Lyons Graduate Fellowships. This publication was also made possible by grant number GM007767 from NIGMS. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of NIGMS.

References

1. (a) Lacrampe, JFA.; Connors, RW.; Ho, CY.; Richardson, A.; Freyne, EJE.; Buijnsters, PJJ.; Bakker, AC. WO. 2004,007,498. 2004. (b) Nagamatsu T, Yamasaki H, Hirota T, Yamato M, Kido Y, Shibata M, Yoneda F. Chem Pharm Bull 1993;41:362. [PubMed: 8500202] (c) Liao T, Baiocchi F, Cheng CC. J Org Chem 1966;31:900. [PubMed: 5907865] (d) Levenberg B, Linton SN. J Biol Chem 1966;241:846. [PubMed: 5905124]
2. Nagamatsu T, Yamasaki H, Yoneda F. Heterocycles 1994;37:1147.
3. Turbiak AJ, Showalter HDH. Tetrahedron Lett 2009;50:1996.
4. Black HT. J Heterocycl Chem 1987;24:1373.
5. (a) Nagamatsu T, Yamasaki H, Hirota T, Yamato M, Kido Y, Shibata M, Yoneda F. Chem Pharm Bull 1993;41:362. [PubMed: 8500202] (b) Yoneda F, Nagamatsu T. Chem Pharm Bull 1975;23:2001. [PubMed: 171091] (c) Yoneda F, Nagamatsu T, Shinomura K. J Chem Soc, Perkin Trans 1976;1:713. [PubMed: 944191]
6. (a) Nagamatsu T, Yamasaki H. J Chem Soc, Perkin Trans 2001;1:130. (b) Yoneda F, Nagamatsu T. J Heterocycl Chem 1974;11:271. (c) Yoneda F, Nagamatsu T. Tetrahedron Lett 1973;14:1577.
7. Dabideen DR, Cheng KF, Aljabari B, Miller EJ, Pavlov VA, Al-Abed Y. J Med Chem 2007;50:1993. [PubMed: 17385848]
8. Yao HC, Resnick P. J Org Chem 1965;30:2832.
9. Deng X, Mani NS. J Org Chem 2008;73:2412. [PubMed: 18278943]
10. Yoneda F, Nagamatsu T. Synthesis 1975:177.
11. Yoneda F, Higuchi M, Nitta Y. J Heterocycl Chem 1980;17:869.
12. Yoneda F, Nagamatsu T. J Heterocycl Chem 1974;11:271.

Figure 1.
Structures of natural and synthetic pyrimidotriazinediones

Substituted Pyrimido[5,4-*e*]-1,2,4-triazine-5,7-(1*H*,6*H*)-diones **10** from Condensation of 6-Chloro-3-methyl-5-nitrotracil (**8**) with Hydrazones **7** Followed by Reductive Ring Closure of **9**

Entry	Ar	R	Product	Yield ^d (%)	Product	Yield (%) from 9	Overall yield (%)
1	4-HOC ₆ H ₄	Ph	9a	30, ^b 72 ^c	10a	58	17, ^b 41 ^c
2	4-MeOC ₆ H ₄	Ph	9b	41, ^b 65 ^c	10b	78 ^d	32, ^b 51 ^c
3	4-ClC ₆ H ₄	Ph	9c	33, ^b 53 ^c	10c	72 ^d	24, ^b 38 ^c
4	4-HOC ₃ H ₄	4-FC ₆ H ₄	9d	82 ^c	10d	82	67
5	4-MeOC ₆ H ₄	4-FC ₆ H ₄	9e	71 ^c	10e	79	56
6	5-methyl-2-furyl	4-FC ₆ H ₄	9f	38 ^c	10f	61	23
7	4-MeOC ₆ H ₄	4-BrC ₆ H ₄	9g	31 ^c	10g	76	24
8	4-FC ₆ H ₄	4-MeOC ₆ H ₄	9h	73 ^c	10h	40	29
9	4-MeC ₆ H ₄	(CH ₂) ₂ OH	9i	66 ^e	10i	70	46
10	3-ClC ₆ H ₄	(CH ₂) ₂ OH	9j	68 ^e	10j	62	42
11	3,4-(MeO) ₂ C ₆ H ₃	(CH ₂) ₂ OH	9k	70 ^e	10k	56	39
12	4-MeOC ₆ H ₄	(CH ₂) ₂ CHMe ₂	9l	52 ^e	10l	62 ^d	32
13	4-MeOC ₆ H ₄	<i>i</i> -Pr	9m	80 ^e	10m	51	41
14	various	Me	— ^e				

^aOverall yield for first two steps.

^bHydrazone isolated in separate step prior to reaction with **8** with AlCl₃ catalysis (Method A).

^cHydrazone formation in situ and subsequent reaction with **8** with AlCl₃ catalysis (Method B). $d_{\text{Ref. 3.}}$

^e See Table 2. Hydrazone formation in situ and subsequent reaction with **8** without AlCl₃ catalysis (Method C).

Table 2

*N*⁸-H Pyrimido[5,4-*e*]-1,2,4-triazine-5,7-(6*H*,8*H*)-diones **10** from Methylhydrazine as a Hydrazine Surrogate

Entry	Ar	Product ^a	Yield (%)	Product	Yield (%) from 9	Overall yield (%)
1	Ph	9n	75	10n	50	38
2	3-FC ₆ H ₄	9o	82	10o	61	50
3	4-FC ₆ H ₄	9p	87	10p	79	68
4	4-ClC ₆ H ₄	9q	78	10q	63	49
5	3-MeOC ₆ H ₄	9r	72	10r	65	46
6	4-MeOC ₆ H ₄	9s	90	10s	43	38
7	3-F-4-MeOC ₆ H ₃	9t	85	10t	47	40
8	4-F-3-MeOC ₆ H ₃	9u	84	10u	86	72

^aHydrazone formation in situ and subsequent reaction with **8** without AlCl₃ catalysis (Method C).

Table 3

N⁸-Substituted Pyrimido[5,4-*e*]-1,2,4-triazine-5,7-(6*H*,8*H*)-diones **11** from Regiospecific Alkylation of **10**

Entry	Substrate	X	R	Product	Yield (%)
1	10q	Cl	Pr	11a	47
2	10s	OMe	(CH ₂) ₂ NEt ₂ ^a	11b	22
3	10n	H	(CH ₂) ₂ OH	11c	43
4	10n	H	(CH ₂) ₂ Ph	11d	39
5	10p	F	4- <i>t</i> -BuC ₆ H ₄ CH ₂	11e	71
6	10q	Cl	2-FC ₆ H ₄ CH ₂	11f	57
7	10n	H	3-FC ₆ H ₄ CH ₂	11g	66
8	10s	OMe	4-FC ₆ H ₄ CH ₂	11h	81
9	10p	F	3,4-F ₂ C ₆ H ₃ CH ₂	11i	91
10	10s	OMe	3,4-F ₂ C ₆ H ₃ CH ₂	11j	87

^a Chloride reacted instead of bromide.