CORRESPONDENCE

The Changing Prevalence of Attention-Deficit/Hyperactivity Disorder and Methylphenidate Prescriptions: A Study of Data From a Random Sample of Insurees of the AOK Health Insurance Company in the German State of Hesse, 2000–2007

by Dr. rer. soc. Ingrid Schubert, Ingrid Köster, Prof. Dr. med. Gerd Lehmkuhl
in volume 36/2010

Is the Treatment the Problem?

The article would have benefited by presenting the data — as it means a restriction of the entire population—as e.g. „prevalence of consultations of AOK members“. The sample displayed in the study excluded AOK insured persons with ADHD who did not utilise the healthcare system during that time period. So the displayed prevalence of consultations is definitely below the actual overall prevalence. Similarly, the selection bias of the AOK population should be discussed since some social classes are underrepresented. To exclude such bias phenomena, prevalence analysis should be limited to the total prevalence from cross sectional studies.

With regard to the prescription frequencies: “The treatment of children and adolescents with methylphenidate ... has been described as an important public health problem.”

This statement solely shows the authors’ emotional attitude to using methylphenidate (MPH). Sticking to the facts it is irritating to call the indication conform use of a licensed drug in an enormously impairing pathology a problem. Is MPH treatment a problem? A discussion would be desirable.

Compared with the total prevalence of 5–7% (2), the treatment prevalence of 1–1.5% in the comparison groups in the study results in a proportion of the medication of 15–20%. Compared with other psychiatric disorders this is rather low and does not give rise to the suspicion that too many patients with the disorder receive medication without a good reason.

Regarding the increase in overall amount of MPH that is being prescribed: the longer duration of treatment up to adulthood follows recent scientific realization that hyperactivity disappears around the 10th birthday whereas the disorder does not. A comparable insight leads to the increased daily dosages. The result of prescribing the drug beyond the time the patients spend at school means that 4–6 additionally treated hours per day increase the prescribed quantities of the drug.

REFERENCES

1. Schubert I, Köster I, Lehmkuhl G: The changing prevalence of attention-deficit/hyperactivity disorder and methylphenidate prescriptions: A study of data from a random sample of insurees of the AOK health insurance company in the German state of Hesse, 2000–2007. *Dtsch Arztebl Int* 2010; 107(36): 615–21.
2. Schlack R, Hölling H, Kurth BM, Huss M: Die Prävalenz der Aufmerksamkeitsdefizit-/Hyperaktivitätsstörung (ADHS) bei Kindern und Jugendlichen in Deutschland. *Bundesgesundheitsbl – Gesundheitsforsch – Gesundheitsschutz* 2007; 50: 827–35.

Dr. med. Karsten Dietrich
Hoeftstraße 32
29525 Uelzen
praxis.dietrich@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

What About the Dam Break Upstream?

Of note is the fact that the number of diagnoses in the 7-year period under study decreased for the age group of 0–5 years (total minus 41), whereas all other age groups showed an almost exponential increase (1). This absolutely cannot be correct if ADHD is a genetically determined disorder as onset would have to occur in early infancy! The association with increased prescription of methylphenidate (MPH) is obvious and points to the fact that the diagnosis of ADHD is inversely—that is, incorrectly—made on the basis of the effects of the medication. With regard to the topic of neuro-enhancement, the authors are too cautious in their assumption that MPH is not used according to the indication. This has always been the case because the diagnosis is made phenomenologically and therefore influenced by subjective interpretations.

With regard to my 35 years of experience with ADHD: in the early 1970s, the “hyperkinetic child” was as good as unknown in Germany. Initial reports about this phenomenon were submitted at the 2nd neuropediatric congress in Kiel in 1976 (2). The differential diagnosis of pediatric hyperkinetic disorder rather than ADHD (neurotic restless disorder, vegetative restless disorder) is hardly considered nowadays. Dr Eichlseder from Munich (3), who pioneered stimulant treatment in Germany, was simply ignored when he expressed serious reservations regarding stimulants.

There are many indications that the path of least resistance is chosen (too) often these days, with a diagnosis of ADHD and subsequent prescription of MPH. The often stipulated multimodal treatment almost never takes place because it takes time and effort. The cause for the many “ADHD children” with behavioral disorders is founded in the changing social and familial environment of our times. Initial studies from the United States have discussed this association.

DOI: 10.3238/arztebl.2010.0918a

Schlack (4) from Bonn has provided a fitting metaphor: many (MPH) helpers jump into a racing river in order to save the drowning children, but no one pays any attention to the dam break upstream.

DOI: 10.3238/arztebl.2010.0918b

REFERENCES

1. Schubert I, Köster I, Lehmkuhl G: The changing prevalence of attention-deficit/hyperactivity disorder and methylphenidate prescriptions: A study of data from a random sample of insureds of the AOK health insurance company in the German state of Hesse, 2000–2007. *Dtsch Arztebl Int* 2010; 107(36): 615–21.
2. Aktuelle Neuropädiatrie – 2. Jahrestagung der Gesellschaft für Neuropädiatrie, Kiel Hrsg. Hermann Dose, Thieme Verlag Stuttgart 1977.
3. Eichlseder W: Zur Behandlung konzentrationsgestörter hyperaktiver Kinder mit DL-Amphetamin. *Päd Praxis* 1974; 14: 109–23.
4. Schlack HG: ADHS: eine Soziogene Epidemie? *Kinderärztl Praxis Sonderheft ADHS* 2004: 6–9.

Dr. med. Joachim Börner
Adenauerring 45
49393 Lohne
dr.med.boerner@ewetel.net

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Uncritical Prescription

If less methylphenidate is prescribed in countries such as France, then this is explained with underdiagnosis or with belonging to an academic social stratum.

A recent Swedish study has shown that social class is important (1).

7960 children aged 6–19 years were investigated who had been prescribed psychostimulant drugs in 2006. They were compared with a national cohort of 1.1 million children.

It turned out that:

- Children of parents living on social security benefits had received psychostimulants 2.3 more often,
- Children of mothers of a low educational status had received psychostimulants 2.3 more often,
- Children of single parents had received psychostimulants 1.5 more often,
- Children of parents who were registered because of a psychiatric disorder had received psychostimulants 1.5 more often than in the national cohort.
- The mentioned socioeconomic and psychosocial factors in combination accounted for 60% of prescriptions. In view of the burden on the families, it is not surprising that the children were restless and their concentration was poor.

Uncritical prescribing of ever increasing daily doses of methylphenidate even in older age groups should give us cause for concern. Are psychosocial background factors overlooked in favor of overhasty prescription of psychostimulant drugs? Can the assumption of neuropsychiatric causes be sustained?

A study (2) that is reported in the book “ADHS-Symptome verstehen—Beziehungen verändern [Understanding the symptoms of ADHD—changing relationships]” shows a different way forward. Of 93 children who received at least 10 sessions of child and family therapy, a minimum of 6 months after the study was completed, only one child had to temporarily take stimulants, and that was because of an acute crisis in his family.

The therapy had taught the children to verbalize their inner states, enabling them to get by without conspicuous behavior.

DOI: 10.3238/arztebl.2010.0919a

REFERENCES

1. Hjern A, et al.: Social adversity predicts ADHD-medication in school children – a national cohort study. *Acta paediatrica* 2010; 99: 920–4.
2. Neraal T, Wildermuth M: ADHS: Symptome verstehen – Beziehungen verändern. Giessen: Psychosozial-Verlag 2008.
3. Schubert I, Köster I, Lehmkuhl G: The changing prevalence of attention-deficit/hyperactivity disorder and methylphenidate prescriptions: A study of data from a random sample of insureds of the AOK health insurance company in the German state of Hesse, 2000–2007. *Dtsch Arztebl Int* 2010; 107(36): 615–21.

Dr. med. Terje Neraal
Höhenstraße 33 c
35435 Wettenberg
t.neraal@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

In Reply:

The authors thank their correspondents for their feedback. Dietrich rightly points out that the data from the health insurance companies document the disorder only for insureds with contact to their doctors and suggests to specify as prevalence estimates as “prevalence of consultations.” We dealt with this aspect in our article with the terms “treatment prevalence” or “administrative” treatment prevalence.

We agree with Dietrich that these administrative prevalence rates are lower than those obtained from population surveys, such as the Robert Koch-Institute’s German Health Interview and Examination Survey for Children and Adolescents (Kinder-und Jugend-Gesundheitssurvey, KiGGS) (1), which may include unknown cases and patients who have not consulted a doctor. However, we regard the routine data that have no selection bias or recall bias as an important addition to the surveys, which can only be conducted at greater time intervals owing to their high cost.

In the discussion section we point out that treatment with methylphenidate is regarded as a public health problem in many countries. From the context it is clear that this statement relates to the increase in prescriptions observed in many countries and thus to the background phenomenon of an increase in diagnoses with all its social, medical, and health economic problems.

At no point in the article did we refer to indication conform treatment as a problem.

We showed in our study that not every coded diagnosis results in a prescription for stimulants. However, the proportion of prescriptions for drug treatment has risen from 35.5% in 2000 to 48% in 2007. Dietrich's assumption that other psychiatric disorders have higher prevalence rates of drug treatment does not provide any clues about the quality with which the indication was defined. Börner also mentions how difficult it is to reach a diagnosis and assumes that the diagnosis is possibly made only on the basis of the effects of MPH. We cannot comment on this because we cannot conclude this from the available data.

The extent to which ADHD is solely genetically determined or whether it is multifactorial in origin (2) was not the subject of our study and we can therefore not discuss it here. However, as Neraal illustrates by explaining the results of a recent Swedish study, many studies have shown an association between socioeconomic class and the development of psychiatric disorders. This applies to Germany too, as detailed in the KiGGS: the lower someone's socioeconomic status, the more often they received a diagnosis of ADHD (1).

DOI: 10.3238/arztebl.2010.0919b

REFERENCES

1. Robert-Koch-Institut (ed): Lebensphasenspezifische Gesundheit von Kindern und Jugendlichen in Deutschland. Ergebnisse des Nationalen Kinder- und Jugendgesundheitssurveys (KiGGS). Beiträge zur Gesundheitsberichterstattung des Bundes. Robert Koch-Institut Berlin, 2008.
2. Sachverständigenrat für die Begutachtung der Entwicklung im Gesundheitswesen. Koordination und Integration – Gesundheitsversorgung in einer Gesellschaft des längeren Lebens. Sondergutachten 2009 [<http://www.svr-gesundheit.de/Gutachten/Gutacht09/gutacht09.htm>]
3. Schubert I, Köster I, Lehmkuhl G: The changing prevalence of attention-deficit/hyperactivity disorder and methylphenidate prescriptions: A study of data from a random sample of insureds of the AOK health insurance company in the German state of Hesse, 2000–2007. *Dtsch Arztebl Int* 2010; 107(36): 615–21.

Dr. rer. soc. Ingrid Schubert

PMV forschungsgruppe an der Klinik und Poliklinik für Psychiatrie und Psychotherapie des Kindes- und Jugendalters
Universität zu Köln
Herderstraße 52
50932 Köln
Ingrid.Schubert@uk-koeln.de

Conflict of interest statement

PMV research group has received project funding from statutory sickness funds, foundations, and pharmaceutical companies (Sanofi-Aventis, Novo-Nordisk, Lilly, Sanofi-Pasteur-MSD, Bayer-Schering, Merz, Janssen-Cilag, Abbott). Dr Ingrid Schubert and Ingrid Köster have not received personal honoraria. Professor Dr Gerd Lehmkuhl has received research funding from Lilly Germany and Bristol-Myers Squibb; he is a member of the advisory board of Eli Lilly Strattera.