Published in final edited form as:

Int J Public Health. 2011 April ; 56(2): 231–235. doi:10.1007/s00038-010-0183-y.

Hypertension and risk of death from external causes in the physicians' health study enrollment cohort

Paul D. Terry¹, Robert J. Glynn², Julie E. Buring², and Howard D. Sesso²

¹ Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA, 30322

² Division of Preventive Medicine, Department of Medicine, Brigham and Women's Hospital, Boston, MA 02215

Abstract

Objectives—To address the recent hypothesis that hypertension increases the risk of death from external causes.

Methods—We examined blood pressure and death from external causes among 82,037 male physicians who were screened for eligibility to enroll in the Physicians' Health Study.

Results—During up to 6.6 years of mortality follow-up, there were 304 deaths from external causes. No association was found overall, although we observed an increased risk of non-passive external causes of death, particularly death due to falls, which was independent of various lifestyle, medical, and pharmacological risk factors.

Conclusions—The results of our study support this novel hypothesis. Further studies are needed to explore potential causal mechanisms between elevated BP and the risk of external death.

Keywords

injuries; external causes of death; epidemiologic studies; hypertension; mortality

Introduction

The possibility that hypertension increases the risk of death from external causes was supported by a recent study that used 25 years of mortality follow-up data from the large cohort of men screened for MRFIT (1). That study showed a 40–50 percent increased risk of death from external causes, particularly injuries, in men with Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7) (2) Stage 2 hypertension, consistent with the results of three (3–5) but not another (6) of four smaller studies. These findings are also consistent with data showing that hypertension-related cerebrovascular damage (7–9) may lead to subtle cognitive and/or functional changes (8,10–15) that may predispose individuals to falls or other injuries (12,16). Nonetheless, epidemiological studies of this issue remain few, and control for potentially confounding variables, such as socioeconomic status (SES), alcohol consumption, physical activity, and use of various medications, has been limited (1). Given the potential public health significance of this issue, we sought to clarify the association between hypertension and risk

Correspondence to: Paul D. Terry, Ph.D., M.P.H., Department of Epidemiology, Emory University School of Public Health, 1518 Clifton Road, NE, Atlanta, GA 30322, Phone: (404) 727-8715, Fax: (404) 727-8737, pdterry@sph.emory.edu.

Competing Interests: The authors declare that they have no competing interests.

of death from external causes using mortality follow-up data from the Physicians' Health Study Enrollment Cohort.

Methods

The Physicians' Health Study was a randomized, double-blind, placebo-controlled trial with a 2×2 factorial design testing two primary prevention hypotheses: 1) whether 325 mg of aspirin taken every other day reduces CVD mortality (17) and 2) whether 50 mg of beta-carotene taken on alternate days decreases the incidence of cancer (18). Starting in 1982, letters of invitation, informed consent forms and baseline questionnaires were mailed to 261,248 potentially eligible U.S. male physicians on the American Medical Association mailing list. By 2/27/1984, 112,528 physicians had responded to the initial enrollment questionnaire. These analyses are confined to the 82,037 respondents aged 39–85 years who at baseline provided self-reported blood pressure measurements and had no prior history of myocardial infarction (MI), stroke, cancer, TIA, peptic ulcers, or liver disease.

On the baseline questionnaire, physicians reported their cardiovascular risk factors including age, height (in inches), weight (in pounds), history of treatment for hypertension (never, past, or current) or hypercholesterolemia (never, past, or current), serum cholesterol (mg/dL), cigarette smoking (never, past, current), frequency of vigorous exercise (<1/week or ≥1/week), alcohol use (rarely/never, <1 per week, ≥1 per week), history of diabetes mellitus (yes/no), current use of aspirin (≥1 per week) and current use of multivitamins (≥1 per week). Body-mass index (in kg/m²) was calculated using weight and height. Self-reported systolic blood pressure (SBP) and diastolic blood pressure (DBP) were collected, which have been highly correlated with measured SBP ($r=0.72$) and DBP ($r=0.60$) in a sample of physicians.(19) Individuals were placed in one of the following JNC 7 categories irrespective of treatment status: normal < 120 mmHg systolic and < 80 mmHg diastolic; pre-hypertension = 120 to 139 mmHg systolic or 80 to 89 mmHg diastolic; stage 1 hypertension = 140 to 159 mmHg systolic or 90 to 99 mmHg diastolic; stage 2 hypertension = 160+ mmHg systolic or 100+ mmHg diastolic.

Our primary endpoint was external cause of death. Death certificates were obtained for the respondents who died before Feb 01, 1988, using the National Death Index. The deaths were classified by trained nosologists using the first revision of the *Ninth International Classification of Diseases* (ICD-9) in conjunction with the *Automated Classification of Medical Entities Decision Tables* to manually select underlying cause of death. Deaths from external causes (ICD-9 E800–E 999) were further categorized as external causes deaths not due to suicide or homicide), i.e., those resulting from unintentional injuries. We also categorized “non-passive” deaths by further excluding unintentional injury deaths that were not likely due to actions taken by the victim, for example, death resulting from being a passenger in a fatal motor vehicle accident, death from falling objects, death during surgery, or death from reaction to therapeutic medication. We did not include the following causes of death in analyses of “non-passive” deaths: E800–E803, E804, E805, E806, E807, E810–E825, E837, E850–E879, and E890–E899, E900–E915, E916, E917.1, E929.4, E929.5, E930–E949, E980–E999.

Means and standard deviations (or percentages, when appropriate) were tabulated for the following individual risk factors according to cause of death: SBP, DBP, age, body mass index, current cigarette smoking, current use of aspirin, vigorous physical activity, alcohol consumption, serum cholesterol, presence of hypertension, and history of the following: diabetes, angina, TIA, renal disease, high cholesterol, and use of medicine for hypertension or high cholesterol. Cox proportional hazards analyses(20) were used to model the associations of blood pressure with all external causes of death (injuries, suicides, and

homicides) and individual causes. Hazard ratios (HRs) and 95% confidence intervals (CIs) are cited.

Hypertension risk categories were based on the classifications outlined by the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7) (2), as was described in detail above.

The multivariate model adjusted for age, cigarette smoking (current, former, never), physical activity (four categories), body mass index, alcohol consumption, aspirin use (yes or no), and history of the following (yes or no): diabetes, renal disease, drug treatment for hypertension, and drug treatment for high cholesterol. We also examined whether there was a non-linear for the categorical SBP and DBP models (separately) in the multivariable model, in which we compared models with just the linear trend term with the model with both the linear trend + quadratic (squared) linear trend term, and the -2 Log Likelihood calculated. All analyses were conducted using SAS 9.1 (SAS Institute Inc, Cary, NC) software. All p values were from two-sided tests.

Results

The average duration of follow-up (from receipt of the PHS enrollment questionnaire through death or January 31, 1988) for all 82,037 men was 5.4 years (a total of 446,641 person-years), and included 304 deaths from external causes (176 injuries, 119 suicides, and 9 homicides). Of the 176 external causes of death not due to suicide or homicide, 122 were considered as not “passive,” e.g., they were drivers and not merely passengers in fatal automobile accidents.

Mean baseline blood pressure (BP) levels and the prevalence of hypertension were greater for men who died from external deaths and subcategories of these deaths than for the cohort as a whole (Table 1). Men who died of external causes were also more likely to smoke cigarettes at baseline, have a sedentary lifestyle, have a history of diabetes, and have a history of transient ischemic attacks.

Hypertension status was not associated with risk of death from combined external causes (Table 2). Compared to individuals who had “normal” BP levels according to JNC 7, the multivariate-adjusted HRs (95% CIs) for pre-hypertension, stage 1 hypertension, and stage 2 hypertension were 0.99 (0.68, 1.45), 1.03 (0.66, 1.62), and 1.24 (0.62, 2.49), respectively. When all external deaths not due to suicide or homicide were separated and examined, the multivariate-adjusted HRs (95% CIs) for pre-hypertension, stage 1 hypertension, and stage 2 hypertension were 0.96 (0.57, 1.61), 1.08 (0.59, 1.96), and 1.81 (0.81, 4.06), respectively. When we examined only “non-passive” deaths, the multivariate-adjusted HR and 95% CIs for pre-hypertension, stage 1 hypertension, and stage 2 hypertension were 1.36(0.69–2.66), 1.64(0.76–3.54), and 4.10 (1.56–10.8), respectively. The latter statistically significant finding suggests the possibility of a positive association between hypertension and risk of death from non-passive unintentional injuries. Other statistically significant inverse predictors of non-passive deaths included age and physical activity level.

Although the number of deaths was too low to examine subcategories of non-passive deaths reliably, our results suggest that the association of hypertension status with death from falls may be particularly strong. The age-adjusted HR (95% CI) of falls for men with stage 2 hypertension (vs. normal) was 11.3 (95% CI, 1.28, 99.4; $p=0.03$), although this was based in very small numbers. The multivariate-adjusted HR for falls was similar (data not shown).

Based on the above results, we decided to further examine the relationship of BP levels with “non-passive” deaths. For these 122 deaths, there was a positive association between each

increasing 10 mmHg category of SBP and DBP and an increased risk of death. The multivariate-adjusted HRs (95% CIs) for men with SBP levels of 110–119 mmHg and each successive 10 mmHg category compared with men with SBP <110 mmHg (the referent category, limited to 2 deaths) were, respectively, 0.87(0.19–3.95), 1.01(0.24–4.26), 1.46(0.35–6.14), 1.57(0.36–6.89), 2.15(0.44–10.6), 3.90(0.74–20.5), and 5.12(0.81–32.4). The test for linear trend was statistically significant ($p=0.001$). Analyses for DBP in increasing 5 mmHg increments up to 100 mmHg compared with the referent of DBP <70 mmHg showed a similar tendency towards an increasing risk of death, although the linear trend ($p=0.24$) was not monotonic among the middle DBP categories. Results were similar when analyses were restricted to participants without history of antihypertensive medication use (data not shown). No significant non-linear associations were observed in separate models for either SBP or DBP.

Discussion

In this cohort of initially healthy, middle-aged and older male physicians, we found an increased risk of external causes of death not due to suicide or homicide among men with high BP, particularly among those with JNC 7 Stage 2 hypertension. However, our results were statistically significant only for those deaths after exclusion of deaths that were classified as being “passive.” This study represents one of the larger cohort studies of BP and external causes of death to date, although a maximum follow-up of 6.6 years resulted in a smaller number of deaths than the MRFIT cohort that benefited from 25 years of follow-up. The shorter follow-up likely helped to reduce misclassification by BP and potential confounding variables because these risk factors (including prescribed antihypertensive medicine formulations) tend to change over time. Notably, the strongest association between BP and risk of external causes of death was observed in MRFIT during the first 10 years of follow-up (1).

A notable strength of this study was the ability to evaluate the potential role of confounding, including that by lifestyle factors (smoking, alcohol consumption, and physical activity level), history of taking aspirin or medications for BP, high cholesterol and diabetes, cholesterol level, and other related factors. Because all study participants were practicing physicians, adjustment for education or SES variables was not necessary. These results suggest that confounding likely does not explain the positive association suggested in our data or that shown in MRFIT and some (3–5), but not all (6) previous studies.

Hypertension may lead to cerebrovascular damage and consequent cognitive or functional decline (7–9), which then increases the risk of death from external causes. Hypertension-related pathological changes in the brain and its vasculature include vascular remodeling, impaired cerebral autoregulation, cerebral microbleeds, white matter lesions, unrecognized lacunar infarcts, and Alzheimer-like changes such as amyloid angiopathy and cerebral atrophy (7–9). These changes, particularly in white matter lesions, have been associated with declines in frontal lobe integrity and executive function (8,10,11), attention and reaction time (12,13), lower extremity mobility and balance (14,15), and, perhaps as a consequence, with potentially fatal injuries, including those from motor vehicle accidents (12) and falls (16).

In conclusion, our study supports the findings from most (1,3–5), but not all (6), studies that examined the roles of BP and hypertension status on external causes of death. We found the strongest positive association among men with JNC 7 Stage 2 hypertension and their risk of “non-passive” deaths, and it was the only statistically significant association. As with MRFIT (1), we observed the strongest magnitude of effect between hypertension and death from falls. Though our conclusions for all external causes of death and its subcategories

must be tempered by relatively small case counts, unlike previous studies we comprehensively adjusted for potential confounders. Further studies are needed to explore potential causal mechanisms between elevated BP and the risk of external death.

Acknowledgments

Supported by research grants CA-097193, HL-26490, HL-34595, HL-42441, CA-34944, and CA-40360 from the National Institutes of Health, Bethesda, MD.

References

1. Terry PD, Abramson JL, Neaton JD. Blood pressure and risk of death from external causes among men screened for the Multiple Risk Factor Intervention Trial. *Am J Epidemiol.* 2007; 165(3):294–301. [PubMed: 17098820]
2. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA.* 2003; 289(19):2560–72. [PubMed: 12748199]
3. Tamosiunas A, Reklaitiene R, Radisauskas R, Jureniene K. Prognosis of risk factors and trends in mortality from external causes among middle-aged men in Lithuania. *Scand J Public Health.* 2005; 33(3):190–6. [PubMed: 16040459]
4. Vartiainen E, Puska P, Pekkanen J, Tuomilehto J, Lonnqvist J, Ehnholm C. Serum cholesterol concentration and mortality from accidents, suicide, and other violent causes. *BMJ.* 1994; 309(6952):445–7. [PubMed: 7920128]
5. Farchi G, Menotti A, Conti S. Coronary risk factors and survival probability from coronary and other causes of death. *Am J Epidemiol.* 1987; 126(3):400–8. [PubMed: 3618576]
6. Strandberg TE, Salomaa VV, Vanhanen HT, Naukkarinen VA, Sarna SJ, Miettinen TA. Mortality in participants and non-participants of a multifactorial prevention study of cardiovascular diseases: a 28 year follow up of the Helsinki Businessmen Study. *Br Heart J.* 1995; 74(4):449–54. [PubMed: 7488463]
7. Manolio TA, Olson J, Longstreth WT. Hypertension and cognitive function: pathophysiologic effects of hypertension on the brain. *Curr Hypertens Rep.* 2003; 5(3):255–61. [PubMed: 12724059]
8. Raz N, Rodrigue KM, Acker JD. Hypertension and the brain: vulnerability of the prefrontal regions and executive functions. *Behav Neurosci.* 2003; 117(6):1169–80. [PubMed: 14674838]
9. DeCarli C, Miller BL, Swan GE, Reed T, Wolf PA, Garner J, et al. Predictors of brain morphology for the men of the NHLBI twin study. *Stroke.* 1999; 30(3):529–36. [PubMed: 10066847]
10. Tullberg M, Fletcher E, DeCarli C, Mungas D, Reed BR, Harvey DJ, et al. White matter lesions impair frontal lobe function regardless of their location. *Neurology.* 2004; 63(2):246–53. [PubMed: 15277616]
11. Prins ND, van Dijk EJ, den Heijer T, Vermeer SE, Jolles J, Koudstaal PJ, et al. Cerebral small-vessel disease and decline in information processing speed, executive function and memory. *Brain.* 2005; 128(Pt 9):2034–41. [PubMed: 15947059]
12. Anstey KJ, Wood J, Lord S, Walker JG. Cognitive, sensory and physical factors enabling driving safety in older adults. *Clin Psychol Rev.* 2005; 25(1):45–65. [PubMed: 15596080]
13. Sierra C, De La Sierra A, Salamero M, Sobrino J, Gomez-Angelats E, Coca A. Silent cerebral white matter lesions and cognitive function in middle-aged essential hypertensive patients. *Am J Hypertens.* 2004; 17(6):529–34. [PubMed: 15177527]
14. Onen F, Feugeas MC, Baron G, De Marco G, Godon-Hardy S, Peretti II, et al. Leukoaraiosis and mobility decline: a high resolution magnetic resonance imaging study in older people with mild cognitive impairment. *Neurosci Lett.* 2004; 355(3):185–8. [PubMed: 14732462]
15. Starr JM, Leaper SA, Murray AD, Lemmon HA, Staff RT, Deary IJ, et al. Brain white matter lesions detected by magnetic resonance [correction of resosnance] imaging are associated with balance and gait speed. *J Neurol Neurosurg Psychiatry.* 2003; 74(1):94–8. [PubMed: 12486275]
16. Kuo HK, Lipsitz LA. Cerebral white matter changes and geriatric syndromes: is there a link? *J Gerontol A Biol Sci Med Sci.* 2004; 59(8):818–26. [PubMed: 15345732]

17. Steering Committee of the Physicians' Health Study Research Group. Final report on the aspirin component of the ongoing Physicians' Health Study. *N Engl J Med.* 1989; 321(3):129–35. [PubMed: 2664509]
18. Hennekens CH, Buring JE, Manson JE, Stampfer M, Rosner B, Cook NR, et al. Lack of effect of long-term supplementation with beta carotene on the incidence of malignant neoplasms and cardiovascular disease. *N Engl J Med.* 1996; 334(18):1145–9. [PubMed: 8602179]
19. Klag MJ, He J, Mead LA, Ford DE, Pearson TA, Levine DM. Validity of physicians' self-reports of cardiovascular disease risk factors. *Ann Epidemiol.* 1993; 3(4):442–7. [PubMed: 8275223]
20. Cox D. Regression models and life tables with discussion. *J R Stat Soc Ser B.* 1972; 34:187–2.

Table 1
Self-Reported Baseline Characteristics of Men in the Physician's Health Study Enrollment Cohort (USA, 1982–1984) According to Cause of Death

	All External Deaths [*]	All Injury Deaths ^{**}	Injury deaths excluding "passive" deaths ^{***}	All Deaths	All Men
SBP (mean mmHg)	130.4	131.8	133.2	135.1	127.6
DBP (mean mmHg)	80.2	80.9	81.6	80.9	79.3
SBP $\geq$ 140 or DBP $\geq$ 90 (%)	31.6	35.2	36.9	44.1	24.2
Age at recruitment (mean years)	58.5	59.7	58.6	65.9	55.3
Body mass index (mean)	25.3	25.4	25.6	25.0	25.0
Any history of hypertension treatment (%)	25.9	28.7	25.0	37.0	18.9
Any history of high cholesterol (%)	13.1	14.2	11.1	16.0	14.6
Any history of high cholesterol treatment	5.3	6.4	3.2	4.1	2.5
Total cholesterol (mean mg/dl)	202.5	203.9	204.4	209.2	211.4
Current smoker (%)	21.8	20.6	17.2	19.3	12.2
Vigorous physical activity <1 time/week (%)	36.5	36.4	36.9	43.6	29.4
History of diabetes (%)	4.9	5.1	2.5	11.1	3.3
History of angina (%)	3.3	2.8	2.5	7.8	3.1
History of transient ischemic attack (%)	3.6	3.5	1.7	6.2	1.5
History of renal disease (%)	1.0	1.7	0.8	3.2	0.9
Current regular use of aspirin	36.0	31.4	27.7	37.3	29.3
No consumption of alcohol (%)	18.2	17.5	17.5	22.1	17.2
Number of Men	304	176	122	3,451	82,037

* All external causes of death, i.e., any death that is other than from natural causes (ICD-9 E800–E 999)

** All external causes of death excluding suicide homicide (E950 – E969)

*** All external causes of death excluding deaths that are not likely due to actions taken by the victim, for example, death resulting from being a passenger in a fatal motor vehicle accident, death from falling objects, death during surgery, death from reaction to therapeutic medication (E800–E803, E804, E805, E806, E807, E810–E825, E837, E850–E879, and E890–E899, E900–E915, E916, E917.1, E929.4, E929.5, E930–E949, E980–E999)

Table 2

Hazard Ratios (95% CIs) of External Causes of Death by Joint National Committee 7 (JNC 7) Blood Pressure Classification (USA, 1982–1988).

Cause of Death	JNC 7 Blood Pressure Classification*			
	Normal (n=12,432)	Pre-Hypertension (n=49,772)	Stage 1 Hypertension (n=17,446)	Stage 2 Hypertension (n=2,387)
All External Causes	39**	169	81	15
- Age-adjusted	1.00	1.00 (0.71–1.42)	1.18 (0.79–1.75)	1.45 (0.78–2.69)
- Multivariate***	1.00	0.99 (0.68–1.45)	1.03 (0.66–1.62)	1.24 (0.62–2.49)
All Injuries	21	93	50	12
- Age-adjusted	1.00	1.00 (0.62–1.61)	1.26 (0.74–2.13)	1.93 (0.92–4.04)
- Multivariate***	1.00	0.96 (0.57–1.61)	1.08 (0.59–1.96)	1.81 (0.81–4.06)
Injuries excluding “passive” injuries	11	66	35	10
- Age-adjusted	1.00	1.41 (0.74–2.68)	1.90 (0.95–3.81)	3.69 (1.51–8.99)
- Multivariate***	1.00	1.36 (0.69–2.66)	1.64 (0.76–3.54)	4.10 (1.56–10.8)

* Individuals were placed in one of the following JNC 7 categories: normal < 120 mmHg systolic and < 80 mmHg diastolic; pre-hypertension = 120 to 139 mmHg systolic or 80 to 89 mmHg diastolic; stage 1 hypertension = 140 to 159 mmHg systolic or 90 to 99 mmHg diastolic; stage 2 hypertension = 160+ mmHg systolic or 100+ mmHg diastolic.

** Number of deaths

*** Model includes covariates corresponding to age, smoking category, vigorous physical activity, body mass index, use of aspirin, use of alcohol, use of medicine for hypertension, use of medicine for high cholesterol, history of diabetes, and history of renal disease.