controversial topics in surgery

Robotic surgery

ADAM JONES AND KRISHNA SETHIA

These two papers succinctly express the core issues of the 'robot debate'. Jones demonstrates the rapid skill acquisition of urologists who can now bypass the laparoscopic stage entirely. Sethia illustrates the huge cost implications, lack of evidence and limited use of the robot.

The very handsome robot in my hospital lies dormant for 3 days or more per week: the predicted use in gynaecological and colorectal surgery remains unmet. Three DGHs and one teaching hospital in a 25-mile radius have these machines: would not one suffice

until sufficient data confirm the health economic as well as clinical benefits?

There is anecdotal evidence that referral rate is falling for prostatic brachytherapy in some robot-enhanced institutions – clinically justified or gizmo fuelled? Apparently, in Belgium, 40% of robots are in 'resting mode'. Is the robot like Concorde: an enormous technological masterpiece, hugely expensive but of benefit only to a few.

TOM DEHN

Why we have bought a robot

ADAM JONES

Urology Department, Royal Berkshire Hospital, Reading, UK

CORRESPONDENCE TO: Adam Jones, Urology Department, Level 1 Centre Block, Royal Berkshire NHS Foundation Trust, London Road, Reading RG1 5AN, UK
E: adam.jones@royalberkshire.nhs.uk

Passing fancy or legitimate treatment option?¹ No, this is not robotic surgery, but the debate that took place with regards to the radically new technique – laparoscopic cholecystectomy – in the early 1990s. No one would argue for open rather than laparoscopic cholecystectomy now but the debate in urology between open and laparoscopic prostatectomy is harder to decide. The subtle variation in both patients and cancer characteristics, the inherent bias in peri-operative care between different centres and the undoubted change in patient management by 'open' surgeons in response to the 'threat' of laparoscopic radical prostatectomy (LRP) makes comparison difficult.²

Most literature reviewed suggests there are small advantages to the laparoscopic approach with regards to blood loss, peri-operative pain and length of stay whereas potency, continence and positive margins are more equal.³ It appears that results are probably more surgeon-dependent than technique-dependent. Despite these similarities, the

whole of surgery is moving to less and less invasive techniques and the next generation of surgeons will not choose to train in open rather than laparoscopic surgery. Given this fact, should the approach be traditional laparoscopy or robot-assisted laparoscopy.

World-wide, and especially in the UK, urology has led the way for robotic surgery, in particular robot-assisted laparoscopic prostatectomy (RALP). Comparisons of RALP with open prostatectomy face the same problems and show similar results to the LRP/open prostatectomy debate.^{4,5} The robotic community does itself no favours with the commercial marketing approach often adopted. The fact that RALP has achieved parity with the open and laparoscopic techniques (refined to their maximum over many years) in such a short time is testament to its value and predicts the results will continue to improve.⁶

The number of units world-wide with a surgical robot has increased exponentially over the last 5 years.

Considering the expense of a robotic programme why is this? First, there is a competitive element. This is not just about the perception that your hospital is as cutting edge (or more than) than your neighbours but also the necessity to maintain your catchment population. Second, whilst skilled laparoscopic surgeons argue, probably correctly, that they can do just as good a job without the expense of a robot, the number that reaches this standard is limited. RALP is difficult, LRP is very difficult. The British Association of Urological Surgeons (BAUS) cancer registry shows only four surgeons with greater than 20 cases per year. Whilst there are a few gifted surgeons who have mastered LRP, there is a long learning curve.⁷ Even with formal fellowships, it is difficult.⁸

When counselling patients, it is the results that individual surgeons can achieve that are important not the often quoted excellent results of a small number of world centres of excellence. Excluding a handful of surgeons with very large series of LRP, most laparoscopic surgeons that have performed both LRP and RALP rapidly find RALP easier. This translates to a shorter learning curve. Hakimi *et al.*⁹ found superior results with RALP compared to LRP with the first 75 robotic cases compared to the last 75 LRP cases in a series of over 300. This number is significant because it is achievable by many urologists in the UK in a short time whereas 300 cases is much harder. Other shortened learning curves for RALP have been reported,¹⁰ allowing for greater trainee participation¹¹ and even allowing open surgeons with no laparoscopic experience to offer a laparoscopic option.¹²

All of these mean that more patients have the ability to benefit from a laparoscopic approach with all its generic advantages that was previously available by standard LRP.

The costs of a robotic programme are often cited as a criticism. At around £1.3 million, this is a significant capital purchase; most centres have used a combination of charitable donations, research programmes and private revenue to offset this. The running costs for the disposables is about to be eliminated by an uplift in robotic HRG codes – a reflection in itself that robotic surgery is here to stay. Maintenance costs are around £70,000 a year. As robotic use crosses from urology to gynaecology and colorectal surgery, as it undoubtedly will, this cost falls to just a few hundred pounds per patient. This is offset by the reduced length of

stay and is a fraction of the cost of many NHS treatments. Only time will tell whether those with robots have egg on their face or whether critics will be regarded as latter day Luddites.

References

1. Miller TA. Laparoscopic cholecystectomy: passing fancy or legitimate treatment option? *Gastroenterology* 1990; **99**: 1527–9.
2. Eggener SE, Guillionneau B. Laparoscopic radical prostatectomy: ten years later, time for evidence-based foundation. *Eur Urol* 2008; **54**: 4–7.
3. Martinez-Salamanca JI, Romero Otero J. Critical analysis between open, laparoscopic and robotic prostatectomy: perioperative morbidity and oncological results (part 1). *Arch Esp Urol* 2007; **60**: 755–65.
4. Krambeck AE, DiMarco DS, Rangel LJ, Bergstralh EJ, Myers RP, Blute ML *et al.* Radical prostatectomy for prostatic adenocarcinoma: a matched comparison of open retropubic and robot-assisted techniques. *BJU Int* 2008; **103**: 448–53.
5. Ficarra V, Novara G, Artibani W, Cestari A, Galfano A, Graefen M *et al.* Retropubic, laparoscopic, and robot-assisted radical prostatectomy: A systemic review and cumulative analysis of comparative studies. *Eur Urol* 2009; **55**: 1037–63.
6. Goldstraw MA, Patil K, Anderson C, Dasgupta P, Kirby RS. A selected review and personal experience with robotic prostatectomy: implications for adoption of this new technology in the United Kingdom. *Prostate Cancer Prostate Dis* 2007; **10**: 242–9.
7. Eden CG, Neill MG, Louie-Johnsun MW. The first 1000 cases of laparoscopic radical prostatectomy in the UK: evidence of multiple 'learning curves'. *BJU Int* 2009; **103**: 1224–30.
8. Brown JA, Sajadi KP. Laparoscopic radical prostatectomy: six months of fellowship training doesn't prevent the learning curve when incorporating into a lower volume practice. *Urol Oncol* 2009; **27**: 144–8.
9. Hakimi AA, Blitstein J, Feder M, Shapiro E, Ghavamian R. Direct comparison of surgical and functional outcomes of robot-assisted versus pure laparoscopic radical prostatectomy: single-surgeon experience. *Urology* 2009; **73**: 119–23.
10. Hakimi AA, Feder M, Ghavamian R. Minimally invasive approaches to prostate cancer: a review of the current literature. *J Urol* 2007; **4**: 130–7.
11. Schroeck FR, de Sousa CA, Kalman RA, Kalia MS, Pierre SA, Haleblan GE *et al.* Trainees do not negatively impact the institutional learning curve for robotic prostatectomy as characterised by operative time, estimated blood loss, and positive surgical margin rate. *Urology* 2008; **71**: 597–601.
12. Ahlering TE, Skarecky D, Lee D, Clayman RV. Successful transfer of open surgical skills to a laparoscopic environment using a robotic interface: initial experience with laparoscopic radical prostatectomy. *J Urol* 2003; **170**: 1738–41.

Why I do not have a robot

KRISHNA SETHIA

Norfolk & Norwich University Trust, Norwich, UK

CORRESPONDENCE TO:

Krishna Sethia, Medical Director, Norfolk & Norwich University Trust, Colney, Norwich NR4 7UY, UK

E: krishna.sethia@nnuh.nhs.uk

The way in which surgeons introduced laparoscopic surgical techniques in the 1990s could be described as an unfortunate illustration of the value which they place on evidence-based medicine. Despite the lack of scientific rigour at the time, we are now at a stage where most intra-abdominal procedures can be performed in a minimally-invasive fashion and, as a result of a few randomised controlled trials performed as an afterthought, there is some evidence of patient advantage. This is largely in the form of reduced discomfort and faster recovery times; surgeons now accept these benefits as justification for the morbidity and, in some cases, mortality of their original uncontrolled activities. There is little information as to whether patients agree.

During the 1990s whilst surgeons were increasing their experience of laparoscopic techniques, robotic technology was developed. By the latter part of the decade, robotically-assisted procedures were attempted in various specialties, most notably cardiac surgery. The suggested advantages of this technique over conventional laparoscopy were, and remain, that it offered better magnification and hence visualisation, increased dexterity with reduction of tremor and that there was a shorter learning curve for the surgeon. The advantages for the patient were, and remain, less obvious and once again there is a lack of objective scientific evidence comparing outcomes after robotic surgery with those of previously-established techniques.

The majority of the available outcome data are for radical prostatectomy, by far the commonest robotically-assisted procedure performed so far. Despite the fact that, to date, 75% of the published series of robotically-assisted radical prostatectomy are from only eight institutions, all therefore treating high volumes of patients, there remains no evidence of superior outcomes of robotic surgery. The margin-positive, PSA recurrence, incontinence and postoperative erectile dysfunction rates are, in fact, similar for robotically-assisted, laparoscopic and, indeed, open operation. Whilst the robotic and laparoscopy approaches result in shorter hospital stays and lower blood loss than open radical prostatectomy, these differences are rarely clinically significant.¹ Given that there is sound evidence that the results of radical prostatectomy are related to patient volumes and that there is a significant learning curve to any new procedure, this must raise the question of whether new centres for robotic surgery will only add to morbidity for some

patients in the short-term for no substantial clinical gain for the majority in the long-term. There are thus clearly significant ethical issues for the surgeon contemplating starting robotic procedures. Justifying innovation and the risks to which it exposes individual patients may be easy if the new procedure is thought to promise tangible benefit. It is much less straightforward when, as is the case with the robot, there is ample evidence that the benefit is marginal.

As well as ethical issues there are proven disadvantages in using the robot – the most obvious of which is its cost. The General Medical Council guidance on *Good Medical Practice*² dictates that doctors have a responsibility to ensure proper use of resource and, in the current world economic climate, this is increasingly relevant. Evidence from a large volume centre in the US shows that, when the effect of the capital cost is included, a robotic is \$1239 more expensive than a laparoscopic prostatectomy which, in turn, is \$487 costlier than an open procedure.³ Although the past few years have seen generous increases in NHS expenditure, we know that resources will now be severely limited for many years. At the time of writing, the capital cost of a da Vinci robot system approaches £1 million and a cancer centre for a population of 1–1.5 million would expect to perform 100–150 radical prostatectomies per year. Even if demand for surgery for prostate cancer increases, the servicing and disposable costs per case for a unit performing 150 prostatectomies per year would be £1500–2000 more than the cost of open surgery and at least £500 more than that of laparoscopic prostatectomy. Compared to laparoscopic prostatectomy there is no significant length of stay or theatre time advantage to compensate for these high costs. Clearly, unless the robot has been donated to a trust, it is impossible to recover the true cost by performing prostatectomies alone. Whilst there may be a place to expand the use the robot in cardiac, paediatric and gynaecological surgery and thus reduce cost per case, these roles are as yet ill-defined. The robot can only begin to be financially-viable in the largest of centres performing several hundred procedures per year. To achieve this volume in the UK would require further significant centralisation and commissioners of healthcare would need to decide whether they can define any advantages of robotic surgery which would justify requiring all their suitable patients to travel significant distances for their surgery.

Despite the adverse economic consequences of robotic surgery, the public perceive it to be a superior technique and it has undoubtedly captured their imagination. However, there is now evidence that it is associated with significantly reduced patient satisfaction. For example, Schroeck *et al.*⁴ showed that patients undergoing robotic prostatectomy were more likely to regret their selection of treatment than were those undergoing open operation with 24% unhappy with the robotic procedure. The major cause of dissatisfaction was thought to be their high expectation of the use of new technology, a fact which surgeons must be aware of when counselling patients about treatment options.

In a recent editorial, Leff and Finucane⁵ described the driving forces which lead clinicians into a culture of 'gizmo idolatry'. They illustrate the potential harm that may result from 'credulous acceptance' of new techniques using the example of the frontal lobotomy. Although this became a procedure which was rapidly accepted by the medical community in the 1930s and 1940s, and won Egas Moniz the 1949 Nobel Prize for medicine, this happened with little evidence of any true benefit and, as subsequently became clear, severe and lasting harm to many patients. The proliferation of robots

in UK hospitals in the past 12 months shows how urologists in particular continue to embrace new technology which has little proven patient advantage. Whilst the robot appears safe (in experienced hands at least), there can be little doubt that it depletes the NHS budget with, in many hospitals, no realistic chance of recovering the investment. Clinicians and managers should consider carefully whether purchasing this gizmo constitutes responsible use of scarce NHS resource.

References

1. Frota R, Turna B, Barros R, Gill I. Comparison of radical prostatectomy techniques: open, laparoscopic and robotic assisted. *Int Braz J Urol* 2008; **34**: 259–69.
2. General Medical Council. Good Medical Practice. London: GMC, 2006; 8.
3. Lotan Y, Cadeddu JA, Gettman MT. The new economics of radical prostatectomy: cost comparison of open, laparoscopic and robot assisted techniques. *J Urol* 2004; **172**: 1431–5.
4. Schroeck FR, Krupski TL, Sun L, Albala DM, Price MM, Polascik TJ *et al.* Satisfaction and regret after open retropubic or robotic-assisted laparoscopic radical prostatectomy. *Eur Urol* 2008; **54**: 785–93.
5. Leff B, Finucane T. Gizmo idolatry. *JAMA* 2008; **299**: 1830–2.