TECHNIQUE

Standard mass closure of the wound is achieved from the extremities towards the centre with two continuous, looped 1-nylon sutures. A probe is passed between adjacent throws of suture 2–3 cm from one end of the wound, along its underside, to emerge through the middle. The closure is then completed and the two sutures tied to each other, with the probe *in situ*. Several further throws are then placed at intervals, to create a loose ‘chain’ of three or four knots about 10 cm in length (Fig. 1). This is fed through the eye of the probe, so that at least two knots have been pulled through (Fig. 2). The probe is then briskly retrieved back through the corner of the closure, taking the suture and knot with it into the abdomen (Fig. 3). The suture is then trimmed at its exit from the abdomen, leaving no palpable nylon in the wound.

DISCUSSION

This technique guarantees that the knot always lies intra-abdominally. This alleviates future anxiety in this group of patients who could otherwise mistake an unburied suture for disease recurrence. It also eliminates the discomfort that could otherwise be associated with a palpable knot.

A useful technique for the control of bleeding following peripheral vascular injury

WILL BJ RUDGE¹, BEN CJ RUDGE², CHRIS J RUDGE³

¹Department of Orthopaedics, St Helier Hospital, Carshalton, UK

²Department of Orthopaedics, Royal National Orthopaedic Hospital, Stanmore, UK

³National Clinical Director for Transplantation, Department of Health, London, UK

CORRESPONDENCE TO

Will Rudge, E: willrudge@doctors.org.uk

BACKGROUND

The majority of arterial bleeding from the peripheral vasculature can be controlled with relative ease by direct pressure. The most effective way of applying pressure is by hand, but this is not practical for prolonged periods. A traditional pressure dressing of gauze and bandage is often ineffective in providing point pressure and may result in a tourniquet effect as bandages are repeatedly tightened. We describe a simple dressing technique for providing localised pressure, using equipment readily available in any ward or emergency department (Fig. 1).

TECHNIQUE

Initially, a small non-adherent dressing is placed over the site of bleeding. Then the lid of a small specimen pot is very firmly packed with cotton wool and positioned directly over the site where pressure is required (Fig. 2). This is secured in place using

Figure 1 Equipment required.

Figure 2 Positioning of a small specimen pot firmly packed with cotton wool.

Figure 3 (A) Application of bandage. (B) Final result.

a bandage around the limb. During the application of the bandage, as it crosses over the lid it is twisted through 180° in order

to provide pressure specifically over the bleeding point (Fig. 3). This gives excellent point pressure exactly where required, and can control bleeding without compromising distal blood flow.

DISCUSSION

We have not found this technique previously described in the literature. It was initially developed for use in dialysis units to control excessive bleeding from arteriovenous fistulae but the authors have also used it successfully in various trauma situations, including radial and brachial artery injuries. It is effective, quick, simple to apply and safe, and provides control of bleeding until definitive intervention.

TECHNICAL TIP

A simple aid to wire localisation procedure in breast surgery: 'covering the hook-wire with a needle

KAUSIK RAY, ASHOK CHOUHAN, AKHIL JOHRI
West Sussex Breast Unit, Worthing Hospital, Worthing, UK

CORRESPONDENCE TO

Kausik Ray, West Sussex Breast Unit, Worthing Hospital, Worthing, UK
E: ray_kausik2003@yahoo.co.in

The hooked wire used for localising impalpable lesions in the breast is soft and malleable. Inadvertent cutting with diathermy

Figure 2 Position of wire tip can be assessed by moving the hub of the needle up, down or sideways.

or scissors¹ and difficulty in locating its tip may be problematic during operation. To overcome this, we use a blunt 18-G needle to cover the wire. It prevents inadvertent cutting of the wire (Fig. 1), helps in locating wire tip accurately by moving the hub of the needle (Fig. 2) and minimises breast deformity as less tissue is removed due to better identification of the position of the lesion. In our experience, this is a useful tip for breast surgeons.

Reference

1. D'Orsi CJ, Swanson RS, Moss LJ, Reale FR, Wertheimer MD. A complication involving a braided hook-wire localization device: *Radiology* 1993; **187**: 580–1.

Figure 1 Protecting needle prevents inadvertent cutting of wire.