movement. In addition, the terms used will be familiar to all radiographers and easily learnt by trauma surgeons.

AUTHORS' RESPONSE

S Chaganti, D Kumar

Department of Trauma and Orthopaedics, Royal Gwent Hospital, Newport, UK

doi 10.1308/003588410X12628812459454

CORRESPONDENCE TO

S Chaganti, E: drchaganti2002@yahoo.co.uk

We appreciate that Day also agrees there is a definite role for unambiguous language between doctors and radiographers. Unfortunately, trauma surgeons do not work with radiographers as closely as radiologists and may have to work with different radiographers with various levels of experience. We respect what appears to have developed in his department. However, that has not become wide-spread as a language and certainly was not published in the literature at the time when we submitted our work for publication to the *Annals*.

As soon as it is appreciated that a common language is required between doctors and radiographers, a shorter, user-friendly version of commands is more than likely to evolve if one is working with the same team. The language suggested in our article is just an example. We accept some of the commands are longer than desired but shorter abbreviated commands leave room for ambiguity especially if one has to work with different radiographers sometimes in the same procedure.

Indications for tonsillectomy

MOHAMMED IQBAL SYED, NATASHA AMIRARAGHI, ALUN T WILLIAMS

Royal Hospital for Sick Children, Edinburgh, UK

doi 10.1308/003588410X12628812459490

CORRESPONDENCE TO

Mohammed Iqbal Syed, E: iqbalms@hotmail.com

COMMENT ON

Toh A, Mullin A, Grainger J, Uppal H. Indications for tonsillectomy: are we documenting them? *Ann R Coll Surg Engl* 2009; **91**: 697–9

doi 10.1308/003588409X12486167521712

We published a three-cycle audit on a similar theme in 2002.¹ We looked at 181 children at a paediatric teaching hospital undergoing tonsillectomy in three cohorts to

determine adherence to the SIGN guidelines for tonsillectomy and to assess adequate record keeping. The first cohort of patients was listed for surgery prior to the publication of the Scottish Intercollegiate Guidelines Network (SIGN) guidelines; the second after. We then analysed the data for these two cohorts, presented them to the relevant clinicians, and kept a reminder poster (of the SIGN guidelines) placed in each out-patient room prior to the third cohort.

Although we found a significant improvement in record keeping (chi-squared test) in cohort 1 to 2 $P < 0.001$, and cohort 2 to 3 $P < 0.001$, we did not find a statistically significant result in adherence to the SIGN guidelines.

In our study we set a standard of 95% adherence to the SIGN guidelines, which we were successful in achieving during the third cycle. There is no specifically mentioned standard in the audit of Toh *et al.* and we note that, after their second cycle, adherence to all four SIGN recommendations was still only 44% (74% during our second cycle), despite dissemination of audit findings of the first cycle to clinical staff, SIGN guideline posters and introduction of rubber stamp; perhaps this could be improved upon further with a third cycle to meet an appropriate standard? We acknowledge that any improvement is desirable; however, one would expect better results when all other indications for tonsillectomy have been excluded and there was an additional change implemented (the rubber stamp) in comparison with our study. We emphasise that a third cycle in the audit would have contributed even further in improving clinical practice.

It is also surprising that the paper makes no reference to the National Prospective Tonsillectomy audit² or Scottish tonsillectomy audit³ that provide valuable reference points for any study on tonsillectomy.

References

1. Williams A, Lee P, Kerr A. Scottish Intercollegiate Guidelines Network (SIGN) guidelines on tonsillectomy: a three cycle audit of clinical record keeping and adherence to national guidelines. *J Laryngol Otol* 2002; **116**: 453–4.
2. Clinical Effectiveness Unit. *National Prospective Tonsillectomy Audit*. London: The Royal College of Surgeons of England, 2005.
3. Blair RL, McKerrow WS, Carter NW, Fenton A. The Scottish tonsillectomy audit. Audit Sub-Committee of the Scottish Otolaryngological Society. *J Laryngol Otol* 1996; **110** (Suppl 20): 1–25.

AUTHORS' RESPONSE

A Toh, J Grainger, H Uppal

Department of Otolaryngology, University Hospital North Staffordshire, Stoke-on-Trent, UK

doi 10.1308/003588410X12628812459535

CORRESPONDENCE TO

Alex Toh, E: alextoh@doctors.org.uk

We thank Syed and his colleagues for their letter and are